

Association of Serum Leptin Levels with Essential Hypertension Independent of Obesity

Dr. M. Subarathi, Dr. D. Gayathri Priya*, Dr. P. Deepa

Assistant professor, Department of Biochemistry, Stanley Medical College, Chennai, Tamilnadu, India

*Corresponding author: Dr. D. Gayathri Priya
DOI: [10.21276/sjams.2019.7.2.70](https://doi.org/10.21276/sjams.2019.7.2.70)

| Received: 15.02.2019 | Accepted: 24.02.2019 | Published: 28.02.2019

Abstract

Original Research Article

Essential Hypertension though being one of the major global health burden, its pathophysiology is still not established well. This is because it is a complex disorder with multiple factors contributing to its evolution. Recently, hormones like Leptin secreted by adipose tissue are gaining significance in research studies linking hypertension and obesity. So we decided to find the association of serum leptin levels with essential hypertension and whether it is independent of obesity. The study included 60 essential hypertensives in the age group of 45-60years with both genders. Anthropometric measurements and blood pressure were recorded. Body mass index were calculated. Serum leptin levels were measured from collected fasting blood sample. Descriptive statistics were derived. Binary logistic regression was done. Serum leptin levels were found to be increased in hypertensives even after adjusting for obesity with p value < 0.000

Keywords: Essential hypertension, Body mass index, Leptin, Obesity.

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

Leptin, an adipose tissue derived peptide hormone is known to regulate food intake and energy expenditure [1]. Therefore it is assumed to play a key role in determining the weight of an individual. In addition to this, many studies have shown that leptin has a direct effect on maintaining blood pressure of an individual [2]. Leptin has both pressor and depressor effects on blood pressure. Its depressor effects are believed to be due to natriuretic effect, stimulation of endothelial nitric oxide and indirectly by increasing insulin sensitivity. Its pressor effects are due to increased sympathetic nervous system activity and due to adverse shifting of pressure-natriuresis curve [3]. Although leptin has dual and opposite roles over blood pressure regulation, in order to maintain the normal vascular tone usually the pressor effects predominate over the depressor effects. Thus when there are increased levels of leptin as in leptin resistance, it results in increased activation of the renal sympathetic system coupled with decreased natriuresis [4]. Thus hyperleptinemia is likely to produce hypertension [5]. Also leptin is found to increase the production of pro-opiomelanocortin which on activation leads to synthesis of alpha melanocyte stimulating hormone. This hormone in turn activates renal sympathetic nervous system and thereby increasing blood pressure [6]. Several studies have shown the positive role of

hyperleptinemia in obesity induced hypertension [7-10]. Many other studies could not support the role of leptin in obese hypertensives [11,12]. So we wanted to conduct a study to know the relationship of leptin levels to hypertension in our population and also to know whether it was independent of obesity.

MATERIALS AND METHODS

A cross-sectional study was carried out at a government tertiary care hospital in north Tamilnadu for a period of 9 months. Ethical committee clearance was obtained from the institution. Convenient sampling was made with 60 newly diagnosed essential hypertensives with both the genders (male =36, female =24) between the age groups of 45 to 60 years. They were not started on any drugs when included in the study. Patients with secondary hypertension, diabetes mellitus, cardiovascular disease, cerebrovascular disease, thyroid disorders, renal disorders, recent drug intake for other ailments were excluded from the study.

After obtaining informed consent in local language, anthropometric measurements such as weight in kilograms and height in metres were measured and body mass index were calculated as (weight in kg)/(height in m)² for each individual. Subjects with BMI more than or equal to 25kg/ m² were considered as obese according to revised guidelines based on

consensus developed through discussions by a Prevention and Management of Obesity and Metabolic Syndrome group [13]. Resting Blood pressure were recorded using sphygmomanometer in lying down position after a resting period of half an hour. Systolic BP more than or equal to 140 mm of Hg and diastolic BP more than or equal to 90 mm of Hg were considered as hypertension.

After an overnight fasting of 12 hours, 5 mL of venous blood sample were collected under sterile conditions in plain tubes. Blood was withdrawn with disposable syringes from the cubital vein of the subjects. Serum was separated by centrifuging the sample at 2500 revolutions per minute. 1mL of serum of each subject was aliquoted in separate eppendorfs and stored at -20°C . Serum leptin levels were measured using Human Leptin ELISA kit procured from dbc – Diagnostics Biochem Canada Inc.

Statistical analysis

All the data were entered in excel sheet and analysed using SPSS software. Simple frequency tests

were calculated. Descriptive statistics of age, gender, blood pressure, body mass index, serum leptin were derived as Mean and standard deviation. Chi square test was done for comparison of variables. Logistic regression was done to determine the relation of leptin, body mass index and hypertension. It was also adjusted for obesity to know the direct association of serum leptin level and hypertension.

RESULTS

Descriptive statistics is given in Table:1. The cases had mean age of 51.8 ± 5.7 years, systolic BP of 150 ± 14 mm of Hg, diastolic BP of 96 ± 8 mm of Hg and body mass index of 25.9 ± 3.9 kg/m^2 , serum leptin levels of 31.9 ± 32.5 . Binary logistic regression for leptin and body mass index showed a significant association $p=0.000$ as shown in Table: 2. and this data were further adjusted for obesity. Even after eliminating the influence of obesity, serum leptin levels showed a statistically significant association with essential hypertension ($p=0.000$).

Table-1: Descriptive statistics

	Mean	Standard deviation
Age	51.8	5.7
Systolic BP	150	14
Diastolic BP	96	8
Body mass index	25.9	3.9
Serum leptin levels	31.9	32.5

Table-2: Binary logistic regression

	B	S.E	Wald	df	Sig.	Exp(B)	95% C.I for EXP (B)	
							Lower	Upper
Leptin cate	-2.276	.626	13.223	1	.000	.103	.030	.350
Obesity	.544	.631	.744	1	.388	1.723	.500	5.932
Constant	2.629	.849	9.599	1	.002	13.859		

P value <0.05 is significant

After adjusting for influence of obesity, most of the hypertensives have high leptin level

DISCUSSIONS

Till date many animal studies have revealed the significant relation of high serum leptin levels with high blood pressure [14]. Human studies are less consistent with the same findings [15,16]. It is a recognized fact that serum leptin levels correlates well with obesity. As expected, this was true in our study. Serum leptin levels were found to be increased in obese hypertensives study which was in line with many other similar studies. In contrast few other studies couldn't obtain a direct positive association between serum leptin levels and blood pressure. Raised serum leptin levels in our study could be due to selective leptin

resistance where the metabolic effects are blunted leading to obesity while the neural action of leptin maintaining the normal vascular tone is impaired, leading to increased pressor effects, thereby increasing blood pressure. One other possibility could be via insulin resistance. Normally leptin is found to enhance the sensitivity of insulin and simultaneously lowering its production. Hyperleptinemia due to leptin resistance could have caused insulin resistance. Insulin resistance and hyperinsulinemia per se are linked to hypertension [17,18]. The difference between serum leptin levels among obese hypertensives and non obese hypertensives was statistically significant in our study. It was also found that even after adjusting for influence of obesity, leptin levels were significantly high in hypertensives. Thus we found that high levels of leptin had direct role in hypertension independent of obesity.

There could be several mechanisms for such a finding. Elevated serum leptin levels could be linked to increased blood pressure via adrenergic or renal mechanisms and have been highly suggested as another possible link for hypertension. Further research studies should be conducted to know the possible mechanism for our finding. The limitations of this study were small sample size. Differences in study design, ethnic and racial differences across study populations may account for the inconsistency of various other similar studies.

CONCLUSION

We conclude from our study that serum leptin levels are high in essential hypertension probably due to selective leptin resistance and this in turn needs further evaluation on a larger population. Leptin is an attractive target for extended research particularly in deriving at the pathophysiology of essential hypertension thereby a potential candidate for pharmacotherapy.

Acknowledgement

We thank Dr. TAMILARASI MD, Assistant professor, Department of Community Medicine, Stanley Medical College for her statistical assistance.

REFERENCES

- Margetic S, Gazzola C, Pegg GG, Hill RA. Leptin: a review of its peripheral actions and interactions. *International journal of obesity*. 2002 Nov;26(11):1407.
- Beltowski J. Role of leptin in blood pressure regulation and arterial hypertension. *Journal of hypertension*. 2006 May 1;24(5):789-801.
- Wang J, Wang H, Luo W, Guo C, Wang J, Chen YE, Chang L, Eitzman DT. Leptin-induced endothelial dysfunction is mediated by sympathetic nervous system activity. *Journal of the American Heart Association*. 2013 Sep 16;2(5):e000299.
- Mark AL. Selective leptin resistance revisited. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2013 Jul 24;305(6):R566-81.
- Kshatriya S, Liu K, Salah A, Szombathy T, Freeman RH, Reams GP, Spear RM, Villarreal D. Obesity hypertension: the regulatory role of leptin. *International journal of hypertension*. 2011 Jan 3;2011.
- da Silva AA, do Carmo JM, Hall JE. Role of leptin and CNS melanocortins in obesity hypertension. *Current opinion in nephrology and hypertension*. 2013 Mar;22(2):135.
- Hall JE, do Carmo JM, da Silva AA, Wang Z, Hall ME. Obesity-induced hypertension: interaction of neurohumoral and renal mechanisms. *Circulation research*. 2015 Mar 13;116(6):991-1006.
- Kerimkulova AS, Lunegova OS, Mirrakhimov AE, Alibaeva NT, Neronova KV, Baïramukova AA, Mirrakhimov EM. Association of leptin with obesity and hypertension in an ethnic Kyrgyz group. *Terapevticheskii arkhiv*. 2014;86(1):49-53.
- Haque Z, Lakho GR, Nafees M, Shahid KU, Mazahir I. Serum leptin levels correlation with high blood pressure in adult females. *Journal of the College of Physicians and Surgeons--Pakistan: JCPSP*. 2006 Jul;16(7):450-4.
- Paleczny, B., A. Siennicka, M. Zacharski, E.A. Jankowska, B. Ponikowska and P. Ponikowski (2016). Increased body fat is associated with potentiation of blood pressure response to hypoxia in healthy men: relations with insulin and leptin. *Clin Auton Res.*, 26(2): 107-16
- Almeida-Pititto BD, Gimeno SG, Freire RD, Ribeiro-Filho FF, Ferreira SR. Leptin is not associated independently with hypertension in Japanese-Brazilian women. *Brazilian journal of medical and biological research*. 2006 Jan;39(1):99-105.
- Wada K, Yatsuya H, Tamakoshi K, Otsuka R, Fujii C, Matsushita K, Sugiura K, Toyoshima H. A positive association between leptin and blood pressure of normal range in Japanese men. *Hypertension research*. 2006 Jul;29(7):485.
- Misra A, Chowbey P, Makkar BM, Vikram NK, Wasir JS, Chadha D, Joshi SR. Consensus statement for diagnosis of obesity, abdominal obesity and the metabolic syndrome for Asian Indians and recommendations for physical activity, medical and surgical management. *JAPI*. 2009 Feb;57(2):163-70.
- Palei AC, Spradley FT, Granger JP. Chronic hyperleptinemia results in the development of hypertension in pregnant rats. *American Journal of Physiology-Heart and Circulatory Physiology*. 2015 Mar 11.
- Simonds SE, Pryor JT, Cowley MA. Does leptin cause an increase in blood pressure in animals and humans?. *Current opinion in nephrology and hypertension*. 2017 Jan 1;26(1):20-5.
- Tsai JP, Hsu BG, Lee CJ, Hsieh YH, Chen YC, Wang JH. Serum leptin is a predictor for central arterial stiffness in hypertensive patients. *Nephrology*. 2017 Oct;22(10):783-9.
- Soleimani M. Insulin resistance and hypertension: new insights. *Kidney international*. 2015 Mar 1;87(3):497-9.
- Manucha W, Ritchie B, Ferder L. Hypertension and insulin resistance: implications of mitochondrial dysfunction. *Current hypertension reports*. 2015 Jan 1;17(1):504.