

## Obturator Hip Dislocation of a Young Sportsman: Case Report and Review of the Literature

Krimech MO\*, Jeddi D, Azirar A, Bassir RA, Boufettal M, Kharmaz M, Lamrani MA, Ouadghiri M, bardouni AE, Mahfoud M, Berrada MS

Traumatology-Orthopedics surgery Department University Hospital Ibn Sina Rabat – Morocco

\*Corresponding author: Krimech MO

| Received: 14.02.2019 | Accepted: 24.02.2019 | Published: 28.02.2019

DOI: [10.21276/sjams.2019.7.2.50](https://doi.org/10.21276/sjams.2019.7.2.50)

### Abstract

### Case Report

**Introduction:** Traumatic dislocation of the coxofemoral joint is defined as the permanent posterior or anterior displacement of the femoral head out of the acetabular cavity. It is generally the consequence of a violent trauma, most often an accident on the public road. Our case it is a post-traumatic hip obturator dislocation, managed in our department. **Clinical case and results:** 16 years old patient, with no particular history, victim of a fall from his motorcycle during a race, causing a closed trauma to his left hip, right ankle and left wrist. He comes with a vicious attitude: left lower limb in flexion, abduction, external rotation. An X-ray requested objective obturator dislocation with a small bone fragment of the head. A bone reconstruction CT was completed in favour of obturator dislocation with head fracture type 1 according to Pipkin's classification. Patient sent directly to the block, he benefits from a reduction under sedation according to Boehler's maneuver, control by scopy, then immobilisation by zimmer splint after testing the stability of dislocation. Patient is hospitalized in our ward for monitoring then follow-up in consultation. Results were satisfactory, removal of the zimmer splint at the 6th week. Total support was possible at week 12, with complete mobility, patient reviewed afterworld without clinico-radiological signs of aseptic necrosis of the femoral-head. **Conclusion:** Obturator dislocation remains rare, especially in sports, diagnosis and early reduction is essential for good progress. However, we should not to ignore the associated fracture which still darkens the prognosis.

**Keywords:** Hip, obturator dislocation, traumatic, reduction, treatment.

**Copyright @ 2019:** This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

## INTRODUCTION

Traumatic dislocation of the coxofemoral joint is defined as the permanent posterior or anterior displacement of the femoral head out of the acetabular cavity. It is generally the consequence of a violent trauma, most often an accident on the public road [1]. Among these anterior dislocations, Epstein and Wiss [3] distinguish type a, upper or pubic and type B, lower or obturator. De Lee *et al.* [4] specified this classification with a type 1, upper with two varieties, one pubic and the other infraspinal, and a type 2, lower also with two varieties, one obturator and the other perineal. In our case it is a post-traumatic hip obturator dislocation, managed in our department.

## OBSERVATIONS

This is a 16 years old patient, with no particular history, victim of a fall from his motorcycle during a race, causing a closed trauma to his left hip,

right ankle and left wrist. He comes with a vicious attitude: left lower limb in flexion, abduction, external rotation. An X-ray requested objective obturator dislocation with a small bone fragment of the head. A bone reconstruction CT was completed in favour of obturator dislocation with head fracture type 1 according to Pipkin's classification. Patient sent directly to the block, he benefits from a reduction under sedation according to Boehler's maneuver, control by scopy, then immobilisation by zimmer splint after testing the stability of dislocation. Patient is hospitalized in our ward for monitoring then follow-up in consultation.

Results were satisfactory, removal of the zimmer splint at the 6th week. Total support was possible at week 12, patient reviewed afterworld with complete mobility and without clinico-radiological signs of aseptic necrosis of the femoral-head.



**Fig-1: Clinical view of the hip dislocation: lower limb in flexion abduction rotation externe**



**Fig-2: Radiography of the dislocated hip**



**Fig-3 : TDM 3D**



**Fig-4 : Radiography post reduction**

## DISCUSSION

Very little work has been done on traumatic obturator dislocations [2]. Percentage of obturator dislocations reported in traumatic dislocation series is between 6 % and 10 %.

It is common experience in hip replacement surgery that hip flexion and an external rotation and adduction force to the femur produces anterior hip dislocation after capsulotomy in patients who are anesthetized. Whether this is the mechanism during a road traffic accident is not known. The strong anterior capsule or Y shaped ligament of Bigelow is a strong disincentive to anterior dislocation, which may be part of the reason that this pattern of dislocation is less common. The anterior hip fracture may be produced as a result of the ligament or capsule failing to yield and causing an avulsion fracture. Inferior dislocation occurs when the hip is forced into abduction and external rotation [6].

Anterior capsular lesions (tearing or disinsertion) are constant. Associated osteoarticular lesions are frequent, foremost among which are cephalic osteochondral fractures [5], attributed to an impact of the femoral head on the lower part of the anterior acetabular blade. Fracture of the anterior acetabular blade is also possible other lesions have been reported: fracture of the ilio and ischio-pubic branches, contralateral hip dislocation, femoral fractures, neck, shaft and major trochanter. Lesions of the femoral vessels and the femoral nerves or obturator must be sought systematically, even if they are rare. Initial management involves an attempt at orthopedic reduction under general anaesthesia with complete muscle relaxation, with the particularity of: the difficulty of reduction and the risk of iatrogenic fracture [2].

Consequences after reduction orthopaedic or surgical procedures are not consensual. Currently there is no scientific argument to justify the interest of traction and unloading in the reduction of the risk of cephalic necrosis of the femoral head. Catonné *et al.*

recommend early support relieved and then total support at j15 with eviction of external rotation for 3 weeks as part of the previous dislocations [7]. The risk of necrosis occurring increases with reduction delay. Hoogard observed 47% necrosis when the reduction time exceeded 6 hours. But these figures are mostly lesions associated with fractures of the acetabulum or femoral head, this rate is certainly lower in isolated dislocations [8].

In 1957, Pipkin<sup>42</sup> established a four-stage classification that often referred to. In 2001, Yoon<sup>57</sup> described a classification that is simpler to understand based on the anatomical description of the lesions and which presents therapeutic implications that should impose it as a reference classification [1]. The publications dealing with femoral head fractures are rare and the series are short, it is difficult to identify a univocal therapeutic strategy.

## CONCLUSION

Obturator dislocation remains rare, especially in sports, diagnosis and early reduction is essential for good progress. However, we should not to ignore the associated fracture which still darkens the prognosis.

## REFERENCES

1. Richardson P, Young JW, Porter D. CT detection of cortical fracture of the femoral head associated with posterior hip dislocation. *AJR. American journal of roentgenology.* 1990 Jul;155(1):93-4.
2. Boyer P, Bassaine M, Hutten D. Traumatic obturator foramen hip dislocation: a case report and review of the literature. *Revue de chirurgie orthopedique et reparatrice de l'appareil moteur.* 2004 Nov;90(7):673-7.
3. Epstein HC, Wiss DA. Traumatic anterior dislocation of the hip. *Orthopedics.* 1985; 8, 130-134.
4. DeLee JC, Evans JA, Thomas JA. Anterior dislocation of the hip and associated femoral-head fractures. *The Journal of bone and joint surgery. American volume.* 1980 Sep;62(6):960-4.

5. Stannard JP, Harris HW, Volgas DA, Alonso JE. Functional outcome of patients with femoral head fractures associated with hip dislocations. *Clinical Orthopaedics and Related Research*. 2000 Aug 1;377:44-56.
6. Phillips AM, Konchwalla A. The pathologic features and mechanism of traumatic dislocation of the hip. *Clinical Orthopaedics and Related Research*. 2000 Aug 1;377:7-10.
7. Catonné Y, Meyer A, Soriali E, Biette G. Pathologie du complexe pelvi-fémoral du sportif. *Pathologie du complexe pelvi-fémoral du sportif*. 2009;88:99.
8. Elouakili I, Ouchrif Y, Ouakrim R, Lamrani O, Kharmaz M, Ismael F, Lahlou A, Bardouni AE, Mahfoud M, Berrada MS, Yaacoubi ME. Luxation obturatrice de la hanche: un traumatisme rare en pratique sportive. *Pan African Medical Journal*. 2015;21(1).
9. Bouya A, Elantri I, Bensaleh MR, Tanane M, Zine A, Jaafar A. Luxation obturatrice de la hanche: survenue rare en milieu sportif. *Journal de Traumatologie du Sport*. 2018 Mar 1;35(1):40-3.