

Prevalence of Thyroid Disorders in Patients of Diabetes Mellitus in a Tertiary Care Hospital in North India

Jaswinder Kaur¹, Kusum Bali^{2*}, Satyam Krishan³, Venus Sharma³, Manmeet Pal Kaur³, Uma Arora⁴¹Assistant Professor, Department of Internal Medicine, Punjab Institute of Medical Sciences, Garha Road, Jalandhar, Punjab 144006, India²Associate Professor, Department of Internal Medicine, Punjab Institute of Medical Sciences, Garha Road, Jalandhar, Punjab 144006, India³Interns, Department of Internal Medicine, Punjab Institute of Medical Sciences & Hospital, Garha Road, Jalandhar, Punjab 144006, India⁴Professor, Department of Biochemistry, Punjab Institute of Medical Sciences & Hospital, Garha Road, Jalandhar, Punjab 144006, India

*Corresponding author: Kusum Bali

| Received: 28.04.2019 | Accepted: 06.05.2019 | Published: 09.05.2019

DOI: [10.21276/sjams.2019.7.5.2](https://doi.org/10.21276/sjams.2019.7.5.2)

Abstract

Original Research Article

Introduction: Diabetes Mellitus is characterised by relative or absolute deficiency in insulin secretion as well as insulin action. It is also associated with carbohydrate, protein and lipid metabolism disturbances. Patients of diabetes mellitus have higher prevalence of thyroid disorders as compared to the general population. **Materials and Methods:** 243 patients of diabetes mellitus, coming to OPD department of Internal Medicine as well indoor patients in PIMS Medical College and Hospital were included. Measurement of weight, height, Body Mass Index, Blood pressure, Fasting plasma Glucose levels, Post prandial glucose, HbA1C, lipid profile, Renal Function Tests, Free Triiodothyronine FT3, Free Tetraiodothyronine (FT4) and Thyrotropin TSH levels tests were done. **Results:** Analysis shows that the prevalence of thyroid disorders is high in patients with type 2 diabetes mellitus. Abnormal thyroid functions were detected in 77(31.7%) patients 38(15.6%) had Subclinical Hypothyroidism. 28(11.5%) patients had Overt Hypothyroidism whereas 11 (4.6%) patients had Hyperthyroidism. **Conclusion:** Type-2 Diabetes Mellitus patients have higher prevalence of thyroid dysfunction in the form of Subclinical Hypothyroidism, overt Hypothyroidism and Hyperthyroidism. So these patients require screening for thyroid functions.

Keywords: Type- 2 Diabetes Mellitus, Hypothyroidism, Subclinical Hypothyroidism.

Copyright @ 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

Diabetes Mellitus is a very common endocrine disorder. In Diabetes Mellitus there is either absolute or relative deficiency in insulin secretion and / or its action. There are also disturbances of carbohydrate, lipid and protein metabolism [1].

The prevalence of Diabetes is expected to rise to 300 million (7.8%) by the year 2030 [2]. The prevalence of thyroid dysfunction is increasing in patients of Diabetes Mellitus.

Insulin secretion is directly controlled by the thyroid hormones. In Hypothyroidism the glucose induced insulin secretion by the beta cells is reduced. Whereas in Hyperthyroidism the catecholamines and insulin resistance both are increased [3, 4]. Several studies have indicated the co- existence of both endocrine disorders. The prevalence of thyroid dysfunction in Diabetic patients have been reported to vary from 2.2 to 17 % [5, 6].

Diabetes Mellitus affects thyroid at two sites, at the level of the hypothalamus by controlling TSH release and also at peripheral tissues by converting T4 to T3 [7, 8].

AIMS AND OBJECTIVES

- To study the prevalence of thyroid disorders either Subclinical hypothyroidism, Overt hypothyroidism, or Hyperthyroidism in patients with type 2 Diabetes Mellitus.
- To find out any correlation between the glycosylated Hemoglobin HbA1C and TSH levels.
- To find out any association between the TSH levels and the duration of diabetes.

MATERIALS AND METHODS

This cross sectional study was conducted in Punjab Institute of Medical Sciences, Jalandhar. A Total of 243 patients coming to Medicine OPD as well as indoor patients having type 2 Diabetes Mellitus were enrolled. After obtaining informed consent, these

patients had to undergo weight and height monitoring, blood pressure readings, routine investigations like fasting blood sugar (FBS), Post prandial plasma glucose levels, HbA1C by CLIA (Chemilumescence immunoassay) technique, Renal function tests. Liver function tests, lipid profile, urine albumin creatinine ratio, Fundus examination, Thyroid Hormones like FT3, FT4 and TSH by CLIA (Chemilumescence Immunoassay) technique by Roshe E411 fully automated analyzer.

This study was approved by the institutional ethics committee.


RESULTS

243 patients having Diabetes Mellitus aged from 17 years to 84 years were enrolled in PIMS hospital coming to Medicine OPDs and indoor patients admitted in medical wards in our hospital.

Table-1: Age distribution

Age	No. of Patients	% of patients
15-30	5	2%
31-45	30	12%
46-60	124	51%
61-75	74	30%
75-90	10	4%


Bar diagram representing age distribution



Out of 243 patients 83 were males (34%) and 160 were females (66%).

Table representing Gender distribution

Gender	No. of Patients
Female	160
Male	83
Grand Total	243



Pie chart representing gender distribution


Out of 243 patients, 32% patients had Diabetic Nephropathy which is a microvascular complication of

Diabetes mellitus where as 68% patients had normal GFR.

Table representing the prevalence of Diabetic Nephropathy in our study group

Patients having Diabetic Nephropathy	Female	Male	Grand Total
Normal	46%	22%	68%
Reduced GFR and increased Creatinine Levels	20%	12%	32%
Grand Total	66%	34%	100%

Bar diagram representing the prevalence of Diabetic Nephropathy according to sex distribution.




Our study found abnormal thyroid functions in 77 patients (31.7%) patients. 166(68.3 %) patients had normal TSH levels.38 patients (15.6%) had Subclinical Hypothyroidism. 28 (11.5%) patients were found to

have Overt hypothyroidism.11 patients (4.5%) patients had Hyperthyroidism. Subclinical Hypothyroidism was the most common thyroid abnormality found in diabetic patients.

Table showing the prevalence of Thyroid disorders in Diabetic patients

Prevalence of Thyroid disorders	% of Patients	No. of Patients
Normal (TSH between 0.4 - 4.6)	68.3%	166
Sub Clinical Hypothyroidism (TSH between 4.6 -10)	15.6%	38
Overt Hypothyroidism (TSH more than 10)	11.5%	28
Hyperthyroidism (TSH less than 0.4)	4.5%	11



Bar diagram representing prevalence of Thyroid disorders in Diabetes Mellitus patients.

Our study however did not find any correlation between the glycosylated haemoglobin HbA1C levels and the TSH levels.

Correlations between HbA1C and TSH levels			
		HBA1C	TSH
HBA1C	Pearson Correlation	1	-.094
	Sig. (2-tailed)		.174
	N	210	210
TSH	Pearson Correlation	-.094	1
	Sig. (2-tailed)	.174	
	N	210	210

Our study also did not find any association between the TSH levels and the duration of diabetes.

Table representing the prevalence of hypothyroidism according to distribution of duration of Diabetes

Category	Prevalence of Hypothyroidism			Grand Total
	Normal	Hyperthyroidism	Overt Hypothyroidism	
Newly Detected	12	1	3	16
Less than 5 years	49	4	25	78
5-10 Years	64	4	22	90
10-15 Years	21	0	8	29
More than 15 Years	20	2	8	30
Grand Total	166	11	66	243

No association found between the prevalence of Thyroid disorders and the Duration of Diabetes Mellitus.

Chi-Square Tests			
	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	17.165 ^a	12	.143
Likelihood Ratio	18.293	12	.107
N of Valid Cases	243		

a. 11 cells (55.0%) have expected count less than 5. The minimum expected count is .72.

DISCUSSION

It has been well known that there is increased prevalence of thyroid disorders in patients of Type 1 Diabetes Mellitus due to common autoimmune origin, But a lot of studies in the past have reported increased prevalence in type 2 Diabetes as well. Thyroid Hormones increase the hepatocyte plasma concentration of GLUT 2, the main glucose transporter in the liver. These increased level of GLUT 2 lead to abnormal glucose metabolism [9].

Our study found a high prevalence of Thyroid disorders in Diabetic patients about 31.7%, with Subclinical Hypothyroidism in 15.6% patients, Overt Hypothyroidism in 11% patients and Hyperthyroidism in 4.6% patients. Several previous studies have also reported a higher prevalence, Papaziffiropoulou *et al.*, (the prevalence of thyroid disorders was 12.3%) [10]. Nobre *et al.*, study also found out prevalence of thyroid disorders 12.7% [11]. Our study however did not find any correlation between the TSH levels and the HbA1C levels. Our study also did not find any association between the TSH levels and the duration of Diabetes mellitus. Our results are in concordance with Diez *et al.*, whose study also did not find any significant relationship between the thyroid dysfunction and the duration of diabetes mellitus type 2 [12].

In hyperthyroidism increased lipolysis is observed which result in increase in FFA which further stimulates hepatic gluconeogenesis. Hyperthyroidism is also associated with increased growth hormone, glucagon and catecholamine levels which further contribute to impaired glucose tolerance [13, 14].

In Hypothyroidism there is reduced rate of liver glucose production which accounts for decrease in the insulin requirement in diabetic patients having

coexisting hypothyroidism. Also both clinical and subclinical hypothyroidism have been recognised as insulin resistant states [15, 16].

In diabetes mellitus there is also increased prevalence of low T3 state. This “low T3 state” could possibly be explained by an impairment in peripheral conversion of T4 to T3. This state normalizes with improved glycemic control.

CONCLUSION

Since there is an increased prevalence of thyroid disorder in Type 2 diabetes, so screening of all the diabetic patients for thyroid profile should be done regularly in order to achieve good glycemic control as well as to decrease the complications of diabetes mellitus Type 2.

Results obtained from our study show that Overt hypothyroidism as well as subclinical hypothyroidism is frequently observed in patients Type 2 diabetes mellitus. However more studies with more number of patients are required in future to study this inter-relationship between these two endocrine disorders.

There is a complex interdependent interaction between these two endocrine disorders. Thyroid function tests should be recommended in patients with diabetes mellitus as failure to recognize the thyroid disorder may be the cause of poor diabetic control.

BIBLIOGRAPHY

1. Tiwari AK, Madhusudanarao J. Diabetes mellitus and multiple therapeutic approaches of phytochemicals: Present status and future prospects. *Current Science* 2002 July; 83(1):30-38.

2. Maratou E, Hadjidakis DJ, Peppas M, Alevizaki M, Tsegka K, Lambadiari V, Mitrou P, Boutati E, Kollias A, Economopoulos T, Raptis SA. Studies of insulin resistance in patients with clinical and subclinical hyperthyroidism. *European journal of endocrinology*. 2010 Oct 1;163(4):625-30.
3. Mitrou P, Raptis SA, Dimitriadis G. Insulin action in hyperthyroidism: a focus on muscle and adipose tissue. *Endocrine Reviews*. 2010 Oct 1;31(5):663-79.
4. Duntas LH, Orgiazzi J, Brabant G. The interface between thyroid and diabetes mellitus. *Clinical endocrinology*. 2011 Jul;75(1):1-9.
5. Sugure DD, McEvoy M, Drury MI. Thyroid disease in diabetics. *Postgrad Med Journal*, 1999; 680-684.
6. Shah SN. Thyroid disease in diabetes mellitus. *The Journal of the Association of Physicians of India*. 1984 Dec;32(12):1057-9.
7. Vikhe VB, Kanitkar SA, Tamakuwala KK, Gaikwad AN, Kalyan M, Agarwal RR. Thyroid dysfunction in patients with type 2 diabetes mellitus at tertiary care centre. *Natl J Med Res*. 2013 Dec;3(4):377-380.
8. Chen HS, Wu TE, Jap TS, Lu RA, Wang ML, Chen RL, Lin HD. Subclinical hypothyroidism is a risk factor for nephropathy and cardiovascular diseases in Type 2 diabetic patients. *Diabetic Medicine*. 2007 Dec;24(12):1336-1344.
9. Kemp HF, Hundal HS, Taylor PM. Glucose transport correlates with GLUT2 abundance in rat liver during altered thyroid status. *Molecular and cellular endocrinology*. 1997 Apr 4;128(1-2):97-102.
10. Papazafiropoulou A, Sotiropoulos A, Kokolaki A, Kardara M, Stamataki P, Pappas S. Prevalence of thyroid dysfunction among greek type 2 diabetic patients attending an outpatient clinic. *Journal of clinical medicine research*. 2010 Apr;2(2):75-78.
11. Nobre E, Jorge Z, Pratas S, Silva C, Castro J. Profile of the thyroid function in a population with type-2 diabetes mellitus. In 21st Joint Meeting of the British Endocrine Societies 2002 Mar 1 (Vol. 3). BioScientifica.
12. Diez JJ, Sánchez P, Iglesias P. Prevalence of thyroid dysfunction in patients with type 2 diabetes. *Experimental and Clinical Endocrinology & Diabetes*. 2011 Apr;119(04):201-7.
13. Miki N, Ono M, Hizuka N, Aoki T, Demura H. Thyroid hormone modulation of the hypothalamic growth hormone (GH)-releasing factor-pituitary GH axis in the rat. *The Journal of clinical investigation*. 1992 Jul 1;90(1):113-120.
14. Sestoft L, Christensen NJ, Saltin B. Responses of glucose and glucoregulatory hormones to exercise in thyrotoxic and myxoedematous patients before and after 3 months of treatment. *Clinical Science*. 1991 Jul 1;81(1):91-99.
15. Dimitriadis G, Miron P, Lambadiari V. Insulin action in hypothyroidism. *Journal of clinical Endocrinology and metabolism*, 2006, 91(12), 4930-4937.
16. Maratou E, Hadjidakis DJ, Kollias A, Tsegka K, Peppas M, Alevizaki M, Mitrou P, Lambadiari V, Boutati E, Nikzas D, Tountas N. Studies of insulin resistance in patients with clinical and subclinical hypothyroidism. *European journal of endocrinology*. 2009 May 1;160(5):785-90.