

## Case Report

### Papillon Lefevre Syndrome-A Case Series of Siblings

Dr. Munaza Shafi<sup>1</sup>, Dr. Suhail Majid<sup>2</sup>, Dr. Roobal Behal<sup>3</sup>, Dr. Reyaz Ahmed<sup>1</sup>

<sup>1</sup>P.G Student, <sup>2</sup>Associate Prof. & Head, <sup>3</sup>Lecturer; Department of Periodontics, Government Dental College, Srinagar.

#### Corresponding author

Dr. Munaza Shafi

Email: [drmunaza.shafi@gmail.com](mailto:drmunaza.shafi@gmail.com)

**Abstract:** Papillon Lefèvre syndrome is a rare disease characterized by skin lesions caused by palmar-plantar hyperkeratosis, and severe periodontal destruction involving both the primary and permanent dentitions. It is transmitted as an autosomal recessive condition and consanguinity of parents is evident in about one third of cases. The palmar plantar keratoderma typically has its onset between the ages of 1 and 4 years and severe periodontitis starts at the age of 3 or 4 years. Two cases of Papillon-Lefevre syndrome in the same family (siblings) are presented. A 15 years old girl and 13 years old boy have all the characteristic features of syndrome.

**Keywords:** Papillon lefevre syndrome, Palmar Plantar keratoderma, severe periodontitis

#### INTRODUCTION

Papillon-Lefèvre syndrome (PLS) is characterized by hyperkeratosis of hands and feet and by a generalized aggressive periodontitis in both the primary and the permanent dentition [1]. The syndrome is a rare autosomal recessive trait with an incidence of one to four persons per million. Parental consanguinity is demonstrated in between 20-40% of the cases [1]. Associated features include calcification of the falx cerebri and the choroid plexus, and retardation of somatic development [2]. The recently identified genetic defect in PLS has been mapped to chromosome 11q14-q21, that involves mutations of cathepsin C [3]. More than 90% reduction in cathepsin C activity has been found in PLS patients [4, 5]. Many reports describe an impaired chemotactic and phagocytic function of polymorphonuclear leukocytes (PMNs) [6-8]. The palmoplantar keratoderma typically has its onset between the ages one and four years. The sharply demarcated erythematous keratotic plaques may occur focally, but usually involve the entire surface of palms and soles resulting in foul-smelling odor [9-10].

The second major feature of PLS is severe periodontitis, which starts at the age of three or four years [5]. The development and eruption of deciduous teeth proceeds normally, but their eruption is associated with gingival inflammation and subsequent rapid destruction of the periodontium [10, 11]. The resulting periodontitis characteristically is unresponsive to traditional periodontal treatment modalities and the primary dentition is usually exfoliated prematurely by the age four years. After exfoliation, the inflammation subsides and gingiva appears healthy. However, with eruption of the permanent dentition the process of gingivitis and periodontitis is usually repeated and there is subsequent premature exfoliation of the permanent teeth, although the third molars are sometimes spared [12].

Radiographic features are characterized by generalized loss of alveolar bone. Gorlin *et al.* have added the third feature of dural calcification [13]. Reyes also observed radiographic evidence of intracranial calcification [14]. Histopathologic findings of affected skin consist of hyperkeratosis, occasional patches of parakeratosis, acanthosis, and slight perivascular inflammatory infiltrate [15].

#### CASE No. 1

A 15-year-old female patient presented to the department of Periodontics and oral Implantology, Government dental college, Srinagar (India) with the chief complaint of multiple loss of teeth. Clinical history revealed that she had normal emergence of deciduous teeth at 8-9 months of age, which started loosening at three years and were all eventually lost by four years of age. Patient was not sure about the time of eruption of permanent teeth, but described gingival bleeding during brushing and eating, after the eruption of permanent teeth. There was loosening of permanent teeth from 10 years of age and eventually all the permanent teeth were lost by 12 years of age, except for one mobile lower third molar tooth (fig 1). Bleeding was also associated at the time of tooth loss.

Dermatological examination revealed dry skin with normal development of hair and nails. There were symmetric, well-demarcated, yellowish, keratotic, and confluent plaques affecting the skin of her palms and soles, also extending onto the dorsal surfaces of hands and feet (fig 1). In the light of these findings Papillon-Lefevre Syndrome was diagnosed in the patient. This diagnosis was confirmed by the dermatology clinic as palmar plantar hyperkeratosis.

Intraoral examination showed premature loss of all teeth, except for the left mandibular third molar which was covered with deposits and it showed severe

mobility. No other abnormality was detected in relation to soft tissues. Patient had a reduced facial height due to resorption of the alveolar ridge (fig 1).

Routine blood investigations and liver function tests were carried out and were found to be within the normative range of values.

### Management

Since the patient was a young teenage girl, it was necessary to manage the esthetic needs of the patient as a priority. It was appropriate to give her complete dentures, considering her functional requirements, such as mastication and speech. Prior to the routine clinical procedure for complete dentures, the mandibular third molar was extracted. The dentures were fabricated in the laboratory and inserted into the patient's mouth (fig 1). Routine postdenture instructions were given to the patient

### CASE No. 2

A 13-year-old boy patient (above patient's brother) reported to the department of periodontics and oral Implantology, Government dental college Srinagar (India) with a chief complaint of loosening of multiple teeth (fig 2).

On examination, there were symmetrical, well demarcated, keratotic and confluent plaques affecting the skin of the palms and soles which extended to the dorsal surface of the feet and hands. The skin was dry and rough on palpation.

On intraoral examination, mobility was present in all the teeth (fig 2). The gingiva was red, soft and edematous with deep periodontal pockets and bleeding on probing. The mucosa in other areas of the oral cavity appears to be normal. OPG of the patient showed severe alveolar bone loss in relation to all the teeth.(Fig 2).



Case 1 (fig. 1)



Case 2 (fig. 2)

Laboratory investigation was carried out, which included haematological and biochemical assessment. The results were within normal limits.

### Management

First of all, the patient was prescribed an antibiotic of doxycycline 100mg OD x 3 weeks and a mouth rinse of 0.2% chlorhexidine gluconate and educated for oral hygiene to prolong the life of his dentition.

### DISCUSSION

Papillon Lefèvre syndrome was first described by Papillon and Lefèvre in 1924. The disease is characterized by diffuse palmoplantar hyperkeratosis and juvenile periodontitis. Haneke used the following three criteria to classify a case as PLS: (a) palmoplantar hyperkeratosis; (b) loss of primary and permanent teeth; and (c) autosomal recessive inheritance.

Although the exact pathogenesis of this syndrome remains relatively obscure, immunologic, microbiologic, and genetic bases have been proposed. Microbiological studies have demonstrated *Actinobacillus actinomycetemcomitans*, *Porphyromon – as gingivalis*, *Fusobacterium nucleatum*, and *Treponema denticola* organisms, suggesting that many pathogens may be involved in the disease process. Previous case reports and studies have reported

that *A. actinomycetemcomitans* plays a significant role in the pathogenesis and progression of the rapid periodontal breakdown seen in PLS [18, 19, 20].

It has been suggested that the presence of periodontal pathogens alone is not sufficient for the expression of PLS, and other factors, such as host response, play an important role in the pathogenesis of the disease process. Several authors have suggested an abnormal neutrophil dysfunction with PLS [21, 22] to explain the pathogenesis.

The differential diagnoses include Hiam-Munk syndrome and hypophosphatasia. Hiam-Munk syndrome also exhibits arachnodactyly, acroosteolysis, atrophy of nails, and deformity of the phalanges in the hands. None of these features were found in the present cases. In hypophosphatasia, deficiency of alkaline phosphatase activity is seen; but in our cases the values were within normal limits and therefore this differential diagnosis could be excluded.

Identification of specific periodontal pathogens and antibiotic therapy appropriate to these microorganisms, along with extraction of severely periodontally compromised teeth, can prolong the viability of nonaffected teeth. Newer therapeutic modalities involve the use of oral retinoids, such as acitretin and isotretinoin [10]. In the future, stem cell

therapy can be expected to open up new vistas in the dental treatment of such children.

### References

1. Kaya FA; Polat ZS; Baran EA; Tekin GG; Papillon-Lefevre Syndrome-3 Years Follow Up: A Case Report. *International Dental And Medical Disorders*, 2008; 1(1): 24-28.
2. Gorlin RJ, Cohen MM, Levin LS; *Syndromes of the Head and Neck*, 3<sup>rd</sup> edition, Oxford: Oxford University Press, 1990: 853–855.
3. Hart T, Hart P, Bowden D, Michalec M, Callison S, Walker S *et al.*; Mutation of the cathepsin C gene are responsible for Papillon–Lefèvre syndrome. *Journal of Medical Genetics*, 1999; 36(12): 881–887.
4. Toomes C, James J, Wood AJ, Wu CL, McCormick D, Lench N *et al.*; Loss-of-function mutations in the cathepsin C gene result in periodontal disease and palmoplantar keratosis. *Nature Genetics*, 1999; 23: 421–424.
5. Hart PS, Zhang Y, Firatli E, Uygur C, Lotfazar M, Michalec MD *et al.*; Identification of cathepsin C mutations in ethnically diverse Papillon Lefèvre syndrome patients. *Journal of Medical Genetics*, 2000; 37: 927–932.
6. D'Angelo A, Margiotta V, Ammatuma P, Sammartano F; Treatment of prepubertal periodontitis. A case report and discussion. *Journal of Clinical Periodontology* 1992; 19: 214–219.
7. Brown RS, Hays G, Flaitz CM, O'Neill PA, Abramovitch K, White RR; A possible late onset variation of Papillon–Lefèvre syndrome: report of 3 cases. *Journal of Periodontology* 1993; 64: 379–386.
8. Tinanoff N, Temprow P, Maderazo EG; Dentaltreatment of Papillon–Lefèvre syndrome: 15 year follow-up. *Journal of Clinical Periodontology*, 1995; 22: 609–612.
9. Yagmur A, Yilmaz G, Ertan U, IKizoglu E, Ozkasap S, Karacan C; Papillon–Lefèvre syndrome: A case report. *Int Pediatr* 2004;19224-5.
10. Janjua SA, Khachemoune A; Papillon-Lefèvre syndrome: case report and review of the literature. *Dermatol Online J.*, 2004; 10(1):13.
11. Galanter DR, Bradford S; Case report. Hyperkeratosis palmoplantaris and periodontosis: The Papillon-Lefèvre syndrome. *J Periodontol.*, 1969; 40(1): 40-47.
12. Hart TC, Hart PS, Michalec MD, Zhang Y, Firatli E, Van Dyke TE *et al.*; Haim-Munk syndrome and Papillon-Lefèvre syndrome are allelic mutations in cathepsin C. *Periodontol.*, 2000; 37: 88-94.
13. Newman MG, Takei HH, Carranza FA; *Clinical periodontology*. 9<sup>th</sup> edition, Pennsylvania, Saunders, 2003.
14. Reys VO, King-Ismael D, Abad-Venida L; Papillon Lefevre syndrome. *Int J Dermatol.*,1998; 37: 268-270.
15. Angel TA, Hsu S, Kornbleuth SI, Kornbleuth J, Kramer EM; Papillon Lefevre syndrome: A case report of four affected siblings. *J Am Acad Dermatol.*, 2002; 46: S8-S10.
16. Bergman R, Friedman-Birnbaum R; Papillon-Lefèvre syndrome: a study of the long-term clinical course of recurrent pyogenic infections and the effects of etretinate treatment. *Br J Dermatol.*, 1988;119:731–736.
17. Haneke E; The Papillon-Lefèvre syndrome: keratosis palmoplantaris with periodontopathy. *Human Genetics*, 1979; 51: 1–35.
18. Wara-Aswapati N, Lertsirivorakul J, Nagasawa T, Kawashima Y, Ishikawa I; Papillon–Lefèvre syndrome: Serum immunoglobulins G (IgG) subclass antibody response to periodontopathic bacteria. A case report. *J Periodontol.*, 2001; 72: 1747–1754.
19. Tinanoff N, Temprow P, Maderazo EG; Dental treatment of Papillon–Lefèvre syndrome: 15 year followup. *J Clin Periodontol.*, 1995; 22: 609–612.
20. Stabholz A, Taichman S, Soskolne WA; Occurrence of actinobacillus actinomycetemcomitans and anti-leukotoxin antibodies in some members of an extended family affected by Papiilon-Lefèvre syndrome. *J Periodontol.*, 1995; 66: 653–657.
21. Rüdiger S, Petersilka, G, Flemming TF; Combined systemic and local antimicrobiol therapy of periodontal disease in Papillon–Lefevre syndrome. *Journal of Clinical Periodontology*, 1999; 26: 847–854.
22. Wiebe CB, Häkkinen L, Putkins EE, Walsh P, Larjava HS; Successful periodontal maintenance of a case with Papillon–Lefèvre syndrome: 12-year follow-up and review of the literature. *Journal of Periodontology*, 2001; 72: 824–830.
23. Glenwright HD, Rock WP; Papillon - Lefevre syndrome. A discussion of etiology and a case report. *Br Dent J.*, 1990;168: 27-29.
24. Tinanoff N, Temprow P, Maderazo EG; Dental treatment of Papillon-Lefevre syndrome: 15 year follow-up. *J Clin Periodontol.*, 1995; 22: 609-612.
25. Gelmetti C, Nazzaro V, Cerri D, Fracasso L; Long term preservation of permanent teeth in a patient with Papillon-Lefevre syndrome treated with etretinate. *Pediatr Dermatol.*, 1989; 6: 222-225.