

Research Article

A study on two important environmental services of urban trees to disseminate the economic importance of trees to student community

P. Pachaiyappan, D. Ushalaya Raj

Institute of Advanced Study in Education, Saidapet, Chennai – 600 015, Tamil Nadu, India

*Corresponding author

P. Pachaiyappan

Email: edutechgreenfa@gmail.com

Abstract: Trees provide innumerable ecosystem services in human-dominated urban environment. Forest disturbances as well as biomass enrichments are tightly linked with atmospheric carbon dioxide concentration. All trees ≥ 5 cm diameter at breast height (dbh) were inventoried from a one hectare area of the Cooum river bank (CRB), Chennai Metropolitan city (CMC), India. Both above and below ground biomass were estimated by widely accepted regression equations with DBH and wood density as inputs. A total of 710 trees belonged to 22 families, 41 genera and 47 species were recorded. Trees accumulated 86.02 Mg dry biomass and 43.01 Mg C in a hectare. Members of Mimosaceae dominated the CRB with 231 individuals. *Tamarindus indica* contributed more (11.744 Mg; 13.7%) to biomass. As to the families Ceasalpiniaceae, Mimosaceae and Papilionaceae altogether contributed 55.61 Mg (64.64%) to total biomass. Tree diameter class 31-45 cm contributed more (35.15 Mg; 40.86%) to total biomass. On average each tree achieved 0.47 ± 0.1 cm dbh growth yr^{-1} . In a year one hectare urban forest sequestered 3999.91 kg biomass and 1999.95 kg C. The data obtained through this study can be useful to educate the importance of urban trees to higher secondary school students and local community people in Chennai city.

Keywords: biomass storage; carbon storage, carbon dioxide absorption; higher secondary school students; human intervention; urban forest

INTRODUCTION

Urban forests and trees do numerable ecosystem services to urban environments. They reduce urban heat island (UHI) effects [1, 2]; pollution [3]; CO₂ concentration [4, 5]. By their autotrophic nature, they sequester carbon (C) thereby they act as carbon sink [7-9]. Furthermore, they also offer an opportunity for conducting research, which can be useful for a more comprehensive understanding of urban ecosystems in general [10].

To date, a very limited data has been made available on biomass storage and carbon sequestration potential of trees growing on the Cooum river bank (CRB) in Chennai metropolitan city (CMC), India. A major proportion of higher secondary school students and local community people have lesser knowledge on urban trees and its importance. Thus, the primary objective of this study was to determine C stockpile and sequestration potential of trees on CRB. Further, the data obtained through this study can be useful to educate the importance of urban trees to higher secondary school students and local community people in Chennai city.

METHODS

Study area

This study was conducted in a highly polluted, urbanized Cooum river bank, Chennai Metropolitan City, India. Chennai is one among the four metropolis of the Indian sub-continent and the capital city of a

southern state, Tamil Nadu. The city feels tropical dissymmetric climate, receives bulk of the rain-fall during north-east monsoon (September-December). Mean temperature and rainfall are 24-37 °C and 1300 mm. Area of the city is 174 km² and the human population is around five million [11]. East part of the city is bounded by the sea, Bay of Bengal; and remaining three sides are surrounded by land, Thiruvallur and Kanchipuram districts. Soil type of the river bed is alluvium; depth of the soil varies from 10 to 20 m thickness and is mostly granular in texture [12].

Field survey

One hundred 10m × 10m plots (Total area = 1 ha) were randomly laid across both northern (50 plots) as well as southern bank (50 plots) of the river. The tree survey was conducted during the month of January on 2011 and 2012. All trees ≥ 5 cm diameter at breast height (dbh) were measured, and tagged with consecutively numbered metal tags for further monitoring and re-assessments. For multi-stemmed individuals, the bole girth was measured separately, basal area calculated and summed. All inventoried species were identified to species level with the help of regional floras and checklist [13-15].

Above and below ground biomass

Above ground biomass was quantified by regression formula of Chave et al., [16]; $[(AGB)_{est} = p \times \exp(-0.667 + 1.784 \text{ LN}(D) + 0.207(\text{LN}(D))^2 - 0.0281(\text{LN}(D))^3)]$; where, -0.667, 1.784, 0.207 and -

0.028 are constants; D = trunk diameter at breast height (cm); LN = Natural logarithm; p = oven-dry wood specific gravity/wood density (g cm^{-3}). The tree allometric relationship is viable, reliably estimate AGB in tropical dry forests around the world. Data on tree wood density data was retrieved from *Global wood density database* [17]. Below ground biomass (BGB) was calculated by regression equation of Cairns *et al.* [18]; $\text{BBD} = \text{Exp} [-1.0587 + 0.8836 \times \text{LN} (\text{ABD})]$ Where, -1.0587, 0.8836 are constants; BBD is below ground biomass density (dry t ha^{-1}); LN = Natural logarithm; ABD = above ground biomass density (dry Mg ha^{-1}).

Quantification of C and CO₂ sequestration

Carbon sequestration values were converted to CO₂ by multiplying with 3.67, the ratio of molecular weights of CO₂ to C (as in [5]). Biomass values were multiplied by 0.50 to get carbon storage value of trees (as in [7]). Stem diameter growth, biomass accumulation yr^{-1} were directly estimated by the difference between survey x and $x+1$. C sequestration potential was estimated by the difference of C storage of trees between year x and $x+1$.

RESULTS

Tree diversity and forest stand

A total of 47 species belonged to 41 genera and 22 families were inventoried from one hectare area of CRB (Table 1). *Muntingia calabura* and *Prosopis juliflora* were dominated the CRB with 139, 137 individuals, respectively, followed by *Ricinus communis* (99) and *Leucaena leucocephala* (72), (Table 2). As to the families, Mimosaceae was dominated with 8 species followed by Caesalpiniaceae (7), Moraceae (4), Bignoniaceae and Myrtaceae (3 each), Anacardiaceae, Annonaceae, Meliaceae, Papilionaceae and Sapotaceae (2 species each), whereas 12 families which include Apocynaceae, Arecaceae and Bombacaceae etc. were represented by just single species (Table 3). Tree stand density and basal area were 710 stem ha^{-1} and 15.10 m^2 , respectively.

Biomass and carbon storage

According to the first survey, study area stored 86.018 Mg biomass in its trees (AGB = 63.66 Mg; BGB = 22.36). Cumulatively, top five species (*Tamarindus indica*, *Prosopis juliflora*, *Pongamia pinnata*, *Albizia saman* and *Peltophorum pterocarpum*) with 197 individuals contributed 45.50 Mg (49.4%) to the total biomass (Table 2). As to the families, Caesalpiniaceae accumulated high biomass 22.06 Mg (25.64%) followed by Mimosaceae (21.91; 25.47%) and Papilionaceae (11.64; 13.53%), (Table 3). Small diameter classes 0-7 cm dbh and 8-15 dominated the CRB; they represented by 223 and 295 individuals, respectively. As to the total biomass, the dbh class 31-45 cm contributed a large amount (35.15 Mg; 40.86%) followed by 16-30 (34.48 Mg; 40.1%) and 8-15 cm dbh (15.30 Mg; 17.79%), (Fig. 1). Large (45+ cm) trees stored approximately 160 times more C ($1225 \pm 105.93 \text{ kg/tree}$) when compared to low diameter class ($7.56 \pm 0.32 \text{ kg/tree}$). Evergreen species contributed more (50.295 Mg; 58.47%) to total biomass than deciduous species (35.722 Mg; 41.53%). Native trees accumulated more biomass (52.24 Mg; 60.74%) compared to introduced trees (33.78; 39.26%). Carbon stored in a tree ranged from a low of 3.78 ± 0.16 (0-7 dbh) to a high of $612.6 \pm 52.96 \text{ C kg/tree}$ (45+ dbh).

Stem horizontal growth

On an average, each tree attained $0.47 \pm 0.1 \text{ cm dbh yr}^{-1}$. Large trees achieved more growth ($0.68 \pm 0.2 \text{ cm dbh}$) than small trees ($0.48 \pm 0.1 \text{ cm dbh}$).

Biomass and carbon sequestration

In a year, inventoried trees accumulated 3999.91 kg biomass ha^{-1} . On an average, each tree accumulated 5.63 kg biomass yr^{-1} . Biomass accumulation/tree/year ranged from 1.4 to 18.41 kg, respectively for small (0-7 cm dbh), and large (31-45 cm dbh) trees (Table 4).

Trees sequestered 1999.95 kg C $\text{yr}^{-1} \text{ ha}^{-1}$. Each tree sequestered 2.82 kg C yr^{-1} . C sequestration ranged from 0.7 ± 0.02 to $9.21 \pm 0.86 \text{ kg/tree/year}$. Roughly, large trees (45+ dbh) sequestered 10 times more C yr^{-1} than small trees (Table 4).

Table 1. Summary of tree inventory ($\geq 5 \text{ cm dbh}$) on Cooum river bank, Chennai, India

Variable	Value
Species richness	47
Number of genera	41
Number of families	22
Tree density (no. ha^{-1})	710
Stand basal area ($\text{m}^2 \text{ ha}^{-1}$)	15.10
Biomass storage (Mg ha^{-1})	86.02
Carbon storage (Mg ha^{-1})	43.01

Table 2. Binomial, family, density, physiognomy, status and biomass storage of trees in Cooum river bank, Chennai, India

Binomial	Family	Density	Physiognomy	Status	Biomass storage, Mg (%)
<i>Muntingia calabura</i>	Elaeocarpaceae	139	Evergreen	Introduced	1.648 (1.92)
<i>Prosopis juliflora</i>	Mimosaceae	137	Evergreen	Introduced	10.119 (11.8)
<i>Ricinus communis</i>	Euphorbiaceae	99	Evergreen	Native	0.373 (0.43)
<i>Leucaena leucocephala</i>	Mimosaceae	72	Deciduous	Introduced	1.484 (1.73)
<i>Pongamia pinnata</i>	Papilionaceae	26	Deciduous	Native	8.204 (9.54)
<i>Azadirachta indica</i>	Meliaceae	24	Deciduous	Native	4.087 (4.75)
<i>Thespesia populnea</i>	Malvaceae	19	Evergreen	Native	0.814 (0.95)
<i>Erythrina variegata</i>	Papilionaceae	14	Deciduous	Native	3.430 (3.99)
<i>Cordia obliqua</i>	Boraginaceae	13	Evergreen	Native	1.257 (1.46)
<i>Peltophorum pterocarpum</i>	Caesalpiniaceae	13	Evergreen	Native	5.838 (6.69)
<i>Albizia saman</i>	Mimosaceae	12	Deciduous	Introduced	6.597 (7.67)
<i>Guazuma ulmifolia</i>	Sterculiaceae	11	Evergreen	Introduced	3.059 (3.56)
<i>Ficus hispida</i>	Moraceae	10	Evergreen	Native	0.255 (0.3)
<i>Cassia fistula</i>	Caesalpiniaceae	10	Deciduous	Native	0.926 (1.08)
<i>Annona squamosa</i>	Annonaceae	9	Evergreen	Introduced	0.363 (0.42)
<i>Tamarindus indica</i>	Caesalpiniaceae	9	Evergreen	Native	11.744 (13.7)
<i>Morinda coreia</i>	Rubiaceae	8	Evergreen	Native	0.780 (0.91)
<i>Delonix regia</i>	Caesalpiniaceae	7	Deciduous	Introduced	1.763 (2.05)
<i>Terminalia catappa</i>	Combretaceae	6	Deciduous	Native	2.250 (2.62)
<i>Ficus religiosa</i>	Moraceae	6	Deciduous	Native	2.465 (2.87)
<i>Polyalthia longifolia</i>	Annonaceae	5	Evergreen	Introduced	1.905 (2.22)
<i>Caesalpinia coriaria</i>	Caesalpiniaceae	4	Evergreen	Introduced	0.999 (1.16)
<i>Moringa pterygosperma</i>	Moringaceae	4	Deciduous	Native	0.096 (0.11)
<i>Spathodea companulata</i>	Bignoniaceae	4	Deciduous	Introduced	1.312 (1.53)
<i>Lannea coromandelica</i>	Anacardiaceae	3	Deciduous	Native	0.792 (0.92)
<i>Psidium guajava</i>	Myrtaceae	3	Evergreen	Introduced	0.171 (0.2)
<i>Acacia auriculiformis</i>	Mimosaceae	3	Evergreen	Introduced	0.285 (0.33)
<i>Melia azedarach</i>	Meliaceae	3	Deciduous	Native	0.306 (0.36)
<i>Ziziphus mauritiana</i>	Rhamnaceae	3	Evergreen	Native	0.231 (0.27)
<i>Cassia roxburghii</i>	Caesalpiniaceae	3	Evergreen	Native	0.183 (0.21)
<i>Ficus racemosa</i>	Moraceae	3	Evergreen	Native	0.363 (0.42)
<i>Tabebuia rosea</i>	Bignoniaceae	3	Evergreen	Introduced	1.083 (1.26)
<i>Adenanthera pavonina</i>	Mimosaceae	3	Deciduous	Native	2.090 (2.43)
<i>Bombax ceiba</i>	Bombacaceae	3	Deciduous	Native	0.905 (1.05)
<i>Mangifera indica</i>	Anacardiaceae	2	Evergreen	Native	1.091 (1.27)
<i>Manilkara zapota</i>	Sapotaceae	2	Evergreen	Introduced	0.078 (0.09)
<i>Cassia siamea</i>	Caesalpiniaceae	2	Evergreen	Native	0.605 (0.7)
<i>Albizia lebbeck</i>	Mimosaceae	2	Deciduous	Native	0.761 (0.89)
<i>Ficus benghalensis</i>	Moraceae	2	Evergreen	Native	1.018 (1.18)
<i>Millingtonia hortensis</i>	Bignoniaceae	2	Evergreen	Introduced	1.329 (1.54)
<i>Wrightia tinctoria</i>	Apocynaceae	1	Deciduous	Native	0.032 (0.04)
<i>Pithecellobium dulce</i>	Mimosaceae	1	Deciduous	Introduced	0.055 (0.06)
<i>Madhuca longifolia</i>	Sapotaceae	1	Deciduous	Native	0.163 (0.19)
<i>Feronia elephantum</i>	Rutaceae	1	Evergreen	Native	0.281 (0.33)
<i>Enterolobium cyclocarpum</i>	Mimosaceae	1	Deciduous	Introduced	0.520 (0.61)
<i>Syzygium cumini</i>	Myrtaceae	1	Evergreen	Native	0.899 (1.05)
<i>Eucalyptus globulus</i>	Myrtaceae	1	Evergreen	Introduced	1.002 (1.17)
Total		710	-	-	86.018 (100)

Table 3. Contribution of families to total biomass in Cooum river bank, Chennai Metropolitan city, India

Family	Density	Biomass (Mg)	Contribution to total biomass (%)
Caesalpinaceae	48	22.058	25.64
Mimosaceae	231	21.913	25.47
Papilionaceae	39	11.635	13.53
Meliaceae	27	4.394	5.11
Moraceae	21	4.102	4.77
Bignoniaceae	9	3.724	4.33
Sterculiaceae	11	3.059	3.56
Annonaceae	14	2.268	2.64
Combretaceae	6	2.25	2.62
Myrtaceae	5	2.072	2.41
Anacardiaceae	5	1.883	2.19
Elaeocarpaceae	139	1.648	1.91
Boraginaceae	13	1.257	1.46
Bombacaceae	3	0.905	1.05
Malvaceae	19	0.814	0.95
Rubiaceae	8	0.78	0.91
Euphorbiaceae	100	0.373	0.42
Rutaceae	1	0.281	0.33
Sapotaceae	3	0.241	0.28
Rhamnaceae	3	0.231	0.27
Moringaceae	3	0.096	0.11
Apocynaceae	1	0.032	0.04
Total	710	86.018	100

Table 4. Biomass, carbon storage and sequestration of trees in Cooum River bank, Chennai Metropolitan city, India


DBH	Density	Biomass storage (Mean \pm SE)	Carbon storage (Mean \pm SE)	Carbon sequestration yr ⁻¹ (Mean \pm SE)
0-7	223	7.56 \pm 0.32	3.78 \pm 0.16	0.7 \pm 0.01
8-15	295	41.49 \pm 1.57	20.74 \pm 0.79	2.14 \pm 0.07
16-30	138	249.88 \pm 12.89	124.94 \pm 6.44	5.07 \pm 0.20
31-45	52	676.09 \pm 48.87	338.04 \pm 29.43	9.21 \pm 0.86
45+	2	1225.2 \pm 105.92	612.6 \pm 52.96	8.22 \pm 0.33
Mean		121.15	60.57	2.82
Total (Mg ha ⁻¹)		86.018	43.01	1.99

Table 5. Carbon storage (Mg ha⁻¹) in Cooum river bank and other urban forests of the world

Country	City	Carbon storage Mg ha ⁻¹ (mean)	Reference
India	Chennai	43.09	Present study
China	Beijing	45.39	[23]
Germany	Liepig	11.00	[26]
India	Pune	54.87	[33]
Korea	4 cities	11.9	[31]
Nepal	Riverine forests	80.47	[36]
Taiwan	Chiayi	13.5	[32]
USA	Shorewood	22.80	[24]
USA	Michigan	40.5	[28]
USA	Los Angeles	10.38	[22]
USA	Oakland	11.00	[29]
USA	3 cities	99.33	[34]
USA	All cities	66.69	[35]
USA	10 cities	21.07	[30]
USA	48 cities	25.10	[27, 30]

Table 6. Money value of carbon storage and sequestration services of trees

DBH classes	C storage (tree/year) as on January 2012	Value (\$)	C sequestration (tree/yr)	Value (\$)
0-7	3.78	0.25	0.7	0.05
8-15	20.74	1.37	2.14	0.14
16-30	124.94	8.25	5.07	0.33
31-45	338.04	22.31	9.21	0.61
45+	612.6	40.43	8.22	0.54

**Fig. 1. Contribution of DBH classes to total dry biomass in Cooum river bank, Chennai, India.**

DISCUSSION

Tree density

Tree density ha^{-1} is two to five folds higher than those of urban forests of Oakland (111.9 trees ha^{-1} ; [19]); Modesto, California (61; [20]); ten cities of USA (mean=147, range, 36 to 276; [7]); Sacramento, USA (73; [21]), Los Angeles (49; [22]); and Beijing, China (79 trees ha^{-1} ; [23]).

Existence of more number of small trees (<45 cm dbh; 708 trees) could be reasoned for high stem density on CRB. Relatively smaller diameter trees such as *Leucaena leucophloea*, *Muntingia calabura*, *Prosopis juliflora*, *Ricinus communis* were abundant and growing luxuriantly on CRB, altogether they have represented by 448 (63.10%) individuals. Furthermore, researchers of other countries studied entire urban forests; we have studied only a small portion of urban forest on CRB. More studies with large study areas are needed to reveal the real picture of the CRB. Approximately, 27% of trees had ≥ 15 cm dbh in our study area; it is less when compared to other urban forests of the world, such as Oakland, California (39%; [19]); Shorewood, Wisconsin (33%; [24]); and Los Angeles of USA (60%; [22]). However, studied forest of Chennai is superior to urban forests of Chicago, Cook and DuPage Counties of USA (23%), [25].

Carbon storage

C storage ($43.09 \text{ Mg C ha}^{-1}$) is greater than those reported from many urban forests of the world, such as Germany [26]; Oakland, Shorewood, Michigan, Los Angeles and several cities of the USA [22, 24, 27, 28, 29, 30]; Korea [31]; and Taiwan [32], (**Table 5**). Also lesser than many urban forests such as Pune, India [33]; Beijing, China [23]; Cook and DuPage counties, and Chicago of USA [34]; cities of the USA [35]; and, riverine forests of Nepal [36] (**Table 4**). Less C storage in CRB could be due to the presence of large number of small trees (≤ 15 cm dbh) and absence of large trees ($50+$ cm dbh).

Stem radial growth

Our findings pertaining to mean stem growth (dbh cm)/tree/year is not agreed with the results of Jo and McPherson [4]; deVries [37] and Nowak [25] they estimated 1.1, 0.61, 0.90 cm dbh growth yr^{-1} respectively for urban trees of USA, central park of New Jersey; and three cities of USA. Presence of more large trees in those forests could be contributed to high dbh growth yr^{-1} . However, trees of CRB attained more growth yr^{-1} than urban trees of Indiana and Illinois of USA, 0.38 cm dbh growth yr^{-1} [38].

Carbon sequestration

Carbon sequestration potential $\text{kg ha}^{-1} \text{yr}^{-1}$ (1999 kg) is superior than those of four cities of Korea (530 to 800 Mg; [31]); Chiayi city of Taiwan (0.71 Mg; [32]); Michigan, USA (810 kg; [28]); Los Angeles, USA (642 kg; [22]); ten cities of USA (mean=747.5 kg $\text{C ha}^{-1} \text{yr}^{-1}$, range 210-1230 kg $\text{C ha}^{-1} \text{yr}^{-1}$; [7]), and as well as 48 cities of USA (mean=800 kg C ha^{-1} [27, 30]. However, low when compared to the Beijing's urban forest (2310 kg $\text{C ha}^{-1} \text{yr}^{-1}$; range 2610 to 6970 kg $\text{C ha}^{-1} \text{yr}^{-1}$; Yang *et al.*, 2005); and riverine forests of Nepal (3210 kg [36]. In this study, an individual tree sequestered approximately 40 to 60% less C than urban trees of three USA cities. Nowak [34] estimated a low of 1.0 kg (0-7 cm dbh) to a high of 92.7 kg (77+ cm dbh) of C sequestration/tree/year for urban trees. Higher stem radial growth yr^{-1} and occurrence of many large trees in urban forests of the USA could have contributed to more biomass and carbon sequestration. The money values of two environmental services namely carbon storage and sequestration have been provided in **Table 6**. The carbon storage value of trees in study site is about 2800 \$ (66 \$/tonne carbon) whereas annual carbon sequestration potential of trees is approximately 150 \$. With this simple calculation one can easily imagine the value of trees which are growing on the Cooum river bank in Chennai city. This study provides the money value of just two environmental services, with studies on all environmental services of trees (such as pollination, water percolation, conservation of soil, maintenance of water quality etc.) one can easily understand the real value of trees.

CONCLUSIONS

This study sheds light on understudied urban forests in Chennai Metropolitan City, India. Tree density is greater as well as lower than in urban forests elsewhere. Whereas, tree dry biomass storage, C storage and sequestration are comparable to other urban forest around the world. Conservation of trees on the CRB is important to protect associated faunal communities, they may play unique ecosystem services such as pollination, seed dispersal etc. The data obtained on carbon storage and sequestration of trees can be useful to educate the economic importance of trees to higher secondary school students and local community people in Chennai metropolitan city, India.

ACKNOWLEDGEMENTS

We thank Mr. M. Udayakumar, DST INSPIRE Fellow, Pachaiyappa's College, Chennai for his help and support during field study and manuscript preparation. We extend our thanks to the metropolitan authorities for their permission to carry out this work on the Cooum river bank.

REFERENCES

1. Akbari H, Rosenfeld AH, Taha H; Summer heat islands, urban trees, and white surfaces. *ASHRAE transactions*, 1990; 96: 1381–1388.
2. Akbari H, Pomerantz M, Taha, H; Cool surfaces and shade trees to reduce energy use and improve air quality in urban areas, *Solar Energy*, 2001; 70: 295–310.
3. Whitford V, Ennos AR, Handley JF; City form and natural process-indicators for the ecological performance of urban areas and their application to Merseyside, UK. *Landscape and Urban Planning*, 2001; 57: 91–103.
4. Jo HK, McPherson EG; Carbon storage and flux in urban residential greenspace. *Journal of Environmental Management*, 1995; 45: 109–33.
5. McPherson EG, Simpson JR; Carbon dioxide reduction through urban forestry: guidelines for professional and volunteer tree planters. General Technical Report PSW–171. USDA Forest Service, Pacific Southwest Research Station, Albany, USA, 1999: 237 pp.
6. Brack CL; Pollution mitigation and carbon sequestration in an urban forest. *Environmental Pollution*, 2002; 116: 195–200.
7. Nowak DJ, Crane DE; Carbon storage and sequestration by urban trees in the USA. *Environmental Pollution*, 2002; 116: 381–389.
8. Chen Y, Jim, CY; Assessment and valuation of the ecosystem services provided by urban forests, in: Carreiro, M.M., Song, Y.C., Wu, J. (Eds.), *Ecology, Planning and Management of Urban Forests, International Perspectives*. Springer, Netherlands, 2008: 53–83.
9. Pickett STA, Cadenasso ML, Grove JM, Groffman PM, Band, LE, Boone CG *et al.*; Beyond urban legends: An emerging frame work of urban ecology, as illustrated by the Baltimore Ecosystem Study. *Bioscience*, 2008; 58: 139–150.
10. McDonnell MJ, Pickett S, Groffman P, Bohlen P, Pouyat R, Zipperer WC, *et al.*; Ecosystem process along urban-to-rural gradients. *Urban Ecosystem*, 1997; 1: 21–36.
11. Population Census India, 2011; Retrieved on 1st March, 2012. Available from <http://www.census2011.co.in/census/district/21-chennai.html>.
12. Giridharan L, Venugopal T, Jayaprakash M; Identification and evaluation of hydrogeochemical processes on river Cooum, South India. *Environmental Monitoring and Management*, 2010; 162: 277–289.
13. Gamble JS, Fischer CEC; *Flora of the Presidency of Madras*. 3 volumes. Adlard and Son Limited, London, 1921-35.
14. Mayuranathan, PV; The flowering plants of Madras city and its immediate neighbourhood. *Bulletin of the Madras government museum*. Government Press, Madras, India, 1929.

15. Udayakumar M, Ayyanar M, Sekar T; Angiosperms, Pachaiyappa's College, Chennai, Tamil Nadu, India. Check List, 2011; 7: 37–48.
16. Chave J, Andalo C, Brown S, Cairns MA, Chambers JQ, Eamus D *et al.*; Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, 2005; 145: 87–99.
17. Zanne AE, Lopez-Gonzalez G, Coomes DA, Ilic J, Jansen S, Lewis SL *et al.*; Global wood density database. Dryad. Identifier: <http://hdl.handle.net/10255/dryad.235>, 2009.
18. Cairns MA, Brown S, Helmer EH, Baumgardner GA; Root biomass allocation in the world's upland forests. *Oecologia*, 1997; 111: 1–11.
19. Nowak DJ; Urban forest development and structure: analysis of Oakland, California. Ph.D. Dissertation. University of California, Berkeley, 1991.
20. McPherson EG, Simpson JR, Peper PJ, Xiao Q; Benefit–Cost Analysis of Modesto's Municipal Urban Forest. *Journal of Arboriculture*, 1999; 25: 235–248.
21. Birdsey RA, Heath LS; Carbon changes in U.S. forests, in: Joyce, L.A. (Eds.), *Climate Change and the Productivity of America's Forests*, General Technical Report RM-GTR-271. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Ft Collins, CO, USA, 1995: 56–70.
22. Nowak DJ, Hoehn RE, Crane DE, Weller R, Davila A; Assessing Urban Forest Effects and Values: Los Angeles' Urban Forest. Resource Bulletin NRS-47. United States Department of Agriculture. USDA Forest Service, Northern Research Station, USA, 2011: 1-35.
23. Yang J, McBride J, Zhou J, Sun Z; The urban forest in Beijing and its role in air pollution reduction. *Urban Forestry and Urban Greening*, 2005; 3: 65–78.
24. Dorney JR, Guntenspergen GR, Keough JR, Stearns F; Composition and structure of an urban woody plant community. *Urban Ecology*, 1984; 8: 69–90.
25. Nowak DJ; 1994a. Urban Forest structure: The state of Chicago's urban forest, in: McPherson, E.G., Nowak, D.J., Rowntree, R.A. (Eds.). *Chicago's urban forest ecosystem: Results of the Chicago Urban Forest Climate Project*. General Technical Report, NE-186. U.S. Department of Agriculture, Forest Service, Radnor, PA, 1994a: 3–18.
26. Strohbach MW, Haase D; Above-ground carbon storage by urban trees of Leipzig, Germany: Analysis of patterns in a European city. *Landscape and Urban Planning*, 2012; 104: 95–104.
27. Dwyer JF, Nowak DJ, Noble MH, Sisinni SM; *Connecting People with Ecosystems in 21st Century: An Assessment of our Nation's Urban Forests*. General Technical Report PNW-GTR-490). US Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR, 2000: 1-490
28. MacFarlane DW; Potential availability of urban wood biomass in Michigan: implications for energy production, carbon sequestration and sustainable forest management in the U.S.A. *Biomass and Bioenergy*, 2009; 33: 628–634.
29. Nowak DJ; Historical vegetation change in Oakland and its implications for urban forest management. *Journal of Arboriculture*, 1993; 19: 313–319.
30. Nowak DJ, Noble MH, Sisinni SM, Dwyer JF; Assessing the US urban forest resource. *Journal of Forestry*, 2001; 99: 37–42.
31. Jo HK; Impacts of urban green space on offsetting carbon emissions from middle Korea. *Journal of Environmental Management*, 2002; 64: 115–126.
32. Huang HH, Huang HW, Chang SH; The study of ecological benefits of Chiayi city, in: Schrenk, M., Popovich, V. V., Engelke, D., & Elisei, P. (Eds.), *Proceedings REAL CORP 2009*, Tagungsband, Taiwan, 2009: 535–540.
33. Waran A; Carbon sequestration potential of trees in and around Pune city. Master's thesis, University of Pune, India; 2001.
34. Nowak DJ; Atmospheric carbon dioxide reduction by Chicago's urban forest, in: McPherson, E.G., Nowak, D.J., Rowntree, R.A. (Eds.), *Chicago's Urban Forest Ecosystem: Results of the Chicago Urban Forest Climate Project*. General Technical Report, NE-186, US Department of Agriculture, Forest Service, Radnor, PA, 1994b; 83–94.
35. Rowntree R, Nowak DJ; Quantifying the role of urban forests in removing atmospheric carbon dioxide. *Journal of Arboriculture*, 1991; 17: 269–75.
36. Baral SK, Malla R, Ranabhat S; Above-ground carbon stock assessment in different forest types of Nepal. *Banko Janakari*, 2009; 19: 10–14.
37. de Vries RE; A preliminary investigation of the growth and longevity of trees in Central Park. MS thesis, Rutgers University, New Brunswick, New Jersey, 1987.
38. Smith WB, Shifley SR; Diameter Growth, Survival, and Volume Estimates for Trees in Indiana and Illinois. Research Paper NC-257. U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station, St. Paul, MN, 1984; 1-10.