

Research Article

Study of Fungi to Evaluate Their Potential for Metabolites Production under Submerge Condition by Fermentation Technology

Chouhan Pawan Kumar*, Nigam Shashwat

Arts and Science Govt. PG College, Ratlam Dist. - 457001, Madhya Pradesh, India

*Corresponding author

Pawan Kumar Chouhan

Email: pawan_chouhan30@yahoo.com

Abstract: *Aspergillus* is filamentous achlorophyllous fungi and was isolated from agriculture crop fields of Mandasaur then identified by microscopic method using cotton blue and lacto phenol. Biomass and metabolites i.e. protein and enzyme were produced by *Aspergillus* under submerge fermentation technology during various incubations. After observation it was found that maximum biomass, protein and invertase was shown as 1.10gm, 4.575mg/ml and 21.80µg/ml after 10 and two days incubation time respectively by *Aspergillus* spp. These metabolites were estimated using standard protocols.

Keywords: *Aspergillus* spp., Biomass, Dinitro salicylic acid Submerge Fermentation technology, Invertase, Incubation time, Lowry reagent, Protein.

INTRODUCTION

Generally fungi are filamentous non-photosynthetic organism and nutritionally multiplied on dead organic materials or living cells as saprophyte or parasite. In growth phase, fungi produced their primary and secondary metabolites and available in their growth medium [9]. Protein is a nitrogenous compound and made up of 20 different amino acids. All living organisms required protein for their growth. Generally fermentation technology was used for commercial protein production using many microorganisms including fungi, yeasts, bacteria and actinomycetes. Proteins are nitrogen or amino acid supplements and most of the enzymes are made of proteins. Besides energy source, protein required for fungi, bacteria, actinomycetes and other unicellular and multicellular cells for their membrane and enzyme synthesis [10].

Filamentous fungi have been widely used in the fermentation industry as it becomes a principal source of protein, enzymes and other metabolites. Therefore, fungi have been widely investigated by various researchers due to low cost and high productivity which attracted many other researchers to improve fungal strains by molecular techniques and also bioprocess [5, 16].

Many cells membrane made of lipoprotein, glycoprotein and other compounds where protein associated with them. Therefore, proteins are produced by fermentation technology and used in a lot of industries such chemical, detergent and food as nutrient supplements for human diet [24, 31].

Invertase catalyze the hydrolysis of sucrose to produce an equimolar mixture of d-glucose and d-fructose at concentrations lower than 10% sucrose, thus

making these enzymes suitable for biotechnological applications [40]. Invertases (EC.3.2.1.26) are members of glycoside hydrolases family [1] and have numerous beneficial favorable functional properties [17].

The maximum rate of invertase enzyme production in presence of sucrose by *A. niger* during submerged culture condition was found as 3.67 U/L/h at 12.5 g/L while in the case of glucose-sucrose mixture; it was 13.95 U/L/h at 25 g/L. It was observed that the enzyme yield was 1.25 times more in presence of sucrose than with combined action of glucose-sucrose [38].

Invertases have been characterized in many filamentous fungi such as *Penicillium*, *Neurospora* and *Aspergillus* spp. [4, 19-20, 36]. The enzymatic activity of invertase has been characterized in other microorganisms i.e. *Thermomyces lanuginosus* [8], *Candida utilis* [7], *Penicillium chrysogenum* [33], *Saccharomyces cerevisiae* and *S. carlsbergensis* [21, 23]. *Saccharomyces cerevisiae* synthesizes two invertases i.e. a glycosylated periplasmic protein and a cytosolic non glycosylated protein [37, 46].

Dworschack and Wickerham [14] demonstrated that extracellular invertase was also produced by yeast and some physiological factors influence the production of extracellular and intracellular invertase by *Saccharomyces uvarum*. Invertase production by *P. chrysogenum* in penicillin fermentation broth was studied by Damele *et al.* [11] and reported that invertase produced extracellularly.

Elorza *et al.* [15], Trimble and Maley [43] reported that *Saccharomyces cerevisiae*-136 produced external invertase in culture filtrate supplemented with

50% sucrose and maltose. Olutiola and Cole [34] described that an extracellular invertase production in *Aspergillus flavus* induced by sucrose containing liquid medium. The biosynthesis of invertase was repressed when repressive hexose sugars (glucose or fructose) were added to sucrose metabolizing cells.

Vainstein and Peberdy [45] investigated that *Aspergillus nidulans* produced an extracellular invertase when cultured on a medium containing sucrose or raffinose and was found maximal on medium containing sucrose after 15 hrs of incubation at 28°C. The amount of invertase in the culture medium was declined after this incubation time.

Rubio and Maldonado [39] studied that invertase produced by *Aspergillus niger* strain showed the maximum enzyme activity at 60°C temperature and pH 5.0. Medium containing glucose (0.1 M), fructose (0.5 M) and sucrose (1.2 M) as compared to that with only sucrose (1.8 M) solutions produced 3-fold more oligosaccharides at pH 7.5 [42].

Herwig *et al.* [21] reported that *S. cerevisiae* invertase as hydrolyzing capacity of sucrose and was controlled by the presence of hexoses such as glucose and fructose in medium. Commercially, invertase is produced on large scale primarily by using submerged fermentation. The well-known methods for production of invertase are submerged and solid state fermentation [19-20, 37]. The conventional method for the production of invert sugar involves acid hydrolysis of sucrose and has low conversion efficiency, high energy consumption and thus high cost of production due to acid hydrolysis whereas microbial invertase splits sucrose into glucose and fructose (sugar syrup) with 100% conversion efficiency [28].

The product obtained by invertase has the advantage of being colorless as compared to that obtained through acid hydrolysis [3, 6, 12, 27]. Kotwal and Shankar [26] studied the importance of invertase for the sucrose hydrolysis that yielded an equimolar mixture of glucose and fructose called as inverted syrup, also commonly used in beverage and food industries. Robledo-Olivo *et al.* [38] studied invertase production by *Aspergillus niger* in submerged culture using different concentrations of glucose and sucrose and found that the initial concentration of sucrose was increased from 6.25 to 50 g/L then a higher biomass production (6.1 g/L) was achieved. The biomass production was increased four times more when a glucose-sucrose combination was used as substrate (26.31 g/L).

Uma *et al.* [44] produced high levels of invertase in *A. flavus* under optimized culture conditions on fourth day of incubation at an optimum pH (5.0), temperature (30°C), and inoculum size (3%) in Czapek Dox medium using fruit peel waste as a substrate by submerged fermentation and also obtained

improved enzyme when nutritional factors such as sucrose and yeast extract were added into the medium.

Invertase is extensively used in confectionaries, food industries and in pharmaceuticals [4]. Invertase is used for the inversion of sucrose in the preparation of invert sugar and syrup. It is one of the most widely used enzymes in food industry where fructose is preferred than sucrose especially in the preparation of jams and candies, because it is sweeter and does not crystallize easily [2]. The objectives of the present study are to produce fungal biomass, total protein and invertase enzyme during various incubations using sucrose and evaluated by standard protocols.

MATERIAL AND METHODS

Soil sample collection

Five to six soil samples were collected from agricultural crop fields around Ratlam (23.32°N and 75.07°E Longitude) in Malwa region, Madhya Pradesh, India and were mix to form one composite sample. Soil sample was crushed using Mortar and Pestle to clean by removing rocks particles and plant debris.

Isolation of fungi

Fungi were isolated from soil by serial dilution technique [13, 22] using 10 gm soil and were dissolved in 0.85% saline water in Erlenmeyer conical flask (500 ml capacity) having 95 ml saline water and beads then mix properly by shaking, mark as 10⁻¹ dilution. Obtain 10⁻³ dilution by serially transferred 10 ml soil sample from 10⁻¹ dilution to conical flasks having 90 ml saline water. 0.1ml sample was transferred from 10⁻³ dilution to sterilize Petri dish and poured melted Potato dextrose agar (PDA) medium. Plate was shaking gently to mix sample and medium then left at room temperature to solidify medium. The solidified plate was incubated at 30°C for 5 days. After that plate was used for observation.

After incubation period, few spores were taken by wire loop and streaked on PDA medium plate by zigzag method and incubated at 30°C for 5 day. Colonies were observed by light microscope using 10 and 40X objective lens. Pure colonies were transferred to PDA plate and used for identification (Fig.- 1).

Identification of fungi

After purification prepared microscopic slide for each pure culture. Fungal culture was taken from colony by forceps on clean glass slide and separated by needles. Culture was stain with Cotton blue and lacto phenol and covered with glass cover slip then observed in light microscope using low power (10X) and high power (40X) objective lens. Morphological characteristics of fungal culture were identified using laboratory manual for introductory mycology [18, 35, 41]. After purification and identification spores of pure colonies was transferred to PDA slant and incubate at

30°C for 5 days then stored in refrigerator at 4°C for 3-6 month.

Spore count

Sterilized water (5.0 ml) was introduced to culture slant and scraped by nichrome wire loop then 0.1ml spore suspension was used to count spore using heamocytometer.

Biomass production

Spores of *Aspergillus* spp. (1 ml) were transferred to steam sterilized production medium (Table- 1) and incubated at 30°C for 2, 4, 6, 8 and 10 days under submerged condition.

Estimation of dry biomass

After incubation period, biomass was harvested by filtration method using Whatman No. 1 filter paper and dried in hot air oven at 60°C for 24 hrs then weigh using electronic balance (Fig.- 2).

Protein production

Poured 5 ml sterilized water in culture tube and then spores were scraped by using nichrome wire loop. One milliliter spore suspension was transferred to five Erlenmeyer conical flasks having 100 ml production medium (Table- 1) and incubated at 30°C for 2, 4, 6, 8 and 10 days under submerged condition for fermentation. After incubation time, culture filtrate was obtained by filtration using Whatman No. 1 filter paper and stored in refrigerator fro protein and enzyme estimation (Fig.- 3).

Estimation of total protein by Lowry method

Lowry reagent C mixes with 0.2ml of culture filtrate in glass test tubes [42]. The tubes were incubated at 37°C for 20 min in water bath then cooled and added and 0.2 ml Folin phenol. The tubes were again incubated at room temperature for 30 min and recorded color density at 660 nm by Spectrophotometer. The absorbance difference was calculated using standard curve of protein (BSA 1 mg /ml).

Enzyme production

Spores of selected fungi i.e. *Aspergillus* spp. were transferred to PDA slants and incubated at 30°C for five days. Five milliliter distilled water was introduced to culture slant and scraped by wire loop. One milliliter spore suspension was transferred to five Erlenmeyer conical flasks having 100 ml production medium (Table- 1) and incubated at 30°C for 2, 4, 6, 8 and 10 days under submerged condition for fermentation. After incubation time, culture filtrate was obtained by filtration using Whatman No. 1 filter paper and stored in refrigerator for enzyme estimation.

Enzyme estimation by DNS method

Enzyme activity was assayed using sucrose as substrate [25]. The assay was performed on the basis of red brown color produced by DNS reagent [30]. In this

assay, 1 ml of 2 mM sucrose solution in 0.05 mM sodium acetate buffer of pH 5 was taken in glass test tube and added 0.5 ml culture filtrate then mixed by using cyclomixer. All test tubes with reaction mixer were incubated in water bath at 30°C for 15min then added 2 ml DNS reagent (DNS- 10 g Phenol- 2 g Sodium succinate- 0.5 g, sodium hydroxide- 10 g and Distilled water- 1 L) to stopped enzyme - substrate reaction and again incubated for 20 min at boiling water bath to developed red brown color. All test tubes were takeout from water bath and cooled then added 1 ml of 1.0% Rochella salt (Sodium potassium tartrate- 10 g and Distilled water- 1 L). All test tubes were left at room temperature for 20 min to stabilized red brown color. Color density was recorded by spectrophotometer at 550 nm using spectrophotometer and compared with standard curve of glucose (1 mg/ml).

RESULTS AND DISCUSSION

The biomass, protein and invertase estimation of various incubation extract was evaluated using standard protocols and have found following estimations given in Table- 2 to 4.

Results obtained to evaluate total dry biomass are shown in Table- 2 and Fig.- 4 that dry weight of biomass was increased from 0.45 to 1.10 g with incubation time. Further Table 2 shows that 0.45 and 0.80 g biomass produced after 2 and 4 days incubation time and was increased exponentially as 1.10 gm after 10 days incubation period. Figure 4 reveals that fungal hyphae utilized nutrient then produced more biomass but after 6 days biomass production was slowly due to late growth phase. The above results for biomass production were supported by [38] that reported higher production of fungi with sucrose used as carbon source.

It is evident from Table 3 and Fig.- 5 that 4.575 mg/ml protein was produced by *Aspergillus* spp. after 2 days and also 1.11 mg/ml protein was recorded after 8 days incubation time. Further Table 3 indicates that fungi produced extracellular protein that used for their building blocks. Hence, the amount of protein was decreased 1.850, 1.0 and 0.905 mg/ml after 4, 6 and 10 days incubation periods respectively. Figure 5 shows that initially amount of protein was produced maximum then decreased because fungal mycelium production utilized these produced protein. Therefore, amount of protein was decreased after 4 to 6 days. After 8 days incubation time, protein amount was increased slightly then again decreased as same. The proteins are nitrogenous compounds and helps in synthesis of enzymes as well as energy source for living organisms. These observations are supported by Finkelstein and Ball [16], Moreira *et al.* [31], Banerjee *et al.* [5] and Karthiresan and Manivannan [24] which were reported that fungi have been used for their metabolite production.

Table- 4 and Fig.- 6 reveals that maximum enzyme activity was occurred after 2 days while shows minimum after 4, 6 and 10 days incubation periods. Further Table 4 indicates that *Aspergillus* spp. invertase production was obtained maximum on the basis of their assay after 48 hrs (21.83µg/ml) and also found minimum after 4 days incubation time. The amount of invertase in culture filtrate was slightly increased after 4 and 6 days incubation time. Initially sucrose was present in medium, function as inducer. Hence, amount of invertase was increase up to maximum but after growth of fungi on medium sucrose was metabolize then produced monosaccharide as glucose and fructose which play a role as inhibitor. Therefore concentration of invertase was decreased. These monosaccharides were utilized by fungi then again remaining sucrose was metabolizing so that invertase concentration was increased after 8 days incubation time.

These results are supported by Elorza *et al.* [15], Trimble and Maley [43], Olutiola and Cole [34], Vainstein and Peberdy [45] reported that fungi produced extracellular invertase by supplemented with sucrose. Rashad *et al.* [37] and Guimaraes *et al.* [19-20] also observed that invertase production was occurred in submerged condition at 30°C temperature and pH 5.0 after 4 days incubation time [44].

Table- 1 Composition of production medium.

S. No.	Ingredients	Quantity (gm/L)
1	Sucrose	20
2	Yeast extracts	10
3	Ammonium sulphate	1
4	Magnesium sulphate	0.75
5	Potassium dihydrogen phosphate	3.5
6	pH	5.0



Figure- 1 (A): *Aspergillus* spp. culture on agar slant



Figure- 1(B): *Aspergillus* spp. culture on agar plate



Figure- 2: Total dry biomass of *Aspergillus* spp. in varying incubation time (2 to 10 days).



Figure- 3 (A): Production media



Figure-3(B): Growth of *Aspergillus* spp. in production media

Table- 2 Estimation of total dry weight of biomass of *Aspergillus* spp.

S. No.	Incubation days	Dry biomass (gm)
1	2	0.45
2	4	0.80
3	6	1.00
4	8	1.02
5	10	1.10

Table- 3 Total protein estimation from culture filtrates of *Aspergillus* spp.

S. No.	Incubation days	Total protein(mg/ml)
1	2	4.575
2	4	1.850
3	6	1.000
4	8	1.110
5	10	0.905

Table- 4 Enzyme activity in culture filtrates of *Aspergillus* spp.

S. No.	Incubation days	Enzyme activity (µg/ml)
1	2	21.80
2	4	19.65
3	6	19.92
4	8	20.30
5	10	19.76

Biomass estimation




Figure- 4: Total dry biomass estimation of *Aspergillus* spp.

Protein estimation




Figure- 5: Total protein estimation from culture filtrates of *Aspergillus* spp.

Invertase estimation




Figure- 6: Invertase estimation from culture filtrates of *Aspergillus* spp.

CONCLUSION

By evaluating it was found that *Aspergillus* spp. metabolites were produced maximally at 30°C temperature and pH 5 after two day incubation times. These metabolites can be used in food and chemical industries for human diet and also preparation of glucose crystals and syrup as energy source.

REFERENCES

1. Alberto F, Bignon, C, Sulzenbacher, G, Henrissat, B and Czjzek, M; The three dimensional structure of invertase (a-

fructosidase) from *Thermotoga maritime* reveals a bimodular arrangement and an evolutionary relationship between retaining and inverting glycosidases. Journal of Biological Chemistry, 2004; 279: 18903-18910.

2. Aranda C, Robledo, A, Loera, O, Contreras-Esquivel, JC, Rodrigues, R and Aguilar, CN; Fungal invertase expression in solid-state fermentation. Food Technol. Biotechnol., 2006; 44(2): 229–233.

3. Arica MY, Senel, S, Alaeddinoglu, NG, Patir, S and Denizli, A; Invertase immobilized on spacer-arm attached poly (hydroxyethyl methacrylate) membrane: preparation and properties. J. Appl. Polymer Sci., 2000; 75: 1685–1692.
4. Ashok kumar B, Kayalvizhi, N and Gunasekaran, P; Optimization of media for a-fructofuranosidase production by *Aspergillus niger* in submerged and solid state fermentation. Process Biochemistry, 2001; 37: 331-338.
5. Banerjee AC, Kundu, A and Ghosh, SK; Genetic manipulation of filamentous fungi. In: Roussos S, editor. New horizons in biotechnology. Dordrecht (Neth) Kluwer Academic Publishers, 2003; 193- 198.
6. Bayramoglu G, Akgol, S, Bulut, A, Denizli, A and Arica, MY; Covalent immobilization of invertase onto a reactive film composed of 2-hydroxyethyl methacrylate and glycidylmethacrylate: properties and application in a continuous flow system. Biochem. Eng. J., 2003; 14: 117–126.
7. Belcarz A, Ginalska, G and LobarzewskiJ, PC; The novel non-glycosylated invertase from *Candida utilis* (the properties and the conditions of production and purification). Biochim. Biophys. Acta., 2002; 1594: 40–53.
8. Chaudhuri A and Maheswari, R; A novel invertase from a thermophilic fungus *Thermomyces lanuginosus*: it requirement of tito and protein for activation. Arch. Biochem. Biophys., 1996; 327: 98-106.
9. Chouhan PK and Das, P; Fungal protein production by small scale fermentation technology using *Aspergillus niger*. Glob. J. of Res. on Medicinal Plants & Indige. Medi., 2013; 2(1): 642- 647.
10. Chouhan PK, Verma, P, Sharma, M and Mali, K; Comparative study of amylase production by small scale fermentation technology under laboratory conditions using *Aspergillus niger*. J. of Bioprocess Techno. Photon, 2013; 98: 260-265.
11. Damle SP, Singh, K and Ghosh, D; Studies on soluble enzymes in penicillin fermentation broth. II. Invertase activity. Antibiotics Symposium, CSIR, New Delhi, 1958; 137-144.
12. Danisman T, Tan, S, Kacar, Y and Ergene, A; Covalent immobilization of invertase on microporous pHEMA–GMA membrane. Food Chem., 2004;85: 461–6.
13. Durowade KA, Kolawole, OM, UddinII, RO and Enonbun, KI; Isolation of ascomycetes fungi a tertiary institutions campus soil. Journal of Applied Science and Environment Management, 2009;12(4): 57-61.
14. Dworschack RG and Wickerham, LJ; Production of extracellular invertase by the yeast, *Saccharomyces uvarum* NRRLY Y-972. Arch. Biochem. Biophys., 1958; 76: 449-456.
15. Elorza MV, Villanueva, JR and Sentadreu, R; The mechanism of catabolite inhibition of invertase by glucose in *Saccharomyces cerevisiae*. Biochem. Biophys. Acta., 1977; 475 (1): 103-112.
16. Finkelstein DB and Ball, C; Biotechnology of filamentous fungi. Boston Butterworth-Heinemann, 1992; 221- 416.
17. Gill PK, Manhas, RK and Singh, P; Purification and properties of a heat-stable exoinulinase isoform from *Aspergillus fumigatus*. Bioresource Technology, 2006; 97: 894-902.
18. Gilman, JC; A manual of soil fungi, Revised 2nd edition, Oxford and IBH publishing Co., 1944.
19. Guimaraes LHS, Somera, AF, Terenzi, HF, Polizeli, ML and Jorge, JA; Production of fructofuranosidase by *Aspergillus niveus* using agroindustrial residues as carbon sources: Characterization of an intracellular enzyme accumulated in the presence of glucose. Process Biochem., 2009; 44: 237-241.
20. Guimaraes LHS, Terenzi, HF, Polizeli, ML and Jorge, JA; Production and characterization of a thermostable extracellular β -dfructofuranosidase produced by *Aspergillus ochraceus* with agroindustrial residues as carbon sources. Enzym. Microb. Technol., 2007; 42: 52–7.
21. Herwig C, Doerries, C, Marison, I and Von Stockar, U; Quantitative analysis of the regulation scheme of invertase expression in *Saccharomyces cerevisiae*. Biotechnol. Bioeng., 2001; 75: 247-58.
22. Ishaq F and Khan, A; Isolation, identification and comparative study of fungal and bacterial strains found in organic and inorganic soils of different agricultural fields. Recent Research in Science and Technology, 2011;3(11): 30-36.
23. Jon KM and Linda, L; Organic acid production by filamentous fungi, Advances in Fungal Biotechnology for Industry, Agriculture and Medicine. Jan and Lene Lange, Kluwer Academic/Plenum Publishers, 2007, 307-339.
24. Kathiresan K and Manivannan, S; α -Amylases production by *Penicillium fellutanum* isolated from mangrove rhizosphere soil. Afr. J. Biotechnol., 2006; 5 (10): 829- 832.
25. Keleke S, Agbor, E, Brauman, A and Griffon, D; Origine de l' activite amylasique dans la fermentation lactique des racines du manioc. In Agbor E *et al.* (eds) Transformation alimentaire du manioc. Orstom, Paris, 1998; 319- 28.

26. Kotwal SM and Shankar, V; Immobilized Invertase. *Biotechnol. Adv.*, 2009;27: 311-322.
27. Kotzelski J and Staude, E; Kinetics of bonded invertase: asymmetric polysulfone membranes. *J. Mem. Sci.*, 1996; 114: 201-14.
28. Kumar S, Chauhan, VS and Nahar, P; Invertase embedded-PVC tubing as a flow through reactor aimed at conversion of sucrose into inverted sugar. *Enzym. Microb. Technol.*, 2008; 43: 517-22.
29. Lowry OH, Rosebrough, NJ, Farr, AL and Randal, RJ; Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry*, 1951; 193: 265-275.
30. Miller, GL; Use of Dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*, 1959; 31: 426-429.
31. Moreira FG, Lenartovicz, V, De Souza, CGM, Ramos, EP and Peralta, RM; The use of α -methyl-D-glucoside, a synthetic analogue of maltose, as inducer of amylase by *Aspergillus* sp. In solid- state and submerged fermentation. *Braz. J. Microbiol.*, 2001; 32: 15- 19.
32. Nguyen QD, Rezessy-Szabo, JM, Bhat, MK and Hoschke, A; Purification and some properties of a fructofuranosidase from *Aspergillus niger* IMI303386. *Process Biochemistry*, 2005; 40: 2461-2466.
33. Nuero OM and Reyes, F; Enzymes for animal feeding from *Penicillium chrysogenum* mycelial wastes from penicillum manufacture. *Lett. Appl. Microbiol.*, 2002; 34: 413-6.
34. Olutiola PO and Cole, OO; *Plant Physiology*, 1980; 50: 26-31.
35. Onion, AHS, Allsop, D and Eggins, HOW; *Smiths Introduction to Industrial Mycology* (7th Ed.) Edward- Arnold Publishers Ltd. London, 1981.
36. Poonawalla FM, Patel, KL and Iyengar, MR; Invertase production by *Penicillium chrysogenum* and other fungi in submerged fermentation. *J. Appl. Environmental Microbiology*, 1965; 13: 749-754.
37. Rashad MM, Mahmoud, AEE, Desouky, MA and Nooman, MU; Purification and characterization of extra and intracellular β -fructofuranosidase from *Saccharomyces cerevisiae* growing on Eichhorniacrassipes leaf extract. *Deutsche Lebensmittel Rundschau*, 2006; 102: 157-166.
38. Robledo-Olivo A, Contreras-Esquivel Juan, C, Rodriguez, R and Cristobal, NA; Induction and repression kinetics of fungal beta-fructofuranosidase in submerged culture. *Int. J. Food Eng.*, 2009; 5(2): 1224.
39. Rubio MC and Maldonado, C; Purification and characterization of invertase from *Aspergillus niger*. *Current Microbiology*, 1995; 31: 80-83.
40. Rubio MC, Runco, R and Navarro, AR; *Phytochemistry*, 2002; 61: 605-609.
41. Smith JE, Berry, DR and Kristiansen, B; *Filamentous fungi Vol. IV, Fungal Technology*. Edward Arnold London, 1983.
42. Somiari RI and Bielecki, S; Effect of fructose and glucose supplementation on invertase-mediated synthesis of oligosaccharised from sucrose. *Bitechnol. Lett.*, 1995; 17(2): 519-521.
43. Trimble RB and Maley, F; *J. Biol. Chem.*, 1977; 252: 4409-4412.
44. Uma C, Gomathi, D, Muthulakhmi, C and Gopalakrishnan, VK; Production, purification and characterization of invertase by *Aspergillus flavus* using fruit peel waste as substrate. *Adv. Biol. Res.*, 2010; 4(1): 31-36.
45. Vainstein MH and Peberdy, JF; Regulation of invertase in *Aspergillus nidulans*: effect of different carbon sources. *J. Gen. Microbiol.*, 1991;137: 315- 321.
46. Vitolo M, Duranti, MA and Pellegrim, MB; Effect of pH, aeration and sucrose feeding on invertase activity of intact *Saccharomyces cerevisiae* cells grown in sugarcane black strap molasses. *J. Ind. Microbiol.*, 1995; 15:75-79.