

Research Article

Comparative Study of Lipid Profile among Stroke Patients with Gender Variations

Anbuselvan V^{1*}, Padmavathi R², Velmurugendran CU³, Shyamala Ramnath⁴, Jayachandran Paulswamy⁵.

¹Associate Professor, Department of Physiology, Sri Ramachandra University, Chennai, India

²Professor and Head, Department of Physiology, Sri Ramachandra University, Chennai, India

³Professor and Head, Department of Neurology, Sri Ramachandra University, Chennai, India

⁴Professor and Head, Department of Physiology Chettinad Medical College and RI, Chennai, India

⁵Occupational and General Physician, Madhavaram, Chennai, India

***Corresponding author**

Dr. Anbuselvan Veeraiah

Email: Drvas2003@yahoo.co.in

Abstract: Stroke is defined according to WHO as rapidly developing clinical signs of focal disturbance of cerebral functions with symptoms lasting for 24hrs or longer or leading to death, with no apparent cause other than vascular origin. This study was aimed to find out the association of one common risk factor of cardiovascular complication risk factor namely lipids level in stroke patient. And the main course of the study is to find the lipid level variations among both genders of stroke population. The stroke patients who were admitted in the Stroke ward of Department of Neurology, SRMC&RI were recruited in this study Patients with diagnosis of stroke comprising 50 consecutive patients of both the genders of age group around 21-60 yrs were included in the study while patients on lipid lowering therapy were excluded from study. A serum sample after 12 hours of overnight fasting was taken on the both groups of patients. Total serum cholesterol, triglycerides, LDL cholesterol, VLDL-cholesterol and HDL-cholesterol was determined, using enzymatic colorimetric method. Statistical analysis - Done by comparison of lipid profile in two subgroups, using proportion test for any significant differences. From our study we concluded that females are in high risk for prevalence of cardiovascular disease. Total serum cholesterol was high in female stroke patient's 210 ± 32.92 mg/dl when compared to male stroke patient's 190 ± 25.82 mg/dl which is statistically significant. Total triglycerides were increased in female stroke patients because of high body mass index when compared to male stroke patients was around 213 ± 23.92 , 197 ± 58.43 mg/dl which is statistically significant. HDL-cholesterol level were very low in female stroke patients when compared to male stroke patients this is statistically more significant. The values were around 23.72 ± 12.06 , 37.43 ± 7.18 mg/dl. LDL cholesterol is the direct marker of large vessel obstruction. LDL cholesterol level was found to be more in female stroke patients when compared to male stroke patients. The values are 141.81 ± 31.89 , 123.92 ± 33.73 mg/dl. VLDL-cholesterol was high in female stroke patients when compared to male stroke patients which are statistically significant the values are 44.93 ± 8.75 , 37.83 ± 9.33 mg/dl.

Keywords: Stroke, Cardiovascular diseases, Total serum cholesterol, Triglycerides, LDL -cholesterol, VLDL-cholesterol and HDL-cholesterol.

INTRODUCTION

More than 2400 years ago the father of modern medicine Hippocrates recognized and described stroke as sudden onset of paralysis. Until recently modern medicine has had very little power over this disease but the world of stroke medicine is changing and new better therapies are being developed every day [1]. Dr. Johann Jacob Wafer (1820) was the first person to demonstrate the stroke by doing post-mortem and bleeding in the brain of patients who died of apoplexy and he also described apart from bleeding, the blockage of the main arteries supplying the brain. Thus stroke became known as "Cerebral vascular Accident"(CVA). After several years of research the medical science has come up with

an array of risk factors and various Path physiological mechanism by which the stroke is caused. The nontraditional factors are like increased Plasma fibrinogen, Hyper Homocysteinemia, High level of Lipoprotein. This has helped to identify the potential candidate for stroke among the patients just by looking at the presence of the risk factors in the blood and to employ the methods of modifying or eliminating these risk factors and thereby to prevent the incidence of stroke in our patients [2]. Serum lipid levels have an established effect on short term mortality due to strokes [3]. It is important to evaluate the difference in serum lipid levels in both the genders of strokes to guide lipid-lowering therapy which can reduce incidence of stroke

and related mortality by adapting primary and secondary preventive measures among the stroke patients [4, 5].

Recent community surveys from many regions of India show a crude prevalence rate for stroke presumed to be of vascular origin in the range of 200/100,000 persons (0.2%) [6].

In approximately 1% of all patients with ischemic stroke and in up to 4% of young adults with stroke, the major precipitant of brain ischemia is mainly by blockage of artery with free radicals and lipid molecules [7-9]. The contribution of dyslipidemia in children with ischemic stroke has been reported to be 20 to 50% in most studies. Since dyslipidemia is also one of the important factor which contribute to the majority of cases of stroke, this study was undertaken to find out the association of lipid profile level in stroke patients.

Serum HDL-cholesterol has anti-atherogenic properties with ability to trigger the flux of cholesterol from peripheral cells to the liver and thus having a protective effect [10].

MATERIALS AND METHODS

The study was conducted in the Department of Physiology in Sri Ramachandra Medical College & Research Institute, Chennai. The stroke patients admitted in the Stroke ward of Department of Neurology, Sri Ramachandra Medical College & Research Institute, Chennai included in this study. The sample size was around 50 and the age groups of the patients were around 21 to 85 yrs. Among stroke patients were 25 males and 25 female's patients. A questionnaire was circulated to the patient attenders

asking about the personal history like smoking, alcoholism, physical exercise and history of diabetes and hypertension. The lipid profile investigations were done fasting state. Venous blood samples were collected from anti cubital vein which is transferred into plain tubes. Samples were centrifuged at 4°C for 20 minutes after incubation of 30 minutes serum was extracted. The sera were analyzed for serum lipid profile including total cholesterol, triglyceride, LDL-cholesterol, VLDL-cholesterol and HDL-cholesterol by standard enzymatic colorimetric method using chemistry auto-analyzer. Mean values of the cholesterol, triglyceride, LDL-cholesterol, HDL-cholesterol and VLDL-cholesterol was determined.

RESULTS

The mean age of presentation of patients with stroke was 54.2±32 years of both the genders. From our study we concluded that females are in high risk for prevalence of cardiovascular disease. Total serum cholesterol was high in females' 210±32.92 mg/dl when compared to male's 190±25.82 mg/dl which is statistically significant. Total triglycerides were increased in females because of high body mass index when compared to males was around 213±23.92, 197±58.43 mg/dl which is statistically significant. HDL-cholesterol level were very low in females when compared to males which is statistically more significant. The values were around 23.72±12.06, 37.43±7.18 mg/dl. LDL cholesterol is the direct marker of large vessel obstruction. LDL cholesterol, level was found to be more in females when compared to males. The values are 141.81±31.89, 123.92±33.73 mg/dl. VLDL-cholesterol was high in females when compared to a male which is statistically significant the values are 44.93±8.75, 37.83±9.33 mg/dl.

Table 1: Shows the Variations among the Lipid Profile of Both the Genders

Variables	TC mg/dl	TGL mg/dl	HDL mg/dl	VLDL mg/dl	LDL mg/dl
Males	190±25.82	197±58.43	37.43±7.18	37.83±9.33	123.92±33.73
Females	210±32.92	213±23.92	23.72±12.06	44.93±8.75	141.81±31.89
P value	<0.005*	<0.001**	<0.005*	<0.01***	<0.005*

There was an increase in Serum TC, TGL, VLDL, and LDL in female patients when compared with males. HDL cholesterol was found to be very less in females when compared with males. This is statistically significant. Significant* highly significant **

DISCUSSION

Stroke is a clinical syndrome characterized by loss of cerebral functions, with symptoms lasting more than 24 hours or leading to death with no apparent cause other than that of vascular origin [11]. Cerebral atherosclerosis with atheroma formation is the basic underlying patho-physiologic mechanism in ischemic stroke [12]. Conflicting results exist in the literature about the correlation between the total plasma

cholesterol of patients and the risk of stroke [13]. Different environmental factors, genetic influences, and aging are playing a wide role in the prevalence of stroke among the old age group.

Hypertension, DM, AF, and high cholesterol are the highest ranking controllable medical risk factors for stroke, whereas smoking, alcohol consumption, and obesity rank as the most important lifestyle stroke risk factors [14].

Dyslipidemia is a major risk factor for coronary heart disease [15], and atherosclerosis not only in stroke patients, but also in old age group who are nearing the age around 60yrs of both the age groups. Qizilbash *et*

al. from his study reviewed the the relationship between serum total cholesterol and subsequent stroke. They concluded that there was a significant association between serum lipid profile and prevalence of stroke [16]. Khan *et al.* and Tanveer *et al.*, from there study they proved that Hyperlipidemia was present in 16% patients of stroke. They analyzed that the serum lipid profile was 3rd most common risk factor for stroke [7, 17].

Sacco RL investigated cross-sectional differences in sociodemographic and stroke risk factors and suggested that the burden of stroke risk factors is increased in both blacks and Hispanics with stroke [18]. Andersen KK *et al.* from their study they showed Cardiovascular risk factors were generally more prevalent in men. Lifestyle cardiovascular risk factors were more common in the young. Prevalence of hypertension, diabetes mellitus, coronary heart disease, and, in men, also atrial fibrillation go down after the age of 70 to 80 years [19]. The role of lipids in stroke patients in our environments needs further work in carefully selected stroke subjects. This finding may be important in this respect in our environment, as our populace now imbibe the western habit of eating high fatty foods and doing away with the indigenous high fiber diet.

CONCLUSION

Different environmental factors, genetic influences, and aging are playing a wide role in the prevalence of stroke among the old age group. Dyslipidemia is a major risk factor for coronary heart disease ,and arteriosclerosis .Cerebral arteriosclerosis with atheroma formation is the basic underlying patho-physiologic which increases the risk of cardiovascular, and vessel wall disease among stroke patients . In our study female patients had an increase in lipid level when compared to males. This variations is mainly due to increase in body mass index, and imbalance in female sexual hormones after menopause may increase the adipose tissue fat content ,which favours the blockage of vessel walls and causes the ischemic responses. Regular monitor of lipid profile among stroke patients may decrease the risk of arteriosclerosis and cardiovascular disease among the stroke patients.

ACKNOWLEDGEMENT

The author would like to thank theHead of the Department, Professor, Associate professor, Assistant professors and non teaching staff of Physiology and Neurology department of Sri Ramachandra University Pour, Chennai for guiding me for each & every step of this research work by giving useful suggestions and made me complete this work successfully.

REFERENCES:

1. Smolders, Robin Lemmens, Thijs V; Stroke: Hope through Research," NINDS. Publication. July 2004; No. 99-2222. NIH Publication.

2. Burkman K; Stroke: History. In The stroke Recovery Book: A Guide for Patients and Families. 2nd edition, 1992.
3. Zuliani G, Cherubini A, Atti AR, Ble A, Vavalle C, Ditodoro F et al.; Low cholesterol levels are associated with short term mortality in older patients with ischaemic strokes. J Gerontol 2004; 59: 293-297.
4. Kirshner HS; Medical prevention of stroke. South Med J., 2003; 96: 354-358.
5. Laloux P, Galanti L, Jamart J; Lipids in ischaemic stroke subtypes. Acta Neurol Belg., 2004; 104:13-19.
6. Bhalla A, Gupta OP, Gupta SB; Predicting Mortality in stroke. Neurology India, 2002; 50(3): 279-281.
7. Hart RG, Sherman DG, Miller VT, Easton JD; Diagnosis and management of ischemic stroke: Selected controversies. Curr Probl Cardiol.,1983; 8: 43-53.
8. Adams HP, Butler MJ, Biller J, Toffal GJ; Nonhemorrhagic cerebral infarction in young adults. Arch Neurol., 1986; 43:793-796.
9. Klein GM, Seland TP; Occlusive cerebrovascular disease in young adults. Can J Neurol Sci 1984; 11: 302-304.
10. Jia L, Bai H, Fu M, Xu Y, Yang Y, Long S; Relationship between plasma HDL subclasses distribution and apoA-I gene polymorphisms. Clinica Chimica Acta 2005; 360: 37-45.
11. Stroke: diagnosis and initial management of acute stroke and transient ischaemic attack (TIA). NICE guideline Draft for consultation, January 2008. Available from <http://www.nice.org.uk/nicemedia/live/11646/38877/38877.pdf>
12. Khan J, Attique-ur-Rehman, Ali SA, Jielani A; Frequency of hypertension in stroke patients presenting at Ayub Teaching Hospital. J Ayub Med Coll Abbottabad., 2006; 18(1): 59-61.
13. Mahmood A, Sharif MA, Khan MN, Ali UZ; Comparison of Serum Lipid Profile in Ischaemic and Haemorrhagic Stroke. Journal of the College of Physicians and Surgeons Pakistan, 2010, 20 (5): 317-320.
14. Goldstein LB, Adams R, Albert MJ, Appel LJ, Brass LM, Bushnell CD et al.; Primary prevention of ischemic stroke. A guideline from the American Heart Association/American Stroke Association Stroke Council. Stroke, 2006; 37:1583–1633.
15. Goff DC Jr, Bertoni AG, Kramer H, Bonds D, Blumenthal RS, Tsai MY et al.; Dyslipidemia Prevalence, Treatment, and Control in the Multi-Ethnic Study of Atherosclerosis (MESA). Circulation, 2006; 113: 647-656.
16. Qizilbash N, Duffy SW, War low C, Mann J; Lipids are risk factors for ischemic stroke: overview and review. Cerebrovascular Disease, 1992; 3(2): 127-136.

17. Tanveer A; Localization and management in Cerebrovascular accident: a comparison of clinical assessment versus C.T scan [Dissertation]. College of Physicians & Surgeons Pakistan; Karachi: 1996.
18. Sacco RL, Kargmann DE, Zamanillo MC; Race-ethnic differences in stroke risk factors among hospitalized patients with cerebral infarction: the Northern Manhattan Study. *Neurology*, 1995; 45(4): 659–663.
19. Andersen KK, Andersen ZJ, Olsen TS; Age- and Gender-Specific Prevalence of Cardiovascular Risk Factors in 40 102 Patients with First-Ever Ischemic Stroke. A Nationwide Danish Study. *Stroke*, 2010; 41(12): 2768-2774.