

Research Article

Post harvest quality of tomato (*Lycopersicon esculantum* mill.) as influenced by drip irrigation scheduling in changing climatic scenario

Sunil Kumar¹ A. Abdul Haris² and V.Chhabra³

¹Associate professor, Department of Horticulture, North Eastern Hill University, Tura Campus, Chandmari-794 0002, West Garo Hills District, Meghalaya

²Principal scientist (Agronomy), ICAR Research Complex for Eastern Region, Patna

³Research Associate, ICAR, Research Complex for Eastern Region, Patna

***Corresponding author**

Dr. Sunil Kumar

Email: sunu159@yahoo.co.in

Abstract: A field experiment was conducted at Sabajpura farm, ICAR research complex for eastern region, Patna on post harvest quality of tomato as influenced by drip irrigation scheduling. The experiment was conducted in split plot design with irrigation water equivalent to 100, 80 and 60% ET (Evapo transpiration) in main plots and water as per daily, alternate days and once in three days schedule in sub-plots. Observations regarding TSS (0 Brix), acidity and fruit firmness (Kg/m²) was undertaken. These parameters showed better response at 80% ET when plant received irrigation once in two days which was at par with once in three days irrigation. Highest TSS in tomato fruit was recorded under irrigation scheduling once in two days (3.430 Brix). However, Lowest acidity was associated with 100% irrigation (4.00) followed by 60% irrigation (5.21). Irrigation scheduling with once in two days proved maximum firmness of tomato fruit (3.53) which was at par with once in three days irrigation scheduling (3.39 Kg/cm²). Drip irrigation is an effective adaptation technology to maintain quality of vegetables like tomato in warmer climate in future time periods.

Keywords: drip irrigation, evapo-transpiration, post harvest quality, tomato.

INTRODUCTION

Tomato is a major solanaceous crop next to potato considered as remunerative vegetable crop in India. It is a rich source of minerals, vitamins and organic acids, used as raw in salad, fetch greater demand in processing industry for making ketchup, sauce and puree. The post harvest losses in tomato are high and can go up to 60 per cent [1].

Agriculture sector depends on climate and climatic resources leading to the special stress upon the need to study impacts of climate change on agricultural production at local, regional, national as well as on global scales. Elevated CO₂ and changes in temperature, besides affecting the crop affects the environment which in turn may have either beneficial or damaging effect on agricultural production. Due to increase in temperature even by 1°C, fruit set and quality of vegetables like tomato may be affected [2]. Loss of quality includes the physiological and chemical changes associated with ripening resulting in changes in appearance, taste, texture and development of off flavours. Development of colouring pigment lycopene is hampered above 27 °C.

Food and energy security of India are crucially dependent on favourable weather and the timely availability of adequate amount of water. Air temperature and precipitation are principle elements of weather systems, so for the prediction of future climate conditions, level of variability of these two weather elements must be examined. According to the estimates by IPCC (2007) earth's linearly averaged surface temperature has increased by 0.74°C during the period 1901-2005 [3]. Increasing trends of rainfall and minimum temperature in Gangetic plains of Bihar was observed [4; 2010a]. Weather generators can be used to generate long term weather data wherever data is not available for impact studies. One such generator is LARSWG which was used and produced similar observations of rainfall and temperature as actual weather data for Bihar in eastern India [5; 2010b]. Although increase in CO₂ is likely to be beneficial for crops, yet associated temperature and rainfall variability may change the food production situation. Simulation studies are useful in predicting future climate scenario based on changing trends in temperature, CO₂ and rainfall etc [6 (2010c); 7 (2012a)]. Climate change impacts on cereal crops are studied in detail, however scarce information is available about impacts on vegetables and fruits. Cereal crop like wheat in Patna

and Ranchi showed a declining yield trend due to increased temperature (thermal stress) coinciding with grain filling stage of crop [8]. Simulation studies performed for different time periods using HadCM3 factors at four centres located in three different agroecological zones showed that simulated yield of wheat may decline in future time periods [9(2013a),10]. Simulation studies showed that long duration rice varieties are more prone to yield decline under future climate scenarios [11 (2010d); 12 (2012); 13(2012b)]. Climate change will affect future water supplies. Development of production systems geared towards improved water-use efficiency and to mitigate the effects of hot and dry conditions in vegetable production systems are top research and development priorities in changing climatic scenarios. There is wide scope for enhancing irrigation efficiency through adoption of drip irrigation systems and other water-conserving technologies [14, 15]. Drip irrigation system has proved its superiority over other conventional methods of irrigation, especially in fruit and vegetable crops owing to precise and direct application of water in the root zone. 20 litre water per day was ideal for cocoa in mixed cropping system with arecanut [16]. Sujatha and Haris, 2000 [17] reported that ferti-drip and drip irrigation in arecanut increased number of feeder roots four fold and two fold respectively compared to basin irrigation indicating increased root activity. It also influences post harvest quality of fruits and vegetables manifold. Therefore, it is necessary for the vegetable growers to know how effectively drip irrigation system enhances productivity and post harvest quality of tomato. Higher yield under drip irrigation at 80% ET once in 3 days in tomato and once in 2 days in cabbage and daily in okra, is reported [18; 19(2014a), 20(2014b)]. Since, drip irrigation saves 30-40% fertilizer/chemical with enhanced quality of produce as compared to surface irrigation and also helpful in reducing labour cost, salt concentration in the root zone and disease incidence. Therefore, an effort has been laid out keeping all these in view to study the post harvest quality of tomato as influenced by drip irrigation scheduling.

MATERIALS AND METHODS

A field experiment was conducted at ICAR Research Complex for Eastern Region, Patna on tomato. The tomato seed S-41 (Gotya, a Mahyco hybrid) was sown in nursery during first week of September and transplanted at the spacing of 60x60cm in the first week of October. The experiment was laid out in split plot design with three replications. The treatments as the main plot factors were irrigation level at 100% (I_1), 80% (I_2) and 60% (I_3) ET (Evapotranspiration) while as the sub-plot factors were scheduling at daily (S_1), once in two days (S_2) and once in three days (S_3). The drip irrigation systems were consisted of 2lph capacity with online drippers. One lateral for one row of crop with one dripper per plant was adopted in each plot. Each sub-plot consisted of ten

rows and ninety plants. The daily irrigation was applied in drip treatments at 1kg/cm² pressure. The daily pan evaporation was used to compute the daily water requirement of the crop through drip systems. The effective rainfall was computed by considering the pan evaporation and precipitation received during irrigation intervals. Tomatoes were harvested several times and data were added to estimate the TSS, acidity and fruit firmness during experimentation and analyzed using statistical methods [21].

RESULTS AND DISCUSSION

Effect of irrigation scheduling on TSS

The data presented in table-1 indicated that there was no significant difference among the irrigation at 100, 80 and 60% on TSS. Non-significant differences of irrigation treatments on TSS were observed in musk melon [22]. However, significant differences was observed under irrigation scheduling once in two days which recorded highest TSS (3.43⁰ Brix) followed by once in three days irrigation (3.31⁰ Brix), while, least TSS was observed with daily irrigation scheduling (3.09⁰ Brix). However, significant differences was noticed during second year and 60% irrigation recorded highest TSS (4.88⁰ Brix) which was on par with 100% (4.80⁰ Brix) irrigation. These results are in close conformity with the findings of Joshi *et al.* 2012 [23] in litchi. But no significant differences were observed under irrigation scheduling and interactions. Drip irrigation system improved TSS in banana [24].

Effect of irrigation scheduling on acidity

Irrigation at 100, 80 and 60% with daily, once in two days and once in three days irrigation scheduling showed significant difference in acidity of tomato fruit (Table-1). Lowest acidity was associated with 100% irrigation (4.00) followed by 60% irrigation (5.21) during first year. In the irrigation scheduling daily scheduling noted lower acidity (4.28) than once in two days and once in three days irrigation. However, irrigation at 100, 80 and 60% with daily, once in two days and once in three days irrigation scheduling did not showed significant difference in acidity of tomato fruit during second year. Whereas, interaction was found to be significant and 60% irrigation with once in three days irrigation scheduling noted lower acidity (3.05) followed by 80% irrigation with once in two days irrigation scheduling (3.55). Drip irrigation system lowers acidity in banana was observed [24].

Effect of irrigation scheduling on fruit firmness

Significant difference was noted among the irrigation scheduling but there was no significant difference among irrigation at 100, 80 and 60% ET (Table-1). Irrigation scheduling with once in two days proved maximum firmness of tomato fruit (3.53) followed by irrigation scheduling once in three days irrigation (3.39 Kg/cm²) during first year of investigation. Whereas, no Significant difference was noted among the 100, 80 and 60% irrigation with daily, once in two days and once in

three days irrigation scheduling during second year. But irrigation scheduling with once in two days proved maximum firmness of tomato fruit (6.01 Kg/cm²) than

daily and once in three days irrigation. However, irrigation at two days and three days were at par.

Table-1: Effect of drip irrigation scheduling on TSS, acidity and firmness in tomato

Treatments	TSS (^o Brix)		Acidity		Firmness (Kg/cm ²)	
	1 st year	2 nd year	1 st year	2 nd year	1 st year	2 nd year
Main plots						
I ₁ (100%ET)	3.23	4.80	4.00	3.19	3.36	5.81
I ₂ (80%ET)	3.31	4.51	5.44	3.59	3.49	6.05
I ₃ (60%ET)	3.29	4.88	5.21	3.39	3.22	5.97
C.D	N.S	0.24	0.17	N.S	N.S	N.S
Sub-Plots						
S ₁ (Daily)	3.09	4.74	4.28	3.26	3.15	5.84
S ₂ (2 days)	3.43	4.75	5.06	3.54	3.53	6.01
S ₃ (3 days)	3.31	4.69	5.30	3.37	3.39	5.98
C.D	0.19	N.S	0.52	N.S	0.29	N.S
I ₁ S ₁	2.97	4.97	4.73	3.07	3.21	5.95
I ₁ S ₂	3.43	4.97	6.18	3.86	3.52	5.87
I ₁ S ₃	3.30	4.47	5.41	3.23	3.37	5.60
I ₂ S ₁	3.20	4.27	3.36	3.40	3.10	5.58
I ₂ S ₂	3.40	4.52	3.78	3.55	3.79	6.27
I ₂ S ₃	3.33	4.73	4.85	3.82	3.58	6.30
I ₃ S ₁	3.10	5.00	4.76	3.32	3.15	5.58
I ₃ S ₂	3.47	4.77	5.23	3.20	3.29	5.90
I ₃ S ₃	3.30	4.87	5.65	3.05	3.21	6.03
C.D	N.S	N.S	N.S	0.45	N.S	N.S

SUMMARY

Under changing climate scenario yield and quality of most crops are affected due to increasing temperature and decreased water availability. Improved irrigation technologies like drip irrigation are the need of the time to maintain yield and quality of produce especially in vegetables. The result obtained from the experimentation concluded that drip system is very effective and efficient method of irrigation for enhancing post harvest quality of tomato fruit and the scheduling of irrigation once in three days at 80% ET through drip systems was suitable for improving TSS, acidity and firmness of tomato fruits.

REFERENCES

1. Shanmugasundaram, S; Surmountable challenges. The Hindu Survey of Indian Agriculture 2004. M/s Kasturi & Sons Ltd., Kasturi Building, Chennai-600002, pp 126-127.
2. Sato, S; The effects of moderately elevated temperature stress due to global warming on the yield and the male reproductive development of tomato (*Lycopersicon esculentum*. Mill). Hort. Research, 2006; (60): 85-89.
3. IPCC, Summary for policymakers. In Climate Change 2007: The Physical Science Basis (eds Solomon, S. D. et al.), Cambridge University Press, Cambridge, UK [4] Pant, G. B. and Kumar, K. R. (1997). Climates of South Asia, John Wiley, Chichester, UK.
4. Haris AA, Chhabra V, Biswas S; Rainfall and temperature trends at three representative agroecological zones of Bihar. Journal of Agrometeorology, (2010a); 12(1) 37-39.
5. Haris AA, MA Khan, V Chhabra, S Biswas, A Pratap; Evaluation of LARSWG for generating long term data assessment of climate change impact in Bihar. Journal of Agrometeorology, 2010; 12:198-201.
6. A.A Haris, Vandna Chhabra, Sandeep Biswas; Impact of climate change in rice and wheat crops, adaptation strategies and resource conservation measures. In Resource conservation technologies for food security and rural livelihood. Published by Agrotech Publishing Academy, Udaipur. 2010; 102-110.
7. Haris AA, Bhatt BP, Chhabra V; Climate Change and Food Security. In: Status of Agricultural Development in Eastern India. Eds.: B.P. Bhatt, A.K. Sikka, Joydeep Mukherjee, Adlul Islam, 2012. pp. 49-60.
8. Haris AA, Kumari P, Chhabra V, Biswas S; Modeling the impact of anticipated climate change on wheat yields in two different agro-climatic zones of eastern India. Journal of Agrometeorology, 2011; 13(2):116-118.

9. Haris AA, Biswas S, Chhabra V, Elanchezhian R , Bhatt BP; Impact of climate change on wheat and winter maize over sub-humid climatic environment. *Curr. Sci.*, 2013; 104(2): 206-214.
10. Haris AA, Chhabra V; Climate change impact on chickpea yield over a sub humid climatic environment. *International Journal of Research in Agricultural Sciences*, 2014; 1(4): 258-261.
11. Haris AA, Biswas S, Chhabra V; Impact of anticipated climate change on the yield of rice variety MTU 7029 in Patna, Bihar. *Indian Journal of Agronomy*, 2010; 55 (4):42-45.
12. Elanchezhian R, AA Haris, S Biswas, V Chhabra; Simulation of yield and its component traits of rice (*Oryza Sativa L.*) varieties grown in Indo-Gangetic plains of Bihar under projected climate change. *Indian J. Plant Physiol.*, 2012; 17 (3 &4):195-202.
13. Haris AA, Bhatt BP, Chhabra V, Biswas S, Elanchezhian R; Climate change impacts on yields of phenologically different rice varieties over a sub humid climatic environment. *Agri. Res.*, 2012; 2(4):319-329.
14. FAO; Guidelines for designing and evaluating surface irrigation systems. *Irrigation and Drainage Paper 45*. FAO, Rome. 1989.
15. FAO (1961-1990). *Agrostat 1961-1990*. FAO, Rome.
16. Abdul Haris, D. Balasimha, S. Sujatha, Ravi Bhat and K. B. Abdul Khader .The influence of drip irrigation and fertilizer levels on yield and photosynthetic characteristics of cocoa in mixed cropping system with arecanut. *Journal of Plantation Crops*, 1999; 27 (2):131-135
17. Sujatha S and Haris AA; Root distribution as influenced by different methods of irrigation in young areca nut (*Areca catechu L.*) palms. *J. Plantation Crops*. 2000; 28(2):117-122.
18. Haris AA and Kumar S; Drip irrigation scheduling in tomato (*Lycopersicon esculentum Mill.*). *HortFlora Res. Spectrum*, 2014; 3(2): 184-186.
19. Haris AA, Kumar S and Singh AK; Water requirement and irrigation scheduling through drip systems in cabbage (*Brassicaoleracea var. capitata*). *HortFlora Res. Spectrum*, 2014; 3(2): 166-168.
20. Haris AA, Kumar S, Singh AK and Rajan K; Drip irrigation scheduling in Okra (*Abelmoschus esculentus (L) Moench*). *Hort Flora Res. Spectrum*, 2014; 3(3): 274-277.
21. Panse VG and Sukhatme PV; *Statistical methods for agricultural workers*. Indian Council of Agricultural Research, New Delhi. 1985.
22. Hartz TK; *Effects of Drip Irrigation Scheduling on Muskmelon Yield and Quality*. *Scientia Horticulturae*, 1997; 69: 117-122.
23. Joshi G, Singh PK, Srivastava PC and Singh SK; *Effect of Mulching, Drip Irrigation Scheduling and Fertilizer Levels on Plant Growth, Fruit Yield and Quality of Litchi (Litchi chinensis Sonn.)*. *Indian Journal of Soil Conservation*, 2012; 40 (1):46-51.
24. Pawar DD and Dingre SK; *Influence of fertigation scheduling through drip on growthand yield of banana in western Maharashtra*. *Indian Journal of Horticulture*, 2013; 70 (2): 200-205.