

Research Article

Studies on the Effects of Treatments of Methyl Methanesulphonate and Sodium Azide on Induction of Variability in *Hordeum vulgare* L.

Shahnawaz Khursheed, Zarrin Akhtar, Samiullah Khan

Mutation Breeding Lab, Department of Botany, A.M.U., Aligarh-202 002, India

***Corresponding author**

Shahnawaz Khursheed

Email: shahnawazkhursheed95@gmail.com

Abstract: Cereals constitute an important source of food in India. India depends mostly on cereals and pulses for food. Cereals contain high amounts of starch along with considerable amounts of proteins, lipids and minerals. The present experiment was conducted to induce the variability in *Hordeum vulgare* L. by methyl methanesulphonate and sodium azide. The seeds were sown to raise the M₁ generation. Seed germination of mutagen treated seeds in M₁ and cytology in PMC's of M₁ generation plants were observed. Seed germination decreased with increasing concentrations of MMS but it showed irregular trend in case of SA treated seeds. Seeds obtained from M₁ generation plants were sown to raise the M₂ generation. Lower concentrations of both mutagens produced less frequency of meiotic abnormalities than higher concentrations but MMS produced more frequency of meiotic abnormalities than SA. Different quantitative characters were studied in M₂ generation. Quantitative characters of a plant are the most important characters associated with yield. Data on quantitative characters did not show regular trend but followed an irregular trend. But lower concentrations of MMS treated plants showed more positive response than SA treated plants.

Keywords: *Hordeum vulgare* L., methyl methanesulphonate, sodium azide.

INTRODUCTION

Food insecurity and malnutrition are threatening the population worldwide. India ranks 63rd according to Global Hunger Index [1]. More than 40% of the population in India has the highest prevalence to be underweight specially in children under five years old. The current increase in population is also posing a great threat to the nation. Due to more urbanization in India, less land is available for producing sufficient food to cope these problems. Scientists throughout the world are busy in ameliorating to solve these problems. Different methodologies are applied worldwide, of which mutation breeding forms an important part. Induced mutation is a fast method of increasing the variability and so the yield of plants and hence reducing the food insecurity and malnutrition of the fast growing population.

Barley, an important nutritious plant is used mostly in Asia and North Africa. Barley contains an abundant amount of starch (65-68%) along with considerable amount of proteins (10-17%) and minerals (1.5-2.5%) [2]. It belongs to family Poaceae and is an annual herb. The diploid chromosome number of barley is 14 (2n=14). Thus it can be said that it is an important nutritional plant.

Therefore, the present experiment was conducted to induce the variability in the plant and increasing the yield of plant. The mutagens used were methyl methanesulphonate and sodium azide.

MATERIALS AND METHODS

Fresh and healthy seeds of *Hordeum vulgare* L. were used for the experiment. 1% Stock solutions of both the mutagens were first prepared. For methyl methanesulphonate the stock solution was prepared in phosphate buffer of pH 7 while that of sodium azide was prepared in phosphate buffer of pH 3. From this stock solution, different concentrations (0.01%, 0.02%, 0.03% and 0.04%) of both mutagens were prepared. Nine sets of seeds each containing about 30 seeds (3 replications of each treatment) were used for this experiment. Those seeds were pre-soaked in double distilled water for 9 hours. Then the seeds were put in different concentrations of both mutagens for 6 hours while one set was kept in double distilled water to act as control. After the mutagenic treatment, the seeds were washed in running tap water to remove any residual effects of mutagens. The seeds were sown in to raise the M₁ generation. For cytology, flower buds of mutants and control were fixed in Carnoy's fluid (1 part Glacial acetic acid : 3 parts Chloroform : 6 parts Alcohol) for 24 hours. Traces of ferric chloride were added to improve the stainability of chromosomes. The flower

buds were washed with and preserved in 70% alcohol. Anthers were smeared in 1% acetocarmine solution. Meiosis of PMC's was observed in M₁ generation plants. The M₁ seeds were sown to raise the M₂ generation. Different quantitative characters were studied in M₂ generation plants.

Studies in M₁ generation

Seed germination

Germination (%) =

$$\frac{\text{No. of seeds germinated}}{\text{No. of seeds sown}} \times 100$$

Inhibition

Inhibition (%)

$$= \frac{\text{Control} - \text{Treated}}{\text{Control}} \times 100$$

Frequency of meiotic abnormality (%) =

$$\frac{\text{Total no. of abnormal PMC's of particular treatment}}{\text{Total no. of PMC's observed in that particular treatment}} \times 100$$

RESULTS AND DISCUSSION

Sodium azide is one of the most powerful mutagens and its mutagenicity is mediated through the production of azide compound. On the other hand, methyl methanesulphonate is an alkylating agent and alkylates DNA on N⁷-deoxyguanosine and N³-deoxyadenosine predominantly. The methyl methanesulphonate was found to induce more variability than sodium azide in both generations.

M₁ GENERATION STUDY

Seed Germination and Inhibition

Data recorded on seed germination and inhibition is shown in Table-1. As is clear from the Table, the seed germination decreased with increasing concentrations of MMS. Similar results were obtained by Ambreen *et al* [3], but followed an irregular trend in case of SA.

Cytology was performed in buds of M₁ generation plants. The control plants showed normal meiosis while the mutagen treated plants showed abnormal meiosis. The MMS treated plants showed higher frequency of meiotic abnormalities than SA treated plants (Table-2). Gulfishan *et al* [4] observed similar results for MMS but in comparison with diethyl sulphate. The different meiotic abnormalities observed were disturbed metaphase, dipolarity at telophase-I, multivalent, laggard formation at anaphase-I. (Plate-1). Dose dependent abnormalities of both mutagens were observed.

The laggards and fragments which fail to reach to the poles results ultimately into the formation of micronucleus[5-6]. Ring formation of chromosomes has also been observed. It may be due to the breaking of chromosome ends[7].

Disturbed metaphase was also observed in treated plants. Disturbed metaphase has also been already reported in *Allium cepa* L. by Maslam [8] after treating it with some medicinal extracts. Disturbed metaphase has been observed due to the effect on proteins that constitute the spindle. Laggard formation occurs due to the improper organization of spindle[9]. Multivalents as observed may occur due to the translocation and inversion[10].

M₂ GENERATION STUDY

The M₂ generation plants were screened for different quantitative traits like plant height, number of fertile branches per plant and plant yield. (Table-3)

PLANT HEIGHT

The mean plant height in control plants was 69.16 cms while as in MMS treated plants, the plant height ranged between 78.38-96.56 cms and in SA treated plants the height ranged between 74-80.14 cms. The maximum height was observed in 0.04% MMS concentration. All treated plants showed increase in height of plants compared to control.

FERTILE BRANCHES PER PLANT

Mean fertile branches in control plants was observed as 3 while as in MMS treated plants, it ranged between 1.80-5.40 and in SA treated plants it ranged between 1.40-4.80.

100-SEEDS WEIGHT

Mean 100-seeds weight in control plants was 3.69 gms while as in MMS treated plants, it was observed between 2.34-4.70 gms and in SA treated plants it was observed as 2.84-4.01 gms.

Quantitative characters of a plant are the most important characters associated with yield. Data on quantitative characters did not show regular trend but followed an irregular trend. But lower concentrations of MMS treated plants showed more positive response than SA treated plants.

The lower concentrations inducing growth promoting effects has already been reported by many workers in different plants[11-12]. The different variable response has already been reported by Wani and Khan in *Vigna radiate*[13]. Waghmare and Mehra [14] also reported increase in number of branches in *Lathyrus sativus* after mutagenesis.




Plate-1

A: Late anaphase (Control), B: Cytomixis, C: Laggard formation at anaphase-I, D: Early anaphase, E: Disturbed metaphase-I, F & G: Disturbed telophase-I, H: Ring bivalents, I: Multivalents



Plate-2

A: Control Plant, B: High Yielding Bushy Plant (0.03% MMS), C: Spike Length (Control, 0.04% MMS, 0.02% MMS)

Table 1: Effect of different concentrations of MMS and SA on seed germination of *Hordeum vulgare* L. in M₁ generation.

Treatment	Seed germination (%)		Inhibition (%)	
	In Pots	In Petriplates	In Pots	In Petriplates
Control	100	100	-	-
0.01% MMS	92.89	96.87	7.11	3.13
0.02% MMS	90.30	93.75	9.70	6.25
0.03% MMS	86.47	90.62	13.53	9.38
0.04% MMS	79.51	93.75	20.49	6.25
0.01% SA	91.36	96.87	8.64	3.13
0.02% SA	82.91	84.37	17.09	15.63
0.03% SA	85.25	93.75	14.75	6.25
0.04% SA	90.30	96.87	9.70	3.13

Table 2: Frequency of meiotic abnormalities in various mutagenic treatments of MMS and SA of *Hordeum vulgare* L. in M₁ generation.

	Total No. of PMC's observed	Meiotic abnormalities					
		Dm	Metaphase Rb	Mt	Anaphase Lg	Telophase Dp	Ct
Control	444	-	-	-	-	-	-
0.01% MMS	438	0.22	-	-	0.45	0.45	0.68
0.02% MMS	436	0.45	0.22	-	0.68	-	0.45
0.03% MMS	440	0.68	-	0.45	0.22	0.45	0.22
0.04% MMS	435	0.91	0.45	0.22	-	-	0.68
0.01% SA	434	-	-	-	0.46	0.23	0.69
0.02% SA	432	0.23	-	-	0.69	0.46	0.23
0.03% SA	437	0.45	0.45	0.45	0.22	0.22	-
0.04% SA	430	-	0.69	0.46	0.23	-	0.69

Dm: Disturbed metaphase, Rb: Ring bivalents, Mt: Multivalents, Lg: Laggard, Dp: Disturbed polarity, Ct: Cytomixis

Table 3: Effect of different concentrations of MMS and SA on quantitative characters of *Hordeum vulgare* L. in M₂ generation.

Treatment	Plant height(cm) $\bar{x} \pm S.E$	Fertile branches/plant $\bar{x} \pm S.E$	100-seeds weight(g) $\bar{x} \pm S.E$
Control	69.16 \pm 1.10	3.0 \pm 0.76	3.69 \pm 0.26
0.01% MMS	86.96 \pm 1.52	3.20 \pm 0.33	4.17 \pm 0.007
0.02% MMS	78.38 \pm 0.53	4.00 \pm 0.63	4.14 \pm 0.006
0.03% MMS	80.26 \pm 3.80	5.40 \pm 0.82	2.34 \pm 0.011
0.04% MMS	96.56 \pm 0.46	1.80 \pm 0.17	4.70 \pm 0.013
0.01% SA	74.00 \pm 1.81	1.40 \pm 0.21	3.91 \pm 0.014
0.02% SA	76.60 \pm 3.21	4.80 \pm 0.76	2.84 \pm 0.010
0.03% SA	80.14 \pm 0.45	3.40 \pm 0.35	3.91 \pm 0.009
0.04% SA	79.76 \pm 0.55	3.80 \pm 0.33	4.01 \pm 0.012

CONCLUSION

The present experiment was conducted to induce variability in *Hordeum vulgare* L. The mutagens used were methyl methanesulphonate and sodium azide. MMS at lower concentrations produced less frequency of meiotic abnormalities in M₁ generation that can be tolerated by plant while M₂ generation plants showed more positive response in quantitative characters at lower concentrations of MMS than that of SA. So, it is

better to use MMS at lower concentrations than SA to increase the favourable mutations in plant.

ACKNOWLEDGEMENT

The authors are thankful to the chairman, Department of Botany, Aligarh Muslim University, Aligarh for providing facilities for carrying out this work.

REFERENCES

1. Global Hunger Index; The challenge of hunger: Building resilience to achieve food and nutrition security. 2013.
2. Quinde Z, Ullrich SE, Bailk BK; Genotypic variation in colour and discoloration potential of barley-based food products, Cereal Chemistry, 2004; 83:752-758.
3. Akhtar A, Ansari MYK, Hisamuddin A, Robab MI; Induced variations in quantitative traits by EMS and SA treatments in *Linum usitatissimum* L. Archives of Phytopathology and Plant Protection, 2012; 45(6):667–671.
4. Gulfishan M, Khan AH, Jafri IF, Bhat TA; Assessment of mutagenicity induced by MMS and DES in *Capsicum annuum* L, 2012;19(2): 251–255.
5. Kumar S, Dubey DK; Effects of gamma rays, EMS and DES on meiosis in *Lathyrus sativus*. J. Cytol. Genet, 1998; 33:139-147.
6. Vinita S, Kumar G; Meiotic studies in two cultivars of *Cicer arietinum* L. after EMS treatment. Cytologia, 2004; 69:243-248.
7. Kumar G, Tripathi A; Comparative mitotoxicity and mutagenicity of two heavy metals in *Allium cepa* L. J. Cytol. Genet, 2003; 4:169-173.
8. Maslam SSM; Molecular, ultra structural and cytological characteristics of *Allium cepa* meristems treated with some medicinal extracts. MS.c. (Cytology) Faculty of Girls, Ain Shams Uni., Cairo, Egypt. 2004.
9. Atef AAH, Abd EL-Hamid NR, Abd EL-Hady EA, AL-Ansary AM; Cytogenetic effect of insecticide Tellition and Fungicide Dithane M 45 on meiotic cells and seed storage proteins of *Vicia faba*. Journal of American Science, 2011; 7:19-25.
10. Kumar V, Sinha SSN; Genotoxic effects of two pesticides (Rogor and Bavistin) and an antibiotic (Streptomycin) in meiotic cells of grass pea (*Lathyrus sativus*). Cytologia, 1991; 56:209-214.
11. Khan S, Goyal S; Mutation genetic studies in mungbean IV. Selection of early maturing mutants. Thai. J. Agric. Sci, 2009; 42:109-113.
12. Kharkwal MC; Induced mutations in chickpea (*Cicer arietinum* L.). VI. Significance of induced altered correlations. Indian J. Genet, 2003;63:219-224.
13. Wani MR, Khan S; MMS and SA induced genetic variability for quantitative traits in mungbean, Indian J. Pulses Res, 2006; 19: 50-52.
14. Waghmare VN, Mehra RB; Induced genetic variability for quantitative characters in grasspea (*Lathyrus sativus* L.). Indian J. Genet, 2000; 60: 81-87.