

Research Article**Study on variability in the location of nutrient foramen in fibular diaphysis of Indian population**Dr. Gopalakrishna. K^{1*}, Dr. Shrikrishna. G. Kulkarni², Dr. BS. Rathna³¹Assistant Professor, ²Associate professor, ³Professor and Head,

Department of Anatomy, Malabar Medical College and Research Centre, Modakkallur, Atholi, Calicut- 673315. Kerala, India.

***Corresponding author**

Dr. Gopalakrishna. K

Email: gkemail01@gmail.com

Abstract: The location of nutrient foramen is particular and variable to each bone. It is important in orthopedic surgery, forensic science, anthropology and anatomy. This study was aimed to evaluate the variation in the location of nutrient foramen in fibular diaphysis of right and left side and its relation with length of bone. Materials and methods: Simple random sample of 116 unpaired fibulas [R-58, L-58] of Indian population were used. The student's t-test and Pearson's correlation coefficient were used to evaluate the objectives. Results: Totally 116 nutrient foramen were found. Distance of nutrient foramen from the proximal end was 16.45±2.83cm [R] and 14.70±2.71cm [L]. Total length of bone was 35.16±3.32 [R] and 34.35±3.10 [L]. Foramina index was 46.31±3.89 [R] and 42.23±4.22 [L]. Conclusion: Significant variability in the location of nutrient foramen in fibular diaphysis was observed between right and left side groups. Strong and linear positive correlation was found between the location and total length of bone.

Keywords: Fibula, Nutrient Foramen, Index, Diaphysis, Location.

INTRODUCTION:

Fibula is a non weight bearing bone on lateral part of the leg. Its nutrient foramen is usually located proximal to the midpoint of the posterior surface of shaft and directed distally [1]. Its location is particular for each bone [2] and usually on the flexor aspect [3]. Location is variable and may alter during growth [4] in mammals. Location of nutrient foramen where nutrient artery enters the bone is important in anatomy, orthopedic surgery (including bone graft and treatment of fracture), forensic science and anthropology. Aim of this study was the evaluation of variation in the location

of nutrient foramen in fibular diaphysis of right and left side and its relation with length of bone.

MATERIALS AND METHODS:

This cross sectional primary research study was done on one hundred and sixteen fibula (n=116), not paired, of Indian origin. They were collected with following criteria and study was performed in the Department of Anatomy, Malabar Medical College and Research Centre, Modakkallur, Calicut, Kerala, India, during February 2014 to January 2015.

Table-1: Summary of methods and materials

| | | |
|----|----------------------------|---|
| 1 | Sample size (n) | 116 (right-58, left-58), not paired. |
| 2 | Unit of Investigation | Dry Fibula bone |
| 3 | Study population | Indian |
| 4 | Pilot study | On thirty bones |
| 5 | Calculation of Sample size | With minimum 80% power |
| 6 | Inclusion criteria | Bone in regular form |
| 7 | Exclusion criteria | Deformed and fractured bone |
| 8 | Method | Simple random sampling |
| 9 | Instruments & materials | Vernier Caliper, Magnification hand lens, divider, camera, tables of random digits, fine metallic wire |
| 10 | Parameters | (1). Location: distance between proximal end of the bone to the nutrient foramen [D-NUF]. (2). Total length (TL) [5] (3). Foramina index [6] (FI)=(DNUF/TL)*100 |




Fig-1: Nutrient foramen in fibular diaphysis directed to distal end.

Procedure

After careful inspection, nutrient foramen [fig-1] was identified in study [7] by Carroll SE. The measurements of parameters were taken carefully by single investigator. Average of three reading was taken and entered in the corresponding right and left side groups. Master chart was prepared and data analysis was done.

Statistical assessment [8, 9]:

The descriptive measures [table-1] and inferential calculations were done manually. Frequency

distribution [figure-2] was observed and data exploration was carried out. The student t-test and Pearson’s correlation coefficient were used [figure-2, 3] to evaluate the objectives in right and left side groups. Statistical assessment with p-values < 0.05 was considered significant.

RESULT:

Totally 116 nutrient foramina were found in this study. The results of present study were shown in the following tables and figures.




Fig-2: Frequency distribution of distance of nutrient foramen.

Frequency distribution [8, 9]: The frequency distribution curve of location of nutrient foramen (Figure-2) is symmetrical and uni modal with negligible skewness to right side. Outliers were not observed in the data set. Values for location of nutrient foramen

were distributed around the mean (15.58) with depth of 15.62 and spread from 8.02cm to 29.98 cm with 2.75 cm standard deviation. Values of mean and median are approximate with each other.

Table-1: Summary statistics of present study (n=116)

| No | Title | Mean | SD | SE | Median | Min | Max |
|----|-----------|-------|------|------|--------|-------|-------|
| 1 | D-NUF:[R] | 16.45 | 2.83 | 0.38 | 16.11 | 9.01 | 23.98 |
| 2 | D-NUF:[L] | 14.70 | 2.71 | 0.34 | 15.12 | 8.02 | 21.76 |
| 3 | TLB:[R] | 35.16 | 3.32 | 0.43 | 35.54 | 27.03 | 43.16 |
| 4 | TLB:[L] | 34.35 | 3.10 | 0.40 | 34.53 | 26.86 | 42.08 |
| 5 | FI:[R] | 46.31 | 3.89 | 0.51 | 45.68 | 33.33 | 55.91 |
| 6 | FI:[L] | 42.23 | 4.22 | 0.55 | 43.21 | 29.85 | 52.73 |

R= Right, L=Left, D-NUF= Distance of nutrient foramen from proximal end,
 TLB=Total length of bone, FI=Foramen index, SD=Standard Deviation, SE=Standard Error,
 Min= Minimum, Max= Maximum. [Measurements in cm]

The half the (mean) length of the fibula is 17.85cm [R] and 17.17cm [L]. So the nutrient foramen is proximal to the middle of fibular diaphysis.

Comparison right and left side groups: Significant variability in the location was found [table-2] between the bilateral groups at 5% type-1 error.

Table-2: Comparison right and left side groups: by student t-test

| Description | Group-[Right] | Group-[Left] |
|-------------------------|------------------------------|--------------|
| Mean | 16.45 | 14.70 |
| P value [two tail] | 0.049 | |
| Significance | Significant at $\alpha=0.05$ | |
| 95% confidence interval | 0.714 to 2.786 | |

Table-3: Relationship between total length and location of nutrient foramen

| Side | Right side | Left side |
|------------------------------|------------------------------|------------------------------|
| Correlation coefficient [r] | 0.926 | 0.903 |
| Regression model | $y=17.29+1.085x$ | $y=18.91+1.050x$ |
| Coefficient of determination | 0.857 | 0.841 |
| P value | $P<0.05$ | $P<0.05$ |
| Significance | Significant at $\alpha=0.05$ | Significant at $\alpha=0.05$ |

Correlation [8, 9]: The length and position of foramen in each bone were shown significant relationship on right ($r=0.926$) and left side ($r=0.903$) [table-3].The correlation between bivariate values for

the length and location of nutrient foramen in each bone is explored with Scatter plot. Positive and linear correlation is observed on both sides [fig-3, 4].




Fig 3: Scatter diagram with least square line and regression model [Right side Fibula].




Fig 4: Scatter diagram with least square line and regression model [Left side Fibula].

DISCUSSION:

In the present study the mean distance of nutrient foramen from the proximal end was 16.45cm [range 9.01 to 23.98cm] on right side and 14.70cm [range 8.02 to 21.76cm] left side. Significant difference (p=0.049) was observed between right and left side

groups. The location of nutrient foramen and total length of bone showed strong and linear positive correlation (r=0.926) right and (r=0.903) left side groups. The foramina index was 46.31±3.89 on right side and 42.23±4.22 on left side.

Table 4: Comparison of researches on different populations

| Researcher | (n=) | Population | D-NUF | TL | FI |
|----------------------------|------|--------------------|---------------------------------|---------------------------------|---------------------------------|
| Present study | 116 | Indian | 16.45±2.83[R], 14.7±2.71[L] | 35.16±3.32[R], 34.35±3.32[L] | 46.31±3.89[R], 42.23±4.22[L] |
| E. Gumusburun et al.; [10] | 305 | Turkey | - | 34.56 ± 0.14 | 48.13±0.46 [23-70] |
| Collipal E et al.; [11] | 40 | Chile | 17.23±3.21[R], 17.38±3.02[L] | 35.14±2.34[R], 34.10±2.74[L] | - |
| K.W. Ongeti et al.; [12] | 200 | Kenya, Nairobi. | 15.3 ± 2.4 | 36.5 ± 3.0 | - |

(n)=sample size, (D-NUF)=Distance of nutrient foramen from proximal end, (TLB)=total length of bone, (FI)=Foramina index

COMPARISON

1. Location of nutrient foramen: both intra-population and interpopulation variability can be observed [Table-4] regarding location and foramina index. But the total lengths of bones were not showing much difference in different populations. The tendency of nutrient foramen proximal to the midpoint is evident [12] and in present study. Incidence of nutrient foramina on the middle one third of the fibula was recorded by different scholars [13, 14, 10, 15]
2. Sufficient information was not available about the relationships between lengths to location in the previous literatures. Hence this feature was not compared.
3. Developmental evidence on blood vessel and bone formation [1, 16, 17]:
4. Development of blood vessel will occur in two phases: (1) vasculogenesis (2) angiogenesis.
5. When a tissue grows, blood vessels will proliferate. Initial capillary network will be formed. Ossification of fibula begins at eighth intrauterine

week. Vasculogenesis (embryonic), angiogenesis (post embryonic) and ossification are regulated by multiple growth factors like- Vascular Endothelial Growth Factor (VEGF), Fibroblast Growth Factor (FGF), Fibroblast Growth Factor Receptors (FGFRs), Bone Morphogenic Proteins (BMPs), Platelet-Derived Growth Factor (PDGF), Transforming Growth Factor-β (TGF-β) and tissue hypoxia. These growth factors will bind to specific receptors and deliver signals to the target cells by signaling mechanisms in the formation of adult bone and blood vessels. Later the reconstruction or modeling and stabilization of vessels will take place to meet the local demand by the tissues.

SIGNIFICANCE

- Location of nutrient foramen is variable [2]. It is needed in following fields-
1. In calculation of the length of bone and height of person in forensic science and anthropology [3].
 2. In orthopedic surgical procedures, fracture repair, healing process [3, 7] of fracture and bone graft.

Fibula is the most suitable for bone auto graft because of length, shape of bone [18, 19] and with lower donor site morbidity [20].

3. In the reconstruction of other bones like mandible, spine and tibia [21].
4. Prognosis: Any damage to the precise area of nutrient vessel or foramina will affect the healing process [7]. The rate of healing of a fracture is related to the vascular supply of the bone. The segment of bone with good blood supply is more rapidly healed than those with poor blood supply. Hence this information is helpful regarding assessment of prognosis of healing of bone fracture or graft [10, 13, 15].
5. In vascularised fibula bone graft [1, 12, 22].

CONCLUSION

Significant variability in the location of nutrient foramen in fibular diaphysis was observed between right and left side groups. Strong and linear positive correlation was found between the location and total length of bone.

The limitations of study:

The angiography, radiographs were not used. Age and gender wise classification were not done.

ACKNOWLEDGEMENTS

The authors are grateful to researchers and publishers of the articles and text books cited in the references section of this article. Authors gratefully acknowledge all the authorities of institution, teaching staff, non-teaching staff and all the students for extending valuable support, guidance and cooperation. Authors sincerely respect Rangu for motivating the research activities.

REFERENCES

1. Standring S; Gray's Anatomy, The anatomical basis of clinical practice, 40th edition. Churchill livingstone, Elsevier. 2008; 92-96, 1411-1428.
2. Payton CG; The position of the nutrient foramen and direction of the nutrient canal in the long bones of the madder-fed pig. J Anat. 1934; 68(Pt 4): 500–510.
3. Longia GS, Ajmani ML, Saxena SK, Thomas RJ; Study of diaphyseal nutrient foramina in human long bones, Acta Anat (Basel). 1980; 107(4): 399–406.
4. Henderson RG; The position of the nutrient foramen in the growing tibia and femur of the rat. J Anat. 1978; 125: 593-599.
5. Forriol Campos F, Pellico LG, Alias MG, Fernandez-Valencia R; A study of the nutrient foramina in human long bones. Surg. Radiol Anat. 1987; 9: 251-5.
6. Hughes H; The factors determining the direction of the canal for the nutrient artery in the long bones of mammals and birds. Acta Anat. 1952; 15: 261-80.
7. Carroll SE; A study of the nutrient foramina of the humeral diaphysis. J Bone Joint Surg. 1963; 45-B: 176-81.
8. Carol E Osborn. Basic Statistics for health information management technology. Johnes and Bartlett publications, Inc. Sudbury, Massachusetts. 2008; 109-137, 143-158, 166-185, 193-202.
9. Burt Gerstman B; Basic Biostatistics: Statistics for public health practice. Johnes and Bartlett publications, Inc. Sudbury, Massachusetts .2008; 35-56, 63-83, 129-138, 235-246, 295-332.
10. Gumusburun E, Adiguzel E, Erdil H, Ozkan Y, Gulec E; A Study of the Nutrient Foramina in the Shaft of the Fibula. Folia Anat Jpn. 1996; 73(2-3):125-128.
11. Collipal E, Vargas R, Parra X, Silva E, Del Sol M; Diaphyseal nutrient foramina in the femur, tibia and fibula bones. Int J Morphol. 2007; 25(2):305-308.
12. Ongeti MM, Obimbo PK, Bundi J. Ogeng'o; Anatomical Variation of Position and Location of the Fibula Nutrient Foramen in Adult Kenyans. East African Orthopaedic Journal. 2007; 1:16-18.
13. Mysorekar VR; Diaphysial Nutrient Foramina in Human Long Bones. J Anat. 1967; 101:813-822.
14. Gumusburun E, Yucel F, Ozkan Y, Akgun Z; A Study of the Nutrient Foramina of Lower Limb Long Bones. Surg Radiol Anat. 1994; 16: 409-412.
15. McKee NH, Haw P, Vettese T; Anatomic study of the nutrient foramen in the shaft of the fibula. Clin. Orthop Rel Res. 1984; 184: 141-144.
16. Sadler TW; Langman's medical embryology- 12th ed. Lippincott Williams & Wilkins, a Wolters Kluwer business. Philadelphia, PA 19103. 2012; 63-85, 185-199.
17. Arthur C, Guyton, John E; Hall .Text Book of Medical Physiology, 11th edition. Saunders, Elsevier Inc. India. 2006.
18. Springfield D; Autograft reconstructions. Orthop Clin North (Am). 1996; 27:483-92.
19. Doi K, Tominaga S, Shibata T; Bone grafts with micro vascular anastomoses of vascular pedicles: an experimental study in dogs. J Bone Joint Surg (Am). 1977; 59: 809-15.
20. Lee EH, Goh JC, Helm R, Pho RW; Donor site morbidity following resection of the fibula. J Bone Joint Surg [Br]. 1990; 72: 129-31.
21. Choi SW, Kim HJ, Koh KS, Chung IH, Cha IH; Topographical anatomy of the fibula and peroneal artery in Koreans. Int J Oral Maxillofac Surg. 2001; 4: 329-332.
22. Taylor GI, Miller GD, Ham SJ; The Free Vascularized Bone Graft. Plast & Reconstr Surg. 1975; 55:533-544.