

Research Article

Exploratory study to analyze the relationship of socio demographic parameters with postpartum depression in females at Tertiary care centre

Pragna Sorani¹, Abhijit Khanna², Ankit Moga³, Dhaval Prajapati⁴, Kamlesh Patel⁵, Prakash Mehta⁶

^{1,3}Senior Resident, Department of Psychiatry, C.U. Shah Medical College, Surendra nagar, Gujarat, India

^{2,4}Resident, Department of Psychiatry, C.U. Shah Medical College, Surendra nagar, Gujarat, India

⁵Professor & Head, Department of Psychiatry, C.U. Shah Medical College, Surendra nagar, Gujarat, India

⁶Professor & Head, Department of Psychiatry, GMERS Medical College and Hospital, Sola, Ahmedabad, Gujarat, India.

***Corresponding author**

Dr. Pragna Sorani

Email: drabhijitmkhanna@gmail.com

Abstract: It is an exploratory study to analyze the relationship of socio demographic parameters with depressive features in postpartum females. The study included 100 females who had live birth. The depression score was collected using HAM-D scale on the 30th day of delivery. Results show 18% of females suffering from postpartum depression and multiple socio demographic parameters affecting the mental health of the mothers. From the result, we gain insight in the factors which might be responsible for postpartum depression in females.

Keywords: Socio demographic parameters, post partum depression

INTRODUCTION

Pregnancy is a major life event of every adult female. A pregnant female passes through many changes in the body including physical, biological, hormonal as well as psychological. So overall pregnancy, child birth and entering into motherhood become very stressful event for a female and during this process they need much family support. Birth of a child is considered a moment of joy for any female but while passing through such changes some women may suffer from depression.

Symptoms and criteria to diagnose postpartum depression are similar to that of major depressive episode. In DSM-5[1], criteria for major depression includes at least 2 weeks period of sad mood, or reduced interest in all activities and 4 symptoms out of the following – changes of weight, sleep, psychomotor activity, feelings of guilt, worthlessness, forgetfulness, difficulty in making decisions, lethargy and recurrent suicidal ideas, thoughts or attempts.

Many of the epidemiological studies consider postpartum depression occurring within 6-12 weeks of postpartum period [2]. Researches carried out in India reports the prevalence rate of postpartum depression 5-23 % [4-10]. One study reported the prevalence rate of postpartum depression to be 12.5 % in Gujarat [7].

Researches carried out report many different risk factors that can predispose women towards the

development of depression in post partum period. They are socio demographic factors like age of mother[3, 12], socio economic status [15, 16, 18, 19, 20], area of residence [3], level of education [12], employment [12], obstetric factors like planning of pregnancy [8], mode of delivery[8, 10],gender of infant [5], and parity [22].

OBJECTIVE

The objective of the study is to analyze the relationship of socio demographic parameters with postpartum depression in females at the end of 1st month of delivery in a tertiary centre at Surendra nagar.

METHODOLOGY

The current study was conducted at C.U. Shah medical college, Surendra nagar. Women attending gynecology dept. during their postpartum period after 1 month of the delivery were included in the study. After obtaining written informed consent their socio demographic and obstetric details were collected. HAM-D scale was used to detect presence of depression.

Hamilton Rating Scale for Depression [23] (HAM-D):

HAM-D is a clinician rating scale for rating the severity of depression. HAM – D provides an indication for depression and over time, guide to recovery. It is one of the most commonly used and accepted outcome measures for evaluating the severity of depression symptoms. The HAM – D was designed

to be administered by a trained professional using a semi-structured interview. Even though Hamilton provides no specific guideline for the administration and scoring of the scale or any standardized questions for eliciting information from the patients, high inter-rater reliability has been observed²⁴. It takes about 20-

30 minutes to complete the interview and score the results. Ten items are scored on a 5 point scale ranging from 0 = not present to 4 = severe. Two items are scored from 0-3, and nine items are scored from 0-2. Sum the total to arrive at a conclusion.

RESULTS

Table-1

	Depressed n=frequency (%)	Not Depressed n=frequency (%)	Total n=frequency (%)
Age (in years)			
≤ 20	0 (0)	11 (13.42)	11 (11)
21-30	15 (83.3)	64 (78.05)	79 (79)
31-40	3 (16.7)	7 (8.53)	10 (10)
Total	18 (18)	82 (82)	100 (100)
Education			
Illiterate	5 (27.78)	29 (35.36)	34 (34)
Educated	13 (72.22)	53 (64.64)	66 (66)
Total	18 (18)	82 (82)	100 (100)
Occupation			
Working	3 (16.7)	15 (18.29)	18 (18)
Housewife	15 (83.3)	67 (81.71)	82 (82)
Total	18 (18)	82 (82)	100 (100)
Area of residence			
Urban	4 (22.2)	23 (28.05)	27 (27)
Rural	14 (77.8)	59 (71.95)	73 (73)
Total	18 (18)	82 (82)	100 (100)
Gender of new born			
Male	3 (16.7)	42 (51.21)	45 (45)
Female	15 (83.3)	40 (48.79)	55 (55)
Total	18 (18)	82 (82)	100 (100)
Parity			
Primipara	2 (11.1)	51 (62.19)	53 (53)
Multipara	16 (88.9)	31 (37.81)	47 (47)
Total	18 (18)	82 (82)	100 (100)
Planning about pregnancy			
Planned	8 (44.4)	16 (19.51)	24 (24)
Unplanned	10 (55.6)	66 (80.49)	76 (76)
Total	18 (18)	82 (82)	100 (100)
Mode of delivery			
LSCS	7 (38.9)	32 (39.02)	39 (39)
Vaginal delivery	11 (61.1)	50 (60.98)	61 (61)
Total	18 (18)	82 (82)	100 (100)
Socio economic status			
Upper class	0 (0)	6 (7.32)	6 (6)
Middle class	8 (44.4)	47 (67.07)	55 (55)
Lower class	10 (55.6)	29 (25.61)	39 (39)
Total	18 (18)	82 (82)	100 (100)




Fig-1: Percentage of females having PPD and its relationship with Age of Mother




Fig- 4: Percentage of females having PPD and its relationship with Residence of Mother




Fig- 2: Percentage of females having PPD and its relationship with Education of Mother




Fig- 5: Percentage of females having PPD and its relationship with Gender of new born




Fig- 3: Percentage of females having PPD and its relationship with Occupation of Mother




Fig- 6: Percentage of females having PPD and its relationship with Parity




Fig- 7: Percentage of females having PPD and its relationship with Planning of pregnancy




Fig- 8: Percentage of females having PPD and its relationship with Mode of Delivery




Fig- 9: Percentage of females having PPD and its relationship with Socioeconomic Status

DISCUSSION

Out of the 100 subjects in study, 18 females (18%) were found to be having postpartum depression. In those 18 subjects, we evaluated for the factors which might suggest the probable causes for PPD. 83.3% women from depressed group from the age group 21-30 years showed higher score in HAM-D (Table 1, Figure

1). This corroborates with the previous studies [3, 11, 12]. In relation to education of mother, women who were educated seemed to be more depressed than the women who were illiterate as they constituted 72.22% of the total depressed women (Table 1, Figure 2). This contrasted from the previous studies [12, 17- 20]. We observed that those women who were housewives were more depressed than those who were working and constituted 83.3% of the total depressed women (Table 1, Figure 3). This corroborates with the previous studies [12, 14, 16, 18-20]. Women hailing from rural area were found to be more depressed than females living in urban areas and constituted 77.8% of the total depressed women (Table 1, Figure 4). This corroborates with the previous study³. Women who delivered a female child were found to be more depressed than the females who delivered a male child and constituted 83.3% of the total depressed women (Table 1, Figure 5). These results are similar to those of previous studies [5, 7-10].

Also, it was observed that multiparous mothers were more depressed than primipara mothers and constituted 88.9% of the total depressed females (Table 1, Figure 6). This corroborates with previous studies [6, 7, 13]. There was no much difference found in depression in females in context to planning of pregnancy (Table 1, Figure 7). Those females who delivered via LSCS suffered depression more than those who delivered by vaginal route and constituted 61.1% of the total depressed females (Table 1, Figure 8). These results are similar to those of previous studies [18, 21]. There was no much difference found in depression in females in context to socioeconomic status of the mother (Table 1, Figure 9).

CONCLUSION

More depressive features were noted in women of age group 21-30 years, those who were educated; those were housewives, living in rural areas, who delivered a female child. Females who were multiparous and delivered by LSCS were found more depressed. The limiting factor for present study was its smaller sample size. There was no conflict of interest in the present study.

REFERENCES

1. Diagnostic and Statistical Manual of Mental Disorders, 5th edition, 160-161.
2. Kumar R, Robson K; A prospective study of emotional disorder in childbearing women. Br J Psychiat, 1984, 166:191-195.
3. Lanes A, Kuk JL, Tamim H; Prevalence and characteristics of postpartum depression symptomatology among Canadian women: a cross-sectional study. BMC Public Health, 2011; 11(1), 302.
4. Thangappah Radhabai Prabhu, T V Asokan, Rajeshwari A; "Postpartum Psychiatric Illness" J Obstet Gynecol, 2005; 55(4): 329-332.

5. Chandran M, Tharyan P, Muliylil J, Abraham S; Postpartum depression in a cohort of women from a rural area of Tamil-Nadu, India: Incidence and risk factors” British Journal of Psychiatry, 2002; 181(6): 499-504.
6. Hegde; Postpartum depression: Prevalence and Associated features among women in India. J women’s health, Issue care, 2012; 1: 1.
7. Desai Nimisha, Mehta Ritambhara, Ganjiwale Jai shree; Study of prevalence and risk factors of Postpartum Depression. National Journal of Medical Research, 2012; 2(2): 194-198.
8. Sankapithilu GJ, Nagaraj AKM, Bhat SU, Raveesh BN, Nagaraja V; A comparative study of frequency of postnatal depression among subjects with normal and caesarean deliveries. Online Journal of Health and Allied Sciences, 2010; 9(2).
9. Mamta sood, Sood AK; Depression in Pregnancy and Postpartum period. Indian Journal of Psychiatry, 2003,45(1), 48-51.
10. Vicar Patel, Merlyn Rodriguez, Nadia Desouza; Gender, Poverty and Postnatal Depression. A study of mothers in Goa, India. Am J Psychiatry, 2002; 159: 161.
11. Sword W, Kurtz Landy C, Thabane L, Watt S, Krueger P, Farine D, et al.; Is mode of delivery associated with Postpartum depression at 6 weeks: a prospective cohort study. British Journal of Obstet & Gynecol, 2011; 118(8): 996-977.
12. Bener; “Prevalence of psychiatric disorder and associated risk in women during postpartum period” I J of women’s health, 2012; 4: 191-200.
13. Lanzi RG, Bert SC, Jacobs BK; Depression among a sample of first time adolescent and adult mother, J Child Adolesce Psychiat Nurse, 2009; 22(4): 194-202.
14. Asli Goker, Emre Yanik Kerem; Postpartum depression: Is mode of delivery a risk factor? International Scholary Research Network. ISRN Obis and Gyn, 2012.
15. Leigh B, Milgrom J; Risk factors antenatal depression, postnatal depression and parenting stress. BMC Psychiatry 2008; 8: 24.
16. Mohamed NA, Mahmoud GA; Postpartum depression: Prevalence and predictors among EL Eman’s specialized hospital, Journal of American Science, 2011: 7(12).
17. Horowitz JA, Murphy CA, Gregory KE, Wojcik J; A community based screening initiative to identify mother at risk of postpartum depression. J Obstet Gynecol Neonatal nurs, 2011; 40(1):52-61.
18. Bener A, Burgut FT; A study of Postpartum depression in a fast developing countries: Prevalence and related factors.
19. Khooharo Y, Majeed T, Das C, Majeed N, Majeed N, Choudhry AM; Associated risk factors for postpartum depression presenting at a teaching hospital” Annals of King Edward Medical University, 2010; 16(2): 87-90.
20. Goyal D, Gay C, Lee KA; “How much does LSES increase the risk of pre and postnatal depression symptoms in first time mothers” Women’s Health Issue, 2010; 20(2): 96-104.
21. Fisher J, Asbury J, Asbury J, Smith A; Adverse psychological impact of operative obstetric intervention: a prospective longitudinal study. Australian and Newzeland J Psychiat, 1997; 31: 728-738.
22. Beck CT, Reynolds MA, Rutowski P; Maternity blues and postpartum depression. J Obstet Gynecol, 1992; 21: 287-93.
23. The Hamilton rating scale for depression, Journal of Operational Psychiatry, 1979; 10(2):149-165.
24. Muller MJ, Dragicevic A; Journal of Affective disorder, 2003; 77(1): 165-169.