
Research Article**Floristic Composition and Diversity Assessment of Home garden Plants in a Rural Village, Swamithoppe, Kanyakumari District, Tamil Nadu, India****R. Neelamegam*, K.B. Premkumar, S. Ancel Sowmiya, A.R. Fathima Sabana**

P.G. Department of Botany & Research Centre, S.T. Hindu College, Nagercoil -629002, Kanyakumari District, Tamil Nadu, India.

***Corresponding author**

R. Neelamegam

Email: rmegamsthcnl@gmail.com

Abstract: The main objective of this study was to assess the status, composition and diversity of plants in a rural village homegardens with the help of socio economic factors of households. The study was carried out in Swamithoppe village, Kanyakumari district, Tamil Nadu, India, in 121 randomly selected homegardens were measured. Complex plant species inventories were carried out to assess the number and abundance of plant species (i.e., all useful plant species) and ornamentals. All gardeners were individually interviewed about homegarden management and plant utilization, among other information. In addition, to plant species information, species diversity, richness, evenness and dominance indices were also calculated. In Swamithoppe village, the homegarden area varied from 20.23m² to 627.28m² with an average of 73.21m² to 519.36m². In the HG survey, a total of 119 plant species comes under 108 genus belongs to 58 families with a total number of individual is 3540 were recorded. The number species recorded ranged from 2 to 23/HG species with an average of 8.30 to 9.24species/HG. The species density varied from 1.65/100m² to 64.26/100m² with an average of 2.58 to 13.84/100m². The number of individuals of plant species noted in the homegarden varied from 7 to 78nos with an average of 24.75 to 66.67nos. The overall, plant diversity in homegarden based on number of individuals ranged from 9.88 to 143.35/100m² with an average of 12.99 to 40.06/100m². In plant overall HGs shows a total of 119 species with 3540 individuals in 121 HGs and the plants distributed in HGs ranged from 3 to 22 HGs with an average of 9.39 HG. Out of 58 botanical families recorded Euphorbiaceae is one of the most represented family having 6 genes, 10 species with 245 individuals with an average of 2450 individuals per species and the maximum diversity ($H' = 0.342$) and low Simpson's diversity ($\lambda = 0.00052$) as compared to other families. Among the top 10 HG species, *Cocos nucifera* shows maximum no of individuals (162) with an average of 7.36/HG and found in 22 HGs. The distribution pattern of HG plants indicate that maximum no of plants found in herbs (36.13%) in life form, earthen plants (77.31%) in habitat, cultivated plants (74.79%) in growth condition, whole plant utilization (39.50%) in useful parts and ornamental use plants (31.93%) in uses category. The diversity indices estimated for over on home garden plants as $H' = 4.627$, (Shannon diversity), species richness ($R = 14.440$), species evenness ($E = 0.968$), and Simpson's diversity ($\lambda = 0.0109$).

Keywords: Homegardens, Plant structure, Plant composition and distribution, Plant diversity, Swamithoppe village, Kanyakumari.

INTRODUCTION

Homegardens make a vital contribution to meet various household needs, especially for poor families in developing countries. However, the importance of the biodiversity of homegardens is the availability of varieties, which are found suitable to human beings under a large variety of social, economic and cultural situations. Research on homegardens gaining interest for their potential as models of economically efficient and ecologically sustainable agroforestry system [1-3], and they emphasize the importance of preserving homegardens as key elements in the conservation and generation of diversity in agricultural species. Personal preference, socio-economic status and culture seems to be the main determinants of the appearance, functions and structure of

homegardens [4]. The rural homegardens usually have more layers of plant canopy and thus, are more complex than the urban gardens [5]. Most homegardens research has focused on homegardens as integrated multi-species system, giving greatest attention to the variation of species diversity among homegardens [6-9]. Additionally, several studies have paid special attention to the vertical variation of species by comparing the different layers of canopy strata constituting homegardens [2, 10, 11]. Little or no attention has been given to analysing the horizontal variability within homegardens [12-14]. The aim of present study is to evaluate the floristic composition and diversity of homegarden plants in the Swamithoppe village, Kanyakumari district, Tamilnadu, India.

MATERIALS AND METHODS

Homegardens in the study area, Swamithoppe village, was surveyed from June 2015 to September 2015 to evaluate the status, composition and diversity of plant species. Methods used in this study mainly focussed on determining certain indicators for the assessment of sustainability with respect to socio-economic condition of households and homegarden plant diversity. The present study was carried out in the Swamithoppe village, Kanyakumari District, Tamilnadu, India. Swamithoppe lies about half-way between the cities of Nagercoil and Kanyakumari on the Nagercoil-Kanyakumari road and located at 8.12°N 77.49°E and elevated as 13m (43ft).

Homegarden survey was conducted in 121 households with homegardens were randomly selected. The selected homegardens were categorized into four types based on their nature –as hutted, tiled, and terraced and storied which are fenced or non-fenced in the study area. Households were identified as sampling units for the survey. Questionnaire was prepared to collect various information from the households (respondents) related to home gardening. For this, the actual respondent of the household was identified as one who involved in most of the decision making in the agriculture (homegarden) related matters. In some case the interview was conducted more than one member as well. Each household was interviewed as basic socio-economic data and homegarden specific data. Finally, the filled questionnaires were checked to confirm the competence and quality of the information collected. The filled questionnaires were thoroughly checked and numerical coding of the filled questionnaires for the data entry and calculation of various parameters was done. Before starting plant inventory, the homegarden type (fenced/non-fenced, hut, tiled, terrace, and multistoried) were noted. The size of HG was measured excluding the area occupied by the houses.

The plant species were identified on the basis of vernacular names, published field inventories, floras [15, 16], experts in plants and consulting available herbaria of the region. In the study the plant species recorded have been arranged alphabetically for each species the binomial name first followed by the local name, family, life-forms, potted/earthen plants, cultivated/wild plants, useful parts of the plants and their utilization are recorded (Annexure table I). From the data collected, various diversity indices like Shannon-Weiner diversity index [17, 18], Simpson's diversity index [19], Pielou's species evenness index [20, 21], Margalef species richness index [22, 23] were estimated using the standard methods.

RESULTS AND DISCUSSION

Socio-economic conditions of Households

The homegardens of Swamithoppe village are categorized into 4 types such as hutted, tiled, terraced and storied house which are sub categories into fenced and non-fenced house (Table 1). Out of 121 houses surveyed in the study area, terrace house represent more in number (55nos.) which is about 45.45% and it is followed by storied house (51nos. with 42.14%).

Among the 121 homegardens surveyed, 74.38% of the house (90nos.) were fenced and remaining are non-fenced house were recorded in the Swamithoppe village. The number of family members exists in the range between as 1 to 7nos. Most of the families (34nos. out of 121) have 4 members in the study area and it was 28.09%. About 55.33% of the households had formal education less than 10th standard while 36.36% had above 10th level formal education and 8.26% of respondents showed illiterate. Most of the households were unskilled workers 35.53% followed by 22.31% skilled workers, 14.87% farmers (Table 1).

Most of the households (38.01%) in the study area were under the annual income ranges from Rs.50,000/- to Rs. 1,00, 000/- (maximum) and most of the households (65nos.) out of 121 (53.71%) spent between Rs. 1000 to Rs. 5000 annually for HG maintenance. It was noted that 34.71% of the households (42nos. out of 121) have experience in homegarden works, while 37.88% households were farm workers (41nos.). Among the households, 34.71% households (42) were used the homegardens as washing area and is followed by 22.31% households utilize the homegarden as children play area. Out of 121 homegardens surveyed, the plants grow well in 47 HGs (38.84%), in 50.41% HGs shown plants with moderately grown and the remaining HGs poorly grown plants. About 84 (69.42%) HGs were categorized into two layered and the remaining 30.58% (30nos.) were three layered (Table 1).

Status and structure of homegarden plant

A complete inventory list of the plant species and the detailed species list is presented in (Annexure Table I) along with common name of the plant species, family name, plant type (habit) growth condition (wild/potted), nature of growth (wild/cultivated), useful part, uses were recorded. A total of 3540 individuals from 119 species under 108 genera belongs to 58 families were recorded and the plant species have been taxonomically verified and identified as distinct species (Table 2).

Table -1: Socio-economic characteristics of households recorded during the survey of homegardens in the Swamithoppe village.

| 1. House types | No of homegardens* | % of homegardens** |
|--|---------------------------|---------------------------|
| i. Hutted house | 01 | 00.83 |
| ii. Tiled house | 14 | 11.57 |
| iii. Terrace house | 55 | 45.45 |
| iv. Storied house | 51 | 42.15 |
| v. Non-fenced house | 31 | 25.62 |
| vi. Fenced house | 90 | 74.38 |
| 2. Household members | No of Households* | % of households** |
| i. ≤ 3 members | 46 | 38.02 |
| ii. 4 to 5 members | 54 | 44.63 |
| iii. ≥ 6 members | 21 | 17.35 |
| 3. Educational status of Households | No of households* | % of households** |
| i. Non-formal education | 10 | 8.27 |
| ii. Formal education up to 10 th level | 67 | 55.37 |
| iii. Formal education above 10 th level | 44 | 36.36 |
| 4. Occupation of Households | No of households* | % of households** |
| i. Farmers | 18 | 14.88 |
| ii. Businessman | 14 | 11.57 |
| iii. Professionals | 19 | 15.70 |
| iv. Skilled workers | 27 | 22.31 |
| v. Unskilled workers | 43 | 35.54 |
| 5. Annual Income of Households | No of households* | % of households** |
| i. Up to Rs. 50, 000/= | 25 | 20.66 |
| ii. Rs. 50, 000/= to 1, 00, 000/= | 46 | 38.02 |
| iii. Rs. 1, 00, 000/= to 2, 00, 000/= | 24 | 19.83 |
| iv. Rs. 2, 00, 000/- to 5, 00, 000/= | 26 | 21.49 |
| 6. Annual Homegarden Expenditure | No of households* | % of households** |
| i. Up to Rs. 1, 000/= | 45 | 37.19 |
| ii. Rs. 1, 000/= to 5, 000/= | 65 | 53.72 |
| iii. Rs. 5, 000/= to 10, 000/= | 11 | 09.09 |
| 7. Activity of Household Members | No of households* | % of households** |
| i. Farm workers | 41 | 33.88 |
| ii. Non-farm workers | 27 | 22.31 |
| iii. Experience in homegarden works | 42 | 34.71 |
| iv. No experience in homegarden works | 11 | 09.09 |
| 8. Homegardens used for other purposes | No of households* | % of households** |
| <u>A. Used as social/ living area</u> | | |
| i. Rest or meeting area | 10 | 08.26 |
| ii. Children's play area | 27 | 22.31 |
| iii. Flower garden | 21 | 17.36 |
| <u>B. Used as Physical/ utility area</u> | | |
| i. Storage area | 11 | 09.09 |
| ii. Washing area | 42 | 34.71 |
| iii. Drying area | 10 | 08.26 |
| 9. Growth condition of HG plants | No of homegardens* | % of homegardens** |
| i. Well grown | 47 | 38.84 |
| ii. Moderately grown | 61 | 50.41 |
| iii. Poorly grown | 13 | 10.74 |
| 10. Layering pattern of HG plants | No of homegardens* | % of homegardens** |
| i. Two layered | 84 | 69.42 |
| ii. Three layered | 37 | 30.58 |
| Total^{*/**} | 121* | 100.00** |

Table -2: Most represented botanical families in number of species, number of genus and number of individuals recorded in the homegardens of Swamithoppe village.

| Family | Number of Species (%) /Family | Number of Genus (%) /Family | Number of Individuals (%) /Family | Number of Individuals (%) /Genus | Number of Individuals (%) /species |
|------------------|-------------------------------|-----------------------------|-----------------------------------|----------------------------------|------------------------------------|
| Euphorbiaceae | 10 (8.40) | 6 (5.56) | 245 (6.92) | 40.83 (1.15) | 24.50 (0.69) |
| Solanaceae | 7 (5.88) | 3 (2.78) | 170 (4.80) | 56.67 (1.60) | 24.29 (0.69) |
| Rubiaceae | 5 (4.20) | 4 (3.70) | 183 (5.17) | 45.75 (1.29) | 36.60 (1.03) |
| Leguminaceae | 4 (3.36) | 4 (3.70) | 72 (2.03) | 18.00 (0.51) | 18.00 (0.51) |
| Cucurbitaceae | 4 (3.36) | 4 (3.70) | 84 (2.37) | 21.00 (0.59) | 21.00 (0.59) |
| Amaranthaceae | 4 (3.36) | 3 (2.78) | 135 (3.81) | 45.00 (1.27) | 33.75 (0.95) |
| 10-Families with | 3 (2.52) | 30 (2.78) | 1051 (2.97) | 35.03 (0.99) | 35.03 (0.99) |
| 13-Families with | 2 (1.68) | 25 (1.78) | 783 (1.70) | 31.32 (0.88) | 30.12 (0.85) |
| 29-Families with | 1 (0.84) | 29 (0.93) | 817 (0.79) | 28.17 (0.80) | 28.17 (0.80) |
| Total -58 | 119 (100.00) | 108 (100.00) | 3540 (100.00) | | |

The homegarden survey reveals that out of 58 families recorded, 29 families have single species, 13 families have two species, 10 families have 3 species, 3 families have 4 species, Rubiaceae family have 5 species, Solanaceae have 7 species and Euphorbiaceae have 10 species. Among the families, Euphorbiaceae represent 8.40% with 5.56% genus (6), 6.92% individuals (245nos.) with an average of 24.50 individuals per species (Table 2). Among the top 5 families recorded, Euphorbiaceae shows maximum

number of species (10sps.), maximum number of individuals (245nos.) and maximum species diversity index ($H' = 0.342$), while *Arecaceae* shows more Simpson's diversity index ($\lambda = 0.00230$) as compared to other families. Among the top 5 species, out of 119 species recorded, *Cocos nucifera* have maximum number of individuals (7.36 per homegarden) and in more number of homegardens (22nos.) as compared to other plants (Table 3).

Table -3: Top 5 families having maximum Simpson's diversity index (λ) and Shannon-Weiner's species diversity index (H') of homegarden species surveyed in the study area.

| Sl. No. | Name of the Family | Species diversity Index (H') | Sl. No. | Name of the Family | Simpson's Diversity Index (λ) |
|---------|--------------------|----------------------------------|---------|--------------------|---|
| 1 | Arecaceae | 0.00230 | 1 | Euphorbiaceae | 0.342 |
| 2 | Rubiaceae | 0.00055 | 2 | Rubiaceae | 0.236 |
| 3 | Euphorbiaceae | 0.00052 | 3 | Arecaceae | 0.233 |
| 4 | Malvaceae | 0.00040 | 4 | Solanaceae | 0.192 |
| 5 | Amaranthaceae | 0.00039 | 5 | Amaranthaceae | 0.176 |

Table -4: Top 5 HG plants showing more number of individuals and found in more number of homegardens of Swamithoppe village.

| Sl. No. | Name of the species | No of In/Sp | Name of the species | Av No of In/Sp/HG | Name of the species | Max No of HG |
|---------|------------------------------------|-------------|-----------------------------------|-------------------|-------------------------------|--------------|
| 1 | <i>Cocos nucifera</i> | 162 | <i>Cocos nucifera</i> | 7.36 | <i>Cocos nucifera</i> | 22 |
| 2 | <i>Abelmoschus esculentus</i> | 61 | <i>Ocimum sanctum</i> | 6.67 | <i>Bambusa arundinacea</i> | 19 |
| 3 | <i>Ocimum sanctum</i> | 60 | <i>Nerium indicum</i> | 5.43 | <i>Abelmoschus esculentus</i> | 18 |
| 4 | <i>Zingiber officinale</i> | 57 | <i>Stachytarpheta jamaicensis</i> | 5.25 | <i>Zingiber officinale</i> | 17 |
| 5 | <i>Amorphophallu spaenoiifolus</i> | 51 | <i>Polyalthia longifolia</i> | 4.64 | <i>Dracaena sps.</i> | 16 |

No –Number; In –Individuals; Sp –Species; HG –Homegarden; Max –Maximum; Av –Average;

Out of 119 species recorded the top 5 species having maximum number of individuals noted in the homegardens are *C. nucifera*, *A. esculentus*, *O. sanctum*, *Z. officinale*, *A. paenoiifolus*. Among these plants, *C. nucifera* have maximum number of individuals (162) found in 22 homegardens with an average of 7.36 individuals per homegarden (Table 4).

The homegarden surveyed in the Swamithoppe village were categorized into three types such as small (<200m²), medium (200 to 400m²) at large (>400m²) based on homegarden area and the data collected were compared in different parameters like number of plant species, species family number of individual plants of their (plant) family and homegarden of area (Table 5).

Out of 121 homegardens, 102 HGs (84.30%) comes under small HGs, while 13 HGs (10.74%) under medium HGs and 6 HGs (4.96%) under large HGs. The overall area surveyed was ranged from 20.23% to 627.28m² with an average area of 116.06m²/HG. About 53.17% of HG area comes under small HGs while it

was 24.64% area under medium HGs and 22.19% area comes under large HGs. Maximum average HGs was noted in large HGs (519.36m²/HG) and it is followed by medium HGs (266.17 m²/HG) and it was low in small HGs (73.21 m²/HG) (Table 5).

Table -5: Comparison of minimum, maximum and average of homegarden plants based on number of species, number of individuals and density in the study area.

| Parameters | Minimum | Maximum | Average | Total Number & % |
|---|---------|---------|---------|-------------------|
| Homegarden area (m²) | | | | |
| Small HGs (102 HGs.) | 20.23 | 182.11 | 73.21 | 7467.25 (53.17) |
| Medium HGs (13 HGs.) | 202.35 | 343.99 | 266.17 | 3468.16 (24.64) |
| Large HGs ((6 HGs.) | 445.17 | 627.28 | 519.36 | 3 16.18(22.19) |
| Overall (121 HGs) | 20.23 | 627.28 | 116.06 | 14043.59 (100.00) |
| Number of Species in HGs | | | | |
| Small HGs (102 HGs.) | 2 | 23 | 8.39 | 856 (76.57) |
| Medium HGs (13 HGs.) | 6 | 21 | 14.08 | 183 (16.37) |
| Large HGs ((6 HGs.) | 8 | 17 | 13.17 | 79 (7.06) |
| Overall (121 HGs) | 2 | 23 | 9.24 | 1118 (100.00) |
| Species Density/100m² | | | | |
| Small HGs (102 HGs.) | 4.12 | 64.26 | 13.84 | 1412.11 (94.25) |
| Medium HGs (13 HGs.) | 2.28 | 9.39 | 5.44 | 70.72 (4.72) |
| Large HGs ((6 HGs.) | 1.65 | 3.82 | 2.58 | 15 .46(1.03) |
| Overall (121 HGs) | 1.65 | 64.26 | 12.38 | 1498.29 (100.00) |
| Number of individuals in HGs | | | | |
| Small HGs (102 HGs.) | 7 | 47 | 24.75 | 2524 (71.30) |
| Medium HGs (13 HGs.) | 37 | 65 | 47.38 | 616 (17.40) |
| Large HGs ((6 HGs.) | 48 | 78 | 66.67 | 400 (11.30) |
| Overall (121 HGs) | 7 | 78 | 29.26 | 3540 (100.00) |
| Plant Density/100m² | | | | |
| Small HGs (102 HGs.) | 11.53 | 143.35 | 40.06 | 4086.31 (92.86) |
| Medium HGs (13 HGs.) | 13.66 | 26.69 | 18.16 | 236.02 (5.36) |
| Large HGs ((6 HGs.) | 9.88 | 16.17 | 12.99 | 77.95 (1.77) |
| Overall (121 HGs) | 9.88 | 143.35 | 36.37 | 4400.29 (100.00) |

Small HGs: <200m²; Medium HGs: 200 to 400m²; Large HGs: >400m²;

The range of plant species in small HGs was 2 to 23 with area average of 8.39 species while in medium HGs it was ranged from 6 to 21 with an average of 14.08 species and 8 to 17species in large HGs with an average of 13.17species. Thus, the average number of species per HGs is higher in medium HGs and large HGs as compared to small HGs. However, the total number of species found in small HGs was higher (76.57%) when compared to medium HGs (16.37%) and large HGs (7.06%).The species density is small HGs was ranged from 4.12 to 64.26/ 100m² with an average species density of 13.84/m². In medium HGs, it was ranged from 2.28/100m² with an average species determination of 5.44/100m² are in large HGs it was ranged from 1.65 to 3.82/100m² with average species density 2.58/100m². The species density/100m² was higher (94.25%) in small HGs density compared to medium HGs and large GH. Among the HGs, maximum species density (94.25%) was noted in small HGs as compared to medium 4.72% in large HGs (1.03%) (Table 5).

The number of individuals of HGs plant species recorded was ranged from 7 to 78nos. with an average of 29.26 individuals per HGs in general. This range was higher in large HGs (48 to 78nos. with are average of 66.67nos) as medium HGs (37 to 65nos. with an average of 24.75nos.). Among the HGs, maximum number of individuals (71.30%) was recorded in small HGs while it was only 17.40% in medium HGs at 11.30% in large HGs. The plant (vegetation) density/100m² in general was ranged from 9.88/100m² to 143.35/100m² with an average of 36.37/100m² HGs. About 92.86% plant density was noted in small HGs as it ranged from 11.53/100m² to 143.35/100m² HGs while it was lower in medium large HGs (Table 5).

Distribution pattern of HG plants

The plant distribution pattern of HG species and their number of individuals, the range and average number of HGs in which the plants present were categorised based on the life form s (climbers, creepers, herbs, shurbs and trees), habitat (earthern/potted), nature of growth (cultivated/wild), usefull parts (flower,

fruit, leaf, seed, stem, wood and wholeplant) and uses (edible, fuel, medicine, multipurpose, ornamental, and vegetable). Based on life form category, maximum number of species (43nos .) in herbs (36.13%) and the number of individuals noted as 1235 (34.89%). The HGs, in which the species present was ranged from 5 to 18 HGs with an average of 9 to 16/HG. But, the range and average number of HGs in which the species present was more in case of tree species which found in 6 to 22 HGs with an average of 10.32 HGs (Table 6).

In the Habitat (Earthen/Potted plants) category, most of the HG plants (77.31%) were earthen plants (92 species) while the potted plants were only 16 species (21.85%) and both earthen and potted plants have only 1 species (0.84%). The number of individuals found more (77.94%) , i .e., earthen plants (2759nos .) and it was 756 nos. (21.36%) noted in potted plants. The minimum and maximum number of HGs in which the species present in case of earthen plants was 3 to 22

HGs with an average of 9.35 HGs, while to potted plants found in 6 to 18 HGs with an average of 9.65 HGs, and the earthen and potted plants found only in 7 HGs (Table 6).

Based on Growth Conditions (cultivated/wild), cultivated plants represented 74.79% (89 species) , then the wild plants (2 species; 68%) in HG's surveyed, whereas both cultivated and wild plants noted as 23.53% (28 species). The total number of individuals of HG plants shows 78.28% (2771 nos .) in cultivated category and it was 66nos . (1.86%) only in wild category, while both represented as 19.86% (703 nos .). The number of HGs in which the cultivated species present ranged from 3 to 22 HGs with an average of 9.64 HGs and it was 7 to 9 HGs with an average of 8 HGs for wild plants. The presence of both cultivated and wild were ranged from 5 to 16 HGs with an average of 8.71 HGs (Table 6).

Table -6: Distribution categories of homegarden species and their individuals in the study area surveyed.

| Plant distribution categories | No. sp. (%) | No of Indi. (%) | No of HGs in which the species present | |
|-------------------------------------|--------------------|---------------------|--|-------------------|
| | | | Range of HGs | Average No of HGs |
| <u>I. Life forms</u> | | | | |
| i. Climbers | 13(10.92) | 276(07.80) | 3 to 10 | 7.38 |
| ii. Creepers | 3(02.52) | 86(02.43) | 6-13 | 9.00 |
| iii. Herbs | 43(36.13) | 1235(34.89) | 5-18 | 9.16 |
| iv. Shrubs | 26(21.85) | 796(22.49) | 5-17 | 9.61 |
| v. Trees | 34(28.57) | 1147(32.40) | 6-22 | 10.32 |
| <u>II. Habitat</u> | | | | |
| i. Earthen plants | 92(77.31) | 2759(77.94) | 3-22 | 9.35 |
| ii. Potted plants | 26(21.85) | 756(21.36) | 6-18 | 9.65 |
| iii. Earthen/Potted plants | 1(00.84) | 25(00.71) | 7 | 7.00 |
| <u>III. Nature of Growth</u> | | | | |
| i. Cultivated | 89(74.79) | 2771(78.28) | 3-22 | 9.64 |
| ii. Wild | 2(01.68) | 66(01.86) | 7-9 | 8.00 |
| iii. Cultivated/Wild | 28(23.53) | 703(19.86) | 5-16 | 8.71 |
| <u>IV. Useful part</u> | | | | |
| i. Flower | 29(24.37) | 795(22.46) | 3-15 | 8.86 |
| ii. Fruit | 25(21.00) | 674(19.04) | 5-18 | 8.96 |
| iii. Leaf | 14(11.76) | 394(11.13) | 5-16 | 9.29 |
| iv. Seed | 2(01.68) | 41(01.16) | 7-10 | 8.50 |
| v. Stem | 1(00.84) | 45(01.27) | 14 | 14.00 |
| vi. Wood | 1(00.84) | 39(01.10) | 16 | 16.00 |
| vii. Whole plant | 47(39.50) | 1552(43.84) | 6-22 | 9.79 |
| <u>V. Uses</u> | | | | |
| i. Edible | 16(13.45) | 419(11.84) | 5-14 | 8.50 |
| ii. Fuel | 2(01.68) | 67(01.89) | 7-16 | 11.50 |
| ii. Medicine | 15(12.61) | 414(11.69) | 6-13 | 8.80 |
| iii. Multipurpose | 30(25.21) | 1057(29.86) | 6-22 | 10.23 |
| iv. Ornamental | 38(31.93) | 1067(30.14) | 3-16 | 9.00 |
| v. Vegetable | 18(15.13) | 516(14.58) | 5-18 | 9.89 |
| Total | 119(100.00) | 3540(100.00) | 3-22 | 9.39 |

In case of useful parts of plants, whole plant utilization category contain 47 species (39.50%) and is followed by flowers (29 species; 24.37%), fruits (25 species; 21%) and leaf (14 species; 11.76%). The total

number of individuals of HG plants higher (1552nos; 43.84%) in whole plant use category as compared to others. The number of HG s in which the whole plant used species present was ranged from 6 to 22 HG s with

an average of 9.79 HGs. Next to this, the leaf part used plants present in HG s range from 5 to 16 HGs with an average of 9.29 HGs; the flower used plant found in HGs ranges from 3 to 15 (average 8.86) HGs; the fruit used plant in HGs ranges from 5 to 18 (average 8.96) HGs; and the seed used plant present in HG ranges from 7 to 10 (average 8.5) HGs. But , the stem and woody part used plants found only 14 and 16 HGs, respectively (Table 6).

Under use category, plants used ornamentally showed a maximum of 38 species (31.93%) and is

followed by multipurposly used plants (30 species; 25.21%), vegetable plants (18 species; 15.13%), edible plants (16 species; 13.45%), medicinal plants (15 species; 12.61%) and the fuel plant (1 species;1.68%). A maximum number of 1067(30.14%) individuals was noted in ornamental category and is followed by 1057 (29.86%) number of plants in multipurpose use of plants. Fuel purpose plants found in 7 to 16 HGs with an average of 11.50 HGs and is followed by multipurposly used plants noted in 6-22 HGs with an average of 10.23 HGs (Table 6).

Table -7: Diversity indices estimated for homegarden species based on homegarden species distribution in Swamithoppe village of Kanyakumari District, Tamil Nadu.

| Plant distribution categories | Diversity indices | | | |
|------------------------------------|-------------------|--------------|----------------|---------------|
| | H' | E | λ | R |
| <u>I. Life forms</u> | | | | |
| i. Climbers | 0.393 | 0.153 | 0.00053 | 2.135 |
| ii. Creepers | 0.116 | 0.106 | 0.00021 | 0.449 |
| iii. Herbs | 1.606 | 0.427 | 0.00332 | 5.900 |
| iv. Shrubs | 1.054 | 0.324 | 0.00218 | 3.743 |
| v. Trees | 1.457 | 0.413 | 0.00466 | 4.684 |
| <u>II. Habitat</u> | | | | |
| i. Earthen | 3.582 | 0.792 | 0.00884 | 11.486 |
| ii. Earthen/Potted | 1.010 | 0.310 | 0.00201 | 3.772 |
| iii. Potted | 0.035 | 0.035 | 0.00005 | 0.000 |
| <u>III. Cultivated/Wild</u> | | | | |
| i. Cultivated | 3.616 | 0.806 | 0.00911 | 11.101 |
| ii. Cultivated/Wild | 0.087 | 0.126 | 0.00018 | 0.239 |
| iii. Wild | 0.924 | 0.277 | 0.00161 | 4.119 |
| <u>IV. Useful part</u> | | | | |
| i. Flower | 1.029 | 0.306 | 0.00200 | 4.193 |
| ii. Fruit | 0.910 | 0.283 | 0.00177 | 3.685 |
| iii. Leaf | 0.534 | 0.202 | 0.00094 | 2.175 |
| iv. Seed | 0.059 | 0.085 | 0.00008 | 0.269 |
| v. Stem | 0.055 | 0.055 | 0.00016 | 0.000 |
| vi. Wood | 0.050 | 0.050 | 0.00012 | 0.000 |
| vii. Whole plant | 1.990 | 0.517 | 0.00583 | 6.261 |
| <u>V. Uses</u> | | | | |
| i. Edible | 0.573 | 0.207 | 0.00100 | 2.484 |
| ii. Fuel | 0.088 | 0.127 | 0.00018 | 0.238 |
| iii. Medicine | 0.564 | 0.208 | 0.00097 | 2.323 |
| iv. Multipurpose | 1.326 | 0.390 | 0.00454 | 4.165 |
| v. Ornamental | 1.391 | 0.382 | 0.00273 | 5.306 |
| vi. Vegetable | 0.685 | 0.237 | 0.00148 | 2.722 |
| Total | 4.627 | 0.968 | 0.01090 | 14.440 |

H' -Shannon-Weiner's Diversity Index; E-Pielou's Index of Species Evenness; λ-Simpson's Diversity Index; R-Margalef's Index of Species Richness;

Diversity assessment of HG plants

Diversity indices estimated for HG plants surveyed in the study area, Swamithoppe village based on various distribution categories (such as lifeforms, habitat, growth condition, useful part and uses). In life form categories, herbs shows higher Shannon diversity index (H' =1.606), species evenness index (E =0.427), species richness index (R =5.8), while trees were more Simpson's diversity (λ= 0.00466), than other life forms.

Earthen plants shows maximum diversity (H' =3.582), species evenness (E =0.792), species richness (R =11.486) and more Simpson's diversity index (λ = 0.00884) as compared to potted and earthen/potted category. In Cultivated/Wild plants category, cultivated plants recorded maximum diversity (H' =3.616), species evenness (E =0.806), species richness (R =11.01) and Simpson's diversity index (λ=0.00911) as compared to cultivated/wild and wild plants categories (Table 7).

Among the plant parts used category, wholeplant utilized category shows maximum diversity ($H' = 1.99$), species evenness ($E = 0.517$), species richness ($R = 6.261$) while the Simpson's diversity index ($\lambda = 0.0058$) was higher that indicate less dominance as compared to other categories. Based on uses category of HG plants, ornamental plants show maximum diversity ($H' = 1.391$) and species richness ($R = 5.306$) while multipurposely used plants show maximum species evenness ($E = 0.390$) and Simpson's diversity index ($\lambda = 0.00454$) as compared to other plant categories (Table 7).

In the present work, it was observed in the study area that the HGs were generally maintained by house wives and not the household heads. It was also noted that the different responsibilities of male and female household members revealed different works. Females mostly maintain vegetables, spices, medicinal plants and ornamentals, whereas, males were responsible for shrubs and tree species. Thus, in HGs dominated by herbal plants females did most of the works, but in trees and shrubs dominated HGs women contributed only little works. Similar reports were also made by many workers [24-26]. In addition, the females did most of the HG works like hoeing, planting, weeding, fertilizing, and harvesting, whereas males carry out works like watering, pruning, land preparation, etc., as reported in several studies [27-29]. In some regions, HGs are managed mainly by females, i.e., in Thailand [30]; in Nepal [31]; and in Bangladesh [32-34].

In India, Dash and Mishra [35] reported that male dominating in HG works. In general, in the HGs of the study area, Swamithoppe village, are socio-economically sustainable with regards to labour input. The work input in HGs as compared to agricultural work is rather small, not very heavy, and having no heavy labour requirements. Instead, home gardening was done continuously year-round and in flexible manner as reported in other studies [36-38].

The HGs of Swamithoppe village have high species richness, with 119 species recorded from 121 HGs of 14043.59m² total areas with 108 genus belongs to 58 families containing 3540 individuals. Earlier reports indicated that in Arumanalloor rural village 83species was recorded from 66 HGs of 12,080.30m² total areas with 76 genera belongs to 43 families containing 2227 individuals [39]. The number of species can be related to the size of the HGs surveyed. It may be influenced by a number of factors such as socio-economic status of households, market integration, landholding size, etc., [40]. Homegarden exhibits complex structure, both vertically and horizontally. Vertical stratification is a common structure among homegardens throughout the tropics [41, 42]. The vertical structure composed of 3 to 4 layers based on the height and plant types [1]. In this

survey, it was noted that the HGs were mostly consisting two layered (69.42%), either with herbs and shrubs or shrubs and trees or herbs and tress. But, it was noted in the previous report that the rural HGs of Arumanalloor village shows three to four layered [43] or more layered structure [1, 44]. The wide range of species found in HGs adds their ecological efficiency in terms of use of physical and chemical resources [45, 46]. In this study, the pattern of species distribution in HGs was showed in table 11 & 12. In the study area, the herbs were dominant (36.13%) over other life forms with 77.31% of earthen plant, 74.79% of cultivated plants, 39.50% whole plant utilization species and 31.93% of ornamental plants. Similar observation was also made in Arumanalloor HGs studies by Neelamegam *et al.* [39]. It was also noted that there is no specific pattern of planting in the HGs of study area.

The structure of the homegardens may be determined by the species diversity of the plants present in each HG [47]. The numbers of local plants present in the homegardens provide an obvious starting point in determining the amount of diversity [48]. Eichenberg *et al.* [47] estimated the Shannon-Weiner diversity index value as 1.66 for the older urban HGs in Rio Claro which included 86 species; Brito *et al.* [49] reported a high diversity index of 2.22 in Aripuana; Neelamegam *et al.* [43] noted a diversity index of 3.977 in rural homegardens of Arumanalloor village with 83 species; and the mean SWI vary widely in tropical HGs and are reported to ranges from 0.93 to 3.00 [41] and from 0.69 to 4.01 [50]. In the present study, the overall Shannon-Weiner Index of plant diversity was estimated as $H' = 4.627$, the species evenness as $E = 0.968$, the species richness as $R = 14.44$ and the Simpson's Simpson's diversity index as $\lambda = 0.0109$. Neelamegam *et al.* [43] also reported species evenness as $E = 0.9$, species richness as $R = 10.637$, and Simpson's dominance as $\lambda = 0.022$ in the HGs of rural Arumanalloor village. Similar observations were also made by Sunwar [51] and Abiskar Subedi *et al.* [52]. This may be due to diverse agro-geographical conditions in rural area creating different micro-environments suitable for diverse species to maintain in conditions and limited options available for the households to grow different HG species as suggested by Abishar Subedi *et al.* [52]. Tynsong and Tiwari [50] estimated evenness index and dominance index in five village HGs, and the evenness index ranged from 0.56 to 1.15 and the dominance index was ranged from 0.06 to 1.15. Kabir and Web [53] reported strong relationship between homegarden sizes with species richness in Bangladesh HGs.

According to Saikia *et al.* [54] high diversity and low concentration of dominance in different HG categories may be due to variations in anthropogenic pressure in different HGs. With increase in household size, more varieties in species composition were also reported by Das and Das [40] in Barak valley, Assam. This suggests that the households maintain a diverse

group of plants to fulfill their regular needs regardless of the HG size. Diversity is selected according to the requirements of the families and the homegarden often contain a large number of individuals for certain species that are commonly utilized by the households. Species found in homegardens from the study area are used for primary and secondary needs of the household.

CONCLUSION

Homegarden are generally regarded as a very complex species rich agro-ecosystem managed in a sustainable manner over decade or even countries. The major purposes of homegardens are sustainable production, proper utilization and income generation, if possible, particularly in rural areas, in addition to fulfill the important ecological, social and cultural function. Plant diversity is considered as a basic for homegarden productivity and sustainability, which is dynamic with time. Both species composition and plant diversity are largely influenced by a combination of various socio-economic and ecological factors. In this study, socio-economic condition of the households, structure and composition, distribution pattern and diversity indices of Swamithoppe village homegarden plants have been presented and assessed. Based on the result, it can be concluded that the homegarden surveyed were managed by mostly family members with low labour investment; the homegarden suitable for *in situ* conservation of genetic resources, but plant diversity may highly dynamic over time and in future, may be threatened by modernization and commercialization; the homegarden surveyed having a high plant diversity, evenness and richness with low dominance; the species composition in homegardens mainly influenced by garden size and various households socio-economic factors; and the urban vegetation also has significance in removing atmospheric pollution.

ACKNOWLEDGEMENT

The authors thank to the Management Authorities, the Principal, S.T Hindu College, and the HOD, Department of Botany and Research Centre, S.T. Hindu College, Nagercoil, Kanyakumari District, India for providing necessary facilities and encouragement.

REFERENCES

1. Fernandes ECM, Nair PKR; An evaluation of the structure and function of tropical homegardens. *Agricultural Systems*, 1986; 21: 279-310.
2. Budowski G; Homegardens in Tropical America: a review. In: Landauer K, Brazil M; (eds), tropical Homegardens. United Nation's University Press, Tokyo, pp. 3-8, 1990.
3. Smith NJH; Home gardens as a spring board for Agroforestry Development in Amazonia. *International Tree Crops Journal*, 1996; 9: 11-30
4. Christanty L; Home gardens in tropical Asia, with special reference to Indonesia. In: Landauer K, Brazil M; (eds.). *Tropical Home Gardens*. The United Nations University, Tokyo, Japan, p. 9-20, 1990.
5. Okeke AI, Udofia SI; Income generation from homegardens of Akwalbom State, Nigeria. *Food production in a Developing Economy*. Int Edition, Afro-Euro Centre for Development studies, Granada, Spain, pp.377-383, 2009.
6. Soemarwoto O; Homegardens: A traditional agroforestry system with a promising future. In: Stepler, H.A.; Nair, P.K.R. (eds.) : *Agroforestry: A Decade of Development*. International Council for Research and Agroforestry (ICRAF), Nairobi, Kenya, p. 157-170, 1987.
7. Padoch C, De Jong W; The house gardens of Santa Rosa: Diversity and variability in an Amazonian agriculture system. *Economic Botany*, 1991; 45 (2): 166-175.
8. Peyre AA, Guodal A, Wiersum KF, Bongers F. Dynamics of homegarden structure and function in Kerala, India. *Agroforestry Systems*, 2006; 66: 101-115.
9. Perrault-Archambault M, Coomes OT; Distribution of agrobiodiversity in homegardens along the Corrientes river, Peruvian Amazon. *Economic Botany*, 2008; 62(2): 109-126.
10. Gajaseni J, Gajaseni N; Ecological rationalities of the traditional homegarden system in the Chao Phraya Basin, Thailand. *Agroforestry Systems*, 1999; 46: 3-23.
11. De Clerck, F.A.J.; Negreros-Castillo, P. (2000). Plant species of traditional Mayan homegardens of Mexico as analogs for multistrata agroforests. *Agroforestry Systems* 48: 303-317.
12. Lok R; A better understanding of traditional homegardens though the use of locally defined management zones. *Indigenous Knowledge and Development Monitor*, 2001; 9(2): 14-19.
13. Mendez VE, Lok R, Somarriba E; Inter disciplinary Analysis of Homegardens in Nicaragua: microzonation, plant use and socio-economic importance. *Agroforestry Systems*, 2001; 25(1): 31-58.
14. Abebe T, Wiersum KF, Bongers F; Spatial and temporal variation in crop diversity. *Agroforestry Systems*, 2010; 78: 309-322.
15. Gamble JS; *Flora of Presidency of Madras*, 3 Volumes. Allered and Son Limited, London, 1928.
16. Gamble JS, Fischer CEC; *Flora of Presidency of Madras*. Vol 1-111 (Repr. ed.). Botanical Survey of India, Calcutta, 1957.
17. Shannon CF, Wiener W; *The Mathematical Theory of communication*. University of Illinois Press, Urbana, p. 117, 1963.

18. Magurran AE; Ecological Diversity and Measurement. Croom Helm, London, UK, pp. 179, 1988.
19. Krebs CJ; Ecological Methodology. Benjamin/Cummings, Menlo Park, USA, pp. 620, 1999.
20. Pielou EC; Species diversity and pattern diversity in the study of ecological succession. J. Theo. Biol, 1996; 10: 370-383.
21. Nilima S, Sadika S, Vidyand N; Diversity of soil fungi in a tropical deciduous forest in Mudumalai. Southern India. Curr. Sci., 2007; 93: 67-669.
22. Margalef T; Temporal succession and spatial heterogeneity in Phytoplankton. In: Buzatti Traverso (ED.) Perspective in Marine Biology. University of California Press, Berkeley, 323-347, 1958.
23. Ludwig JA, Reynolds JF; Statistical ecology a primer on method and computing. Wiley-Interscience, New York, USA, pp. 337, 1998.
24. Trinh LN, Watson JW, Hue NN, De NN, Mish NV, Chu P, Sthapit BR, Eyzaguirre PB; Agrobiodiversity conservation and development in Vietnamese homegardens. Agriculture, Ecosystems and Environment, 2003; 97: 317-344.
25. Azurdia C, Leiva JM; Home-garden biodiversity in two contrasting regions of Guatemala. In: Eyzaguirre PB, Linares OF; (eds.). Home Gardens and Agrobiodiversity. Smithsonian Books, Washington, USA, p. 168-184, 2004.
26. Howard PL; Gender and social dynamics in Swidden and homegardens in Latin America. In: Kumar BM, Nair PKR; (eds.). Tropical Homegardens: A Time-Tested Example of Sustainable Agroforestry. Advances in Agroforestry, Vol. 3, Springer Science, Dordrecht, The Netherlands, pp. 159-182, 2006.
27. Rugalema GH, Okting'ATI A, Johnsen FH; Thehomegarden agroforestry system of Bukoba district, North –Western Tanzania. 1. Farming system analysis. Agroforestry System, 1994; 26: 53-64.
28. Tchatat M, Puig H, Fabre A; Genèseetorganisation des jardins de case des zones forestièreshumides du Cameroun. Revue d'Écologie: La Terre et la Vie, 1996; 51: 197-221.
29. Bennett-Lartey SO, Ayernor GS, Markwei CM, Asante IK, Abbiw DK, Boateng SK, Anchirinah VM, Ekpe P; Aspects of home-garden cultivation in Ghana: Regional differences in ecology and society. In: Eyzaguirre PB, Linares OF; (eds.). Home Gardens and Agrobiodiversity. Smithsonian Books, Washington, USA, p. 148-167, 2004.
30. Moreno-Black G, Somnasang P, Thamathawan S; Cultivating continuity and creating change: Woman's home garden practices in North-eastern Thailand. Agriculture and Human Values, 1996; 13: 3-11.
31. Shrestha P, Gautam R, Rana RB, Sthapit B; Managing diversity in various ecosystem homegardens of epal. In Eyzaguirre, P.B., Linares, O.F., (eds.). homegardens and Agrobiodiversity. Smithsonian book, Washinington, USA, p. 95-122, 2004.
32. Oakley E; Home gardens: A cultural responsibility. Leisa Magazine, 2004; 20: 22-23.
33. Ali AMS; Homegardens in smallholder farming systems: Examples from Bangladesh. *Human Ecology*, 2005; 33: 245-270.
34. Oakley E, Mornsen JH; Women and seed management: A study of two villages in Bangladesh. Singapore Journal of Tropical Geography, 2007; 28: 90-106.
35. Dash SS, Misra MK; Studies on hill agro-ecosystems of three tribal villages on the Eastern Ghats of Orissa, India. Agriculture, Ecosystems and Environment, 2001; 86: 287-302.
36. Kimber C; Dooryard gardens of Martinique. Yearbook of the Association Pacific Coast Geographers, 1966; 28: 97-118.
37. Christanty L, Abdoellah OS, Marten GG, Iskandar J; Traditional agroforestry in West java: the pekarangan (homegardens) and kebun –talun (annual-perennial rotation) cropping system. In: Marten GG; (ed.).Traditional Agriculture in southeastern Asia, Boulder, Colorado, USA, 1986.
38. Kyoslef H; Homegardens of Javanese transmigrants in Seberida sub district: description, agroecological constraints and evaluation of potential solutions to declining productivity. In: Sandbukt O, Wiriadinata H; (eds.). Rain Forest and Resource Management. Norindra Seminar, Jakarta, 1994. Indonesian Institute of Sciences (LIPI), Indonesia, p. 127-136, 1994.
39. Neelamegam R, Mathevan Pillai V, Mary Anishal Priyanka A, Roselin S; Status and composition of home garden plants in rural and urban areas in Kanyakumari District, Tamil Nadu, India. Scholars Academic Journal of Biosciences (SAJB), 2015a; 3(8): 656-667.
40. Das T, Das AK; Inventorying plant biodiversity in homegardens: A case study in Barak valley, Assam, North East India. Current Science, 2005; 89: 155-163.
41. Karyono; Home Gardens in Java. Their Structure and Function. In: Landauer K, Brazil M; (eds). Tropical Home Gardens. The United Nations University, Tokyo, Japan, p.138-146, 1990.

42. Gillespie AR, Knudson DM, Geilfus F; The structure of four home gardens in the Petén, Guatemala. *Agroforestry Systems*, 1993; 24: 157-170.
43. Neelamegam R, Roselin S, Mary Anishal Priyanka A, Mathevan Pillai V; Diversity indices of home garden plants in rural and urban areas in Kanyakumari District, Tamil Nadu, India. *Scholars Academic Journal of Biosciences*, 2015; 3(9): 752-761.
44. Zimic I, Saikia P, Khan MI; Comparative study on homegardens of Assam and Arunachal Pradesh in terms of species diversity and plant utilization pattern. *Research Journal of Agricultural Sciences*, 2012; 3(3): 611-618.
45. Wiersum KF; Tree gardening and taungyc in Java: Example of agroforestry techniques in the humid tropics. *Agroforestry system*, 1982; 1: 53-70.
46. Blanckaert I, Swennen RL, Paredes Flores M, Rosas Lopez R, Lira Saade R. Floristic composition, plant uses and management practices in homegardens of San Rafael Coxcatan, valley of Tehuacan-Cuicatlan, Mexico. *Journal of Arid Environment*, 2004; 57: 39-62.
47. Eichemberg MT, Maria Christina, de MeoAmoroza, e Leila Cunha de Moura; Species composition and plant use in old urban home gardens in Ris Claro, southeast of Brazil. *Acta bot Bras*, 2009; 23(4): 1057-1075.
48. Hodgkin T; Home gardens and maintenance of genetic diversity. In: Watson JW, Eyzaguirre PB, (eds.) *Proceedings of the Second International Home Gardens Workshops: Contribution of home gardens to In situ Conservation of plant genetic resources*, Witzenhausen, Federal Republic of Germany, International Plant Genetic Resources, Rome, 2002.
49. Brito MA; In. Eichemberg MT, Maria Christina de MeoAmoroza, e Leila Cunha de Moura; Species composition and plant use in old urban home gardens in Ris Claro, southeast of Brazil. *Acta bot Bras*, 2009; 23(4): 1057-1075.
50. Tynsong H, Tiwari BK; Plant diversity in the homegardens and their significance in the livelihoods of War Khasi community of Meghalaya, Northeast India. *Journal of Biodiversity*, 2010; 1(1): 1-11.
51. Sunwar S; Does Shannon-Weaver Index explains the species diversity in home gardens? In: *Home Gardens in Nepal: Proceeding of a workshop* (eds.) 2006, Hautam R, Sthapit BR, Shrestha, PK, 6-7 August 2004; Pokhara, Nepal, 2004; 324: 66-71.
52. AbishkarSubedi, RojeeSuwa, ReshamGautam, SharmilaSunwar, PratapShrestha; Status and composition of plant genetic diversity in Nepalese Home Gardens. In: *Home Gardens in Nepal: Proceeding of a workshop* (eds.) 2006, Hautam R, Sthapit BR, Shrestha, PK, Pokhara, Nepal, 2004.
53. Kabir ME, Web EL; Household and homegarden characteristics in South Western Bangladesh. *Agroforestry Systems*, 2009; 75: 129-145.
54. Saikia P, Choudhury BI, Khan M; Floristic composition and plant utilization pattern in home gardens of Upper Assam, India. *Tropical Ecology*, 2012; 53(1): 105-118.

Annexure Table

Table -I: List of species recorded in the homegardens of Swamithoppe village, Kanyakumari District, Tamil Nadu, India.

| Sl. No. | Name of the species | Common name | Family | LF | C/W | E/P | UsPa | Uses |
|---------|---|--------------------------|----------------|----|-----|-----|------|------|
| 1 | <i>Abelmoschus esculentus</i> (L.) Moench | Vendai | Malvaceae | H | C | E/P | Fr | Veg |
| 2 | <i>Acalypha hispida</i> Linn. | Fox tail | Euphorbiaceae | H | C | E | Le | Med |
| 3 | <i>Acalypha indica</i> L. | Kuppai meni | Euphorbiaceae | H | W | E | Le | Med |
| 4 | <i>Achras sapota</i> (L.) P.Royan | Sapota | Sapotaceae | T | C | E | Fr | Edi |
| 5 | <i>Allium cepa</i> L. | Ullie | Liliaceae | H | C | E/P | WP | Veg |
| 6 | <i>Aloe vera</i> (L.) Burm F. | Kathazhi | Liliaceae | H | W | E | WP | MP |
| 7 | <i>Adenanthera ficoidea</i> Linn. | Ani Kundamani | Adenanthereae | H | W | E | Le | Med |
| 8 | <i>Amaranthus blitum</i> Linn. | Thandu keerai | Amaranthaceae | H | C | E | WP | Veg |
| 9 | <i>Amaranthus tricolor</i> L. | Spinach | Amaranthaceae | T | C | E | Le | Veg |
| 10 | <i>Amorphophallus paenoiifolus</i> Bl. | Karu-naik-kishangu | Araceae | H | C | E | WP | MP |
| 11 | <i>Anacardium occidentale</i> Linn. | Kollammaram | Anacardiaceae | S | W | E | WP | MP |
| 12 | <i>Ananas cosmosus</i> (L.) Merr. | Annachi | Bromeliaceae | H | C | E | Fr | Edi |
| 13 | <i>Annana squamosa</i> Linn. | Sethapazhlam | Annonaceae | T | C | E | Fr | Edi |
| 14 | <i>Arachis hypogaea</i> L. | Verkadalai | Papilionaceae | H | C | E | Se | MP |
| 15 | <i>Areca catechu</i> L. | Pakkumaram | Arecaceae | T | C | E | Se | MP |
| 16 | <i>Artocarpus heterophyllus</i> (L.f.) Bhandari | Palamaram | Moraceae | T | C | E | WP | MP |
| 17 | <i>Auracaria sps.</i> | Monkey tail | Auracareace | T | C | E/P | WP | Om |
| 18 | <i>Averrhoa bilimbi</i> Linn. | Pulichankai | Geraniaceae | T | C | E | Fr | Edi |
| 19 | <i>Azadirachta indica</i> A.Juss. | Veppamaram | Meliaceae | T | C | E | WP | MP |
| 20 | <i>Bambusa arundinacea</i> (Retz.) Roxb. | Moongil | Gramineae | T | C | E | WP | MP |
| 21 | <i>Bauhinia acuminata</i> Linn. | Anthimantharai | Leguminosae | H | C | E | F1 | Om |
| 22 | <i>Bougainvillea spectabilis</i> L. | Nyctaginaceae | Bignoniaceae | S | C | E | F1 | Om |
| 23 | <i>Borassus flabellifer</i> Linn. | Panamaram | Palmaceae | T | W | E | WP | MP |
| 24 | <i>Calophyllum inophyllum</i> Linn. | Punnaimaram | Guttiferae | T | W | E | WP | MP |
| 25 | <i>Cassia auriculata</i> L. | Avaram | Caesalpinaceae | Cl | C | E | WP | Om |
| 26 | <i>Canna indica</i> L. | Kalvazhlai | Musaceae | T | C | E | WP | MP |
| 27 | <i>Calotropis gigantea</i> Linn. | Erukku | Asclepiadaceae | S | W | E | WP | Med |
| 28 | <i>Carica papaya</i> L. | Pappali | Caricaceae | T | C | E | WP | MP |
| 29 | <i>Catharanthus roseus</i> (L.) G. Don. | Nithiya kalyani | Apocynaceae | S | C/W | E | F1 | Om |
| 30 | <i>Celosia argentea var. cristata</i> | Kozhlivada | Amaranthaceae | H | C | P | F1 | Om |
| 31 | <i>Centella asiatica</i> Linn. | Vallarai | Umbelliferae | Cr | C | E | WP | MP |
| 32 | <i>Chamabaina cuspidata</i> Wt. | Insulin plant | Urticaceae | H | C | E | WP | MP |
| 33 | <i>Cissus quadrangularis</i> | Pirandai | Vitaceae | H | C/W | E | F1 | Om |
| 34 | <i>Citrus aurantium</i> Linn. | Naarthalai(Sour) | Rutaceae | T | C | E | Fr | Edi |
| 35 | <i>Clitoria ternatea</i> Linn. | Chankkupoospa m | Leguminosae | Cl | W | E | F1 | Om |
| 36 | <i>Cocos nucifera</i> Linn. | Thennaimaram | Arecaceae | T | C | E | WP | MP |
| 37 | <i>Crossandra infundibuliformis</i> (L.) Ness. | Kanagamaram | Acanthaceae | S | C | E/P | F1 | Om |
| 38 | <i>Crotalaria verrucosa</i> Linn. | Kilukiluppai | Leguminosae | H | W | E | F1 | Om |
| 39 | <i>Croton sparsiflorus</i> Linn. | Mannanaiveratti | Euphorbiaceae | H | W | E | F1 | Om |
| 40 | <i>Cucumis sativa</i> L. | Vellarikai | Cucurbitaceae | Cl | C | E | Fr | Edi |
| 41 | <i>Cucurbita pepo</i> (L.) Dumort. | Poosani | Cucurbitaceae | Cl | C | E | Fr | Edi |
| 42 | <i>Curcuma longa</i> L. | Manjal | Zingiberaceae | H | C | E/P | WP | MP |
| 43 | <i>Cyamopsis tetragonoloba</i> Taub. | Kothavarai | Leguminosae | Cl | C | E | WP | Veg |
| 44 | <i>Daemia extensa</i> Linn. | Veliparuthi | Asclepiadaceae | Cr | W | E | WP | MP |
| 45 | <i>Datura metal</i> Linn. | Oomatthai | Solanaceae | H | W | E | F1 | Om |
| 46 | <i>Dolichos lablab</i> L. | Avarakai | Papilionaceae | Cl | C | E | Fr | Veg |
| 47 | <i>Dracaena sps.</i> | Happy plant/ Dragan tree | Asparagaceae | S | C | E/P | Le | Om |
| 48 | <i>Duranta pluemeri</i> Jacq. | Goldan-dew-drop | Verbenaceae | S | C | E | F1 | Om |
| 49 | <i>Eclipta alba</i> L. | Karisalankanni | Asteraceae | S | C | E | F1 | Om |
| 50 | <i>Epipremnum aureum</i> (L.) Engl. | Money plant | Araceae | Cl | C | E/P | Le | Om |
| 51 | <i>Euphorbia heterophylla</i> L. | Ellaimelraja | Euphorbiaceae | S | C | E | WP | Med |
| 52 | <i>Euphorbia hirta</i> L. | Ammanpacharisi | Euphorbiaceae | H | W | E | WP | Med |
| 53 | <i>Euphorbia nilli</i> Linn. | Crown of thorns | Euphorbiaceae | H | C | E/P | WP | Med |
| 54 | <i>Ficus religiosa</i> Linn. | Arasamaram | Moraceae | T | C | E | WP | MP |
| 55 | <i>Furcraea gigantea</i> Vent. | Giant aloe | Dioscoreaceae | S | C | E/P | F1 | Om |
| 56 | <i>Gomphrena decumbens</i> L. | Vadamalli | Amaranthaceae | H | W | E/P | F1 | Om |
| 57 | <i>Hemidesmus indicus</i> L. | Nannari | Asclepiadaceae | H | C | E/P | Le | Med |
| 58 | <i>Hibiscus rosa sinensis</i> Linn. | Chembaruthi | Malvaceae | S | C | E | WP | MP |
| 59 | <i>Impatiens balsamina</i> L. | Balsam | Geraniaceae | H | C | E/P | F1 | Om |
| 60 | <i>Ipomoea quamodit</i> L. | Mayilmanikam | Convolvulaceae | Cl | C | E/P | WP | Om |

| | | | | | | | | |
|-----|--|--------------------------|----------------|----|---|-----|-------|-----|
| 61 | <i>Ixora coccinea</i> L. | Thitti | Rubiaceae | S | C | E | WP | Orn |
| 62 | <i>Ixora</i> sps. L. | Idlipoo | Rubiaceae | S | C | E/P | WP | Orn |
| 63 | <i>Jasminum auriculatum</i> Vahl. | Malligai | Oleaceae | H | C | E | Fl | Orn |
| 64 | <i>Jasminum multiflorum</i> sps. | Kattumalligai | Oleaceae | H | W | E | Fl | Orn |
| 65 | <i>Jatropha</i> Linn. | Kattamanakku | Euphorbiaceae | S | W | E | WP | Fue |
| 66 | <i>Lawsonia innermis</i> L. | Maruthani | Lythraceae | T | C | E | Le | Med |
| 67 | <i>Leucas aspera</i> Spreng. | Thumbai | Lamiaceae | H | W | E | WP | Med |
| 68 | <i>Lycopersicon esculentum</i> Mill. | Thakkali | Solanaceae | S | C | E | Fr | Veg |
| 69 | <i>Mangifera indica</i> L. | Mamaram | Anacardiaceae | T | W | E | Fr | Edi |
| 70 | <i>Mentha piperita</i> L. | Pudina | Lamiaceae | S | C | E | Le | Veg |
| 71 | <i>Mirabilis jalapa</i> Linn. | Anthimantharai | Nyctaginaceae | H | W | E/P | Fl | Orn |
| 72 | <i>Momordica charantia</i> L. | Pagarkai | Cucurbitaceae | Cl | C | E | Fr | Veg |
| 73 | <i>Morinda tinctoria</i> Roxb. | Manjanathi | Rubiaceae | T | W | E | Wo | Fue |
| 74 | <i>Moringa oleifera</i> Lamk. | Murungaimaram | Moringaceae | T | C | E | WP | MP |
| 75 | <i>Murraya koengii</i> (L) Spreng. | Karuveppilai | Rutaceae | T | C | E/P | Le | Veg |
| 76 | <i>Musa paradisiaca</i> Linn. | Vazhlaimaram | Musaceae | T | C | E | WP | MP |
| 77 | <i>Mussanda erythrophylla</i> Hutch. | Queen sirikit | Rubiaceae | S | C | E | Fl | Orn |
| 78 | <i>Nerium indicum</i> Mill. | Arali | Apocynaceae | S | C | E | Fl | Orn |
| 79 | <i>Ocimum sanctum</i> L. | Tulasi | Lamiaceae | H | C | E | WP | MP |
| 80 | <i>Oldenlandia umbellata</i> Linn. | Indian madar | Rubiaceae | H | C | E/P | Fl | Orn |
| 81 | <i>Pandanus amaryllifolium</i> Roxb. | Rambai | Pandanaceae | H | C | E/P | Le | Veg |
| 82 | <i>Phaseolus vulgaris</i> Linn. | Beans | Fabaceae | Cl | C | E | Fr | Veg |
| 83 | <i>Phoenix sylvestris</i> Roxb. | Perichchamaram | Arecaceae | T | W | E | Fr | Edi |
| 84 | <i>Phyllanthus emblica</i> L. | Nellikamaram | Euphorbiaceae | T | W | E | WP | MP |
| 85 | <i>Phyllanthus niruri</i> Linn. | Kezhlanelli | Euphorbiaceae | H | C | E | Le | Med |
| 86 | <i>Piper nigrum</i> L. | Kaduku | Piperaceae | Cl | C | E | Fr | MP |
| 87 | <i>Pithecellobium dulce</i> (Roxb) Benth. | Kodukapuli | Mimosaceae | T | W | E | Fr | Edi |
| 88 | <i>Plectranthus amboinicus</i> (L) He'r | Karpuravalli | Labiatae | H | C | E/P | Le | Med |
| 89 | <i>Plumpago zeylanica</i> Linn. | Sky flower | Plumbaginaceae | H | C | E | WP | MP |
| 90 | <i>Polyalthia longifolia</i> (Sonner.) Thw. | Nettulingam | Annonaceae | T | C | E | WP | Orn |
| 91 | <i>Prunus avium</i> (L) L. | Cherry | Rosaceae | T | C | E | Fruit | Edi |
| 92 | <i>Psidium guajava</i> (L). Bot. | Koiyamaram | Myrtaceae | T | C | E | Fr | Edi |
| 93 | <i>Punica granatum</i> Linn. | Mathulai | Punicaceae | S | C | E | Fr | Edi |
| 94 | <i>Rhoeo spathacea</i> sps. | Boat lilly | Commelinaceae | H | C | E/P | Fl | Orn |
| 95 | <i>Ricinus communis</i> Linn. | Ammanakku | Euphorbiaceae | S | W | E | WP | Med |
| 96 | <i>Rosa</i> sps. | Roja | Rosaceae | H | C | E/P | Fl | Orn |
| 97 | <i>Saccharum officinarum</i> Linn. | Karumpu | Gramineae | S | C | E/P | St | Edi |
| 98 | <i>Santalum album</i> L. | Santanamaram | Santalaceae | T | C | E | WP | MP |
| 99 | <i>Solanum capsicum</i> Linn. | Patchamillagai | Solanaceae | H | C | E/P | Fr | Veg |
| 100 | <i>Solanum melongena</i> Linn. | Kathirikai | Solanaceae | H | C | E | Fr | Veg |
| 101 | <i>Solanum torvum</i> Swarta. | Sundakai | Solanaceae | H | C | E | WP | Veg |
| 102 | <i>Solanum trilobatum</i> L. | Thuthuvalai | Solanaceae | Cl | C | E | Le | Veg |
| 103 | <i>Solanum nigrum</i> Linn. | Manathakalli | Solanaceae | H | C | E/P | Fr | Veg |
| 104 | <i>Stachytarpheta jamaicensis</i> Vahl. | Kattunairivi | Verbenaceae | H | W | E | WP | Med |
| 105 | <i>Syzygium cumini</i> (L) Skeels. | Navalmaram | Myrtaceae | T | W | E | WP | MP |
| 106 | <i>Tabernaemontana divaricata</i> R. Br. Ex. Roem. & Schult. | Nanthiya vattai | Apocynaceae | S | C | E | Fl | Orn |
| 107 | <i>Talinum portulacifolium</i> (Forssk.) Asch. Ex. Schweinf. | Galaghati | Portulacaceae | H | C | E/P | Fl | Orn |
| 108 | <i>Tamarandus indica</i> Linn. | Puliyamaram | Cesalpiniaceae | T | C | E | WP | MP |
| 109 | <i>Tecoma stans</i> | Manjapoo | Bignoniaceae | S | C | E | Fl | Orn |
| 110 | <i>Tecomaria capensis</i> | Cape honey Suckle | Bignoniaceae | S | C | E | Fl | Orn |
| 111 | <i>Tectona grandis</i> Linn. | Thekkumaram | Verbenaceae | T | C | E | WP | MP |
| 112 | <i>Tephrosia purpurea</i> (L)Pers. | Kollukainelai | Papilionaceae | H | W | E | Fl | Orn |
| 113 | <i>Terminalia catappa</i> Linn. | Badam | Combretaceae | S | C | E/P | Fr | Edi |
| 114 | <i>Thespesia populnea</i> L. Soland. | Poovarsamaram | Malvaceae | T | C | E | WP | MP |
| 115 | <i>Thuja occidentalis</i> L. | Arbor vitae/ White cedar | Cupressaceae | T | C | E | WP | Orn |
| 116 | <i>Trichosanthes anguina</i> L. | Pudalankai | Cucurbitaceae | Cl | C | E | Fr | Edi |
| 117 | <i>Vernonia cinerea</i> Linn. | Punarva | Asteraceae | H | W | E | Fl | Orn |
| 118 | <i>Wedalia trilobata</i> Jacq. | Pottralaikaiyyan | Asteraceae | Cr | C | E | WP | Med |
| 119 | <i>Zingiber officinale</i> Roscoe | Inji | Zingiberaceae | S | C | E | Fr | Veg |

LF-Life forms; T-Tree; H-Herb; S-Shrub; Cl-Climbers; Cr-Creepers; C-Cultivated; W-Wild; WP-Whole Plant; E-Earthern; P-Potted; Veg-Vegetable; Orn-Ornamental; MP-Multipurpose; Le-Leaf; Fr-Fruits; Fl-Flower; Med-Medicinal; Edi-Edible; Se-Seeds; St-Stem;