

Case Report

Space Occupying Lesion of the Liver: The Feline Connection

Cyriac Abby Philips¹, Chhagan Bihari Sharma², Apurva Pande¹, Juned Khan¹

¹The Liver Intensive Unit

²Department of Hepatology and Transplant Medicine, Institute of Liver and Biliary Sciences, New Delhi

*Corresponding author

Cyriac Abby Philips

Email: abbyphilips@gmail.com

Abstract: Eosinophilic granulomatous abscesses, the spectrum of visceral larva migrans are rare. Most reports are from the Far-East and India where the most commonly attributed cause is *Toxocara canis* or *cati*. We present here, the case of a middle aged woman with cats as pets who developed 3 to 4 weeks history of abdominal pain, fever and pruritus, initially managed conservatively and in whom, a simple but in depth history analysis led to the diagnosis of hepatic visceral larva migrans confirmed on fine needle aspiration cytology. We also provide a short algorithmic approach to differential diagnosis of liver SOLs in adults.

Keywords: visceral larva migrans, liver abscess, eosinophilia, *Toxocara*, fine needle aspiration cytology, fever of unknown origin.

INTRODUCTION

Visceral larva migrans occurs during the migratory second larva stage of nematodes in human beings. Dogs and cats are the most common primary hosts for the disease causing nematodes *Toxocara canis* and *Toxocara cati* respectively[1]. Uncommonly, *Capillaria hepatica*, *Ascaris suum*, and *Ancylostoma* species also cause similar hepatic disease[2]. Accidental ingestion of embryonated eggs present in the soil or the arrested second-stage larvae of nematodes from the primary host leads to infections in humans. Ingested eggs or larvae can evolve into migrating larvae, which are released into the small bowel, entering the portal venous system, and migrate to the liver, lungs, brain, heart, and eyes. Larva migrans in the liver present as eosinophilic abscesses or granulomas[3]. On ultrasonography, appearance is usually multiple rounded or oval hypoechoic lesions with conglomeration. Contrast computed tomography or magnetic resonance imaging reveals ill-defined enhancing fuzzy walls of liquefied conglomerating lesions which may be oval, rounded, or asymmetrical seen predominantly in the venous and equilibrium phases. A unique characteristic feature of these abscesses on plain MRI study hyperintense rim on T1-imaging and diffusion restriction on echo planar imaging[4].

CASE REPORT

A 48 year old housewife, presented with 2 weeks history of fever and right upper quadrant abdomen pain. She recovered from fever, severe pruritus and urticaria two weeks before after course of montelukast and cetirizine, 6 weeks prior to current presentation. Four months prior, she bought 4 cats from a local pet shop and was caring for them at home as a past time. Examination revealed mild pallor without organomegaly or skin changes. Ultrasound (USG) done elsewhere showed large heterogeneous lesion in the right lobe of liver. Blood investigations revealed anaemia (haemoglobin 9.8 g/L), total leukocyte count $13 \times 10^9/L$ with 24% eosinophils. Contrast MRI abdomen revealed multiple heterogeneous coalescing, peripherally enhancing well defined T1 iso-intense (Figure 1, Panel A) and T2 hyper-intense lesions with central non-enhancing areas and thick peripheral wall involving segments IVB and V (Figure 1, Panel B). Serum tumor markers were within normal limits. USG guided fine needle aspiration from the liver mass (Figure 1, Panel C) revealed large number of eosinophils (Panel C dashed arrow), in the background admixed with multiple Charcot-Layden crystals (Panel C, black arrow) leading to final diagnosis of visceral larva migrans (VLM) – eosinophilic liver abscess.




Fig 1: Contrast MRI abdomen showing multiple heterogeneous coalescing, peripherally enhancing well defined T1 iso-intense (Panel A) and T2 hyper-intense lesions with central non-enhancing areas and thick peripheral wall involving segments IVB and V (Panel B). USG guided fine needle aspiration from the liver mass showing large number of eosinophils (H and E stain, 20X Panel C dashed arrow), in the background admixed with multiple Charcot-Layden crystals (H and E stain, 20X, Panel C, black arrow)

DISCUSSION

VLM is caused by *Toxocara canis* (dog round worm) or *catis* (cat round worm) and even *Fasciola hepatica* or fungal infections such as *Basidio bolusranarum*. The 2nd stage larva of nematode when accidentally ingested by human host enter portal circulation through small bowel wall, infesting distant organs such as lung, liver and eyes[5]. Visceral larva migrans can present as eosinophilic liver infiltrates or as granulomatous abscesses. On CT, hepatic lesions present as small, ill defined, oval or elongated hypo dense nodules best seen on portal venous phase[6]. On MR imaging, the lesions can have variable intensities,

but usually hypo-intense or iso-intense on T1 and hyper-intense on T2 sequences (Panel B, white arrow). Peculiar findings include coalescent lesions (Panel A, dotted arrow) with peripheral rim enhancement in portal venous phase[7]. Differentials include hepatocholangiocarcinoma or infected liver metastases[8]. A short algorithm for approach to focal liver lesions is shown in Figure 2. When dealing with atypical liver abscess, systemic symptoms such as easy fatigability, weight loss, skin rash and associated constitutional symptoms with peripheral blood work showing eosinophils or atypical cells point toward specific, but rare aetiology.




Fig 2: Differential diagnosis algorithm of liquid and solid space occupying lesions in the liver

CONCLUSION

Hepatic visceral larva migrans refers to a condition characterized by granulomatous liver lesions containing eosinophils and inflammatory cells secondary to second stage larval migration of *Toxocara canis* or *catis*. Typical imaging features include small, ill-defined, oval or elongated, low-attenuating nodules with shaggy margins, non-spherical shape, and absent or insignificant rim enhancement on contrast-enhanced computed tomography. Early resolution of lesions is seen with prompt utilization of anti-helminthic treatment for 3 to 6 months.

REFERENCES

1. Lim JH. Toxocariasis of the liver: Visceral larva migrans. *Abdom Imaging* 2008; 33:151-6.
2. Laroia ST, Rastogi A, Sarin S. Case series of visceral larva migrans in the liver: CT and MRI findings. *Int J Case Rep Images* 2012; 3:7-12.

3. Chang S, Lim JH, Choi D, Park CK, Kwon NH, Cho SY, et al. Hepatic visceral larva migrans of *Toxocara canis*: CT and sonographic findings. *AJR Am J Roentgenol* 2006; 187:W622-9.
4. Ramachandran J, Chandramohan A, Gangadharan SK, Unnikrishnan LS, Priyambada L, Simon A. Visceral larva migrans presenting as multiple liver abscesses. *Trop Doct* 2013; 43:154-7.
5. Santiago Rivera L, Figueroa Rivera I, Toro DH, Gutierrez J, Acosta E; Eosinophilic Liver Infiltration. *ACG Case Reports Journal*. 2015; 3(1):63-65.
6. Hassan HA, Majid RA, Rashid NG, Nuradeen B.E, Abdulkarim Q.H, Hawramy T.A *et al.*; Eosinophilic granulomatous gastrointestinal and hepatic abscesses attributable to basidio bolomycosis and fasciolias: a simultaneous emergence in Iraqi Kurdistan. *BMC Infectious Diseases*. 2013; 13(1):1.

7. Mukund, A. Arora, Y. Patidar, Mangla V, Bihari C, Rastogi A *et al.*; "Eosinophilic abscesses: a new facet of hepatic visceral larva migrans," *Abdominal Imaging*, 2013; 38(4):774-777.
8. Laroia ST, Rastogi A, Bihari C, Bhadoria AS, Sarin SK; Hepatic visceral larva migrans, a resilient entity on imaging: Experience from a tertiary liver center. *Tropical Parasitology*. 2016; 6(1):56-68.