

Original Research Article

Insomnia & Serum Cortisol level in Major Depressive Disorder

Soma Gupta¹, Ranjana Mandal², Somsubhra Chattopadhyay³, Saswati Nath⁴, Harendra Nath Das⁵

¹Professor, Department of Biochemistry, N.R.S. Medical College, (formerly attached to College of Medicine & Sagore Dutta Hospital), Kolkata, West Bengal, India

²Associate Professor, Department of Physiology, College of Medicine & Sagore Dutta Hospital, Kolkata, West Bengal, India

³Assistant Professor, Department of Psychiatry, College of Medicine & Sagore Dutta Hospital, Kolkata, West Bengal, India

⁴Associate Professor, Department of Psychiatry, R.G. Kar Medical College, Kolkata, West Bengal, India

⁵Ex Professor & Head, Department of Biochemistry, R.G. Kar Medical College, Kolkata, West Bengal, India

***Corresponding author**

Prof. Soma Gupta

Email: guptasomarusa@yahoo.com

Abstract: A total of 207 subjects (101 cases of depression & 106 age & sex matched controls) were studied by measuring Athens Insomnia scale (AIS) and serum cortisol. Among 101 cases of depression, 19 cases were of mild grade, 37 were of moderate and 45 were suffering from severe depression. It has been found that 91 out of 101 MDD cases had AIS score ≥ 6 . Thus the prevalence of insomnia was calculated to be 90.1%. Mean concentration of cortisol was found to be 10.2 ± 5.5 and 6.8 ± 1.6 $\mu\text{g/dl}$ and AIS score was found to be 11.5 ± 4.3 & 3.8 ± 1.6 in cases & controls respectively. Both the parameters were found to be statistically significant. The cases were further classified according to severity. The correlation of cortisol and AIS score with severity of depression was assessed by Spearman rank correlation test. AIS score was found to be significantly correlated with depression showing a distinct relationship between insomnia and depression. Serum cortisol level however, was not found to correlate with the severity of depression.

Keywords: Major Depressive Disorder (MDD), Insomnia, serum cortisol, Athens Insomnia Scale (AIS)

INTRODUCTION

Sleep disorder in form of Insomnia is one of the most common features of depression and the intimate nexus between sleep and depressive symptoms have been studied elaborately. Epidemiological studies show that symptoms of sleep disturbances precede an episode of depression in 40% of cases. It has been observed that people suffering from depression with prominent insomnia are likely to have a poorer response to treatment. Insomnia is also found to be predictive of a greater risk of suicidal behavior [1].

A distinct circadian rhythm is observed in humans, which determine multiple biological and behavioral processes performed by them. One such biological process is sleep which follows a diurnal pattern. This diurnal pattern plays a key role to adapt a person to environmental circumstances. Adaptation to extrinsic and intrinsic forces is a survival necessity for all living organisms. The hypothalamic-pituitary-adrenal (HPA) axis is an adaptive system with the purpose of maintaining a dynamic equilibrium or

homeostasis in a constantly changing environment. Sleep is regulated by the HPA axis in multiple ways, and a growing body of research suggests reciprocal associations between sleep and the activity of the HPA axis [2].

It has been observed that a similar diurnal pattern exists in the stress response, which may be of particular relevance to stress related disorders such as depression. This observation is further enforced as almost 40–60% of patients suffering from depression have been found to have a disturbance in the hypothalamic-pituitary-adrenal (HPA) axis, which manifest as changes in patterns of diurnal cortisol release [3].

The hypothalamus first manufactures corticotrophic-releasing hormone (CRH), which stimulates pituitary gland to release adreno cortico trophic hormone (ACTH). This hormone then makes the adrenal glands to secrete cortisol in the blood. In normal condition, the hypothalamus monitors the level

of cortisol in the blood. When the level rises, the hypothalamus slows down the production of CRH. When cortisol levels become reduced, the hypothalamus stimulates the pituitary gland to produce more CRH. But in a person who is depressed, the hypothalamus may continuously influence the pituitary to produce CRH without regard to the amount of cortisol present in the blood [4].

In this backdrop, this study was undertaken to find out

1. Prevalence of Insomnia in patients suffering from Major depressive disorder (MDD).
2. To estimate serum cortisol level and Insomnia score in patients suffering from MDD.
3. To find out correlation of serum cortisol level and Insomnia score with MDD.

MATERIALS & METHODS

This Case control study was undertaken in Department of Biochemistry, College of Medicine & Sagore Dutta Hospital in collaboration with Department of Psychiatry of same Institute. The study period was from June, 2014 to Dec, 2015. The study was approved by Institutional Ethics Committee.

Selection of study subjects

All patients who were suspected to suffer from Major depressive disorder (MDD) were selected from the Psychiatry outdoor of College of Medicine & Sagore Dutta Hospital. These patients were first evaluated by detailed history taking and clinical examination through a structured proforma designed for this study. Then they were screened with WHO Five well being index. The raw score was calculated. When raw score was below 13 or if the patient had answered 0 to 1 to any of the 5 items, they were further tested. Patients were diagnosed as having major depressive disorder according to the Structured Clinical Interview for DSM-IV [5] and who scored at least 14 points on Major Depression Inventory (MDI). This inventory was also used to classify the patients according to ICD 10 criteria for depression [6].

The exclusion criteria were significant psychiatric co-morbidity, organic mental disorder, mental retardation, bipolar disorder, intake of any psychotropic drugs during and at least 1 week before the study, substance abuse, history of endocrine disorders, pregnancy, postpartum depression, lactation, and any condition other than depression-related insomnia.

Apparently healthy age and sex matched individuals were assessed using General Health Questionnaire (GHQ 12). A score of less than or equal to 15 were considered as not to suffer from major

psychiatric illness [7]. Such individuals were selected as control group.

Informed consents were taken from the patients or legal guardians and from the control subjects. Insomnia was detected using Athens Insomnia scale (AIS 8) from all study subjects (cases & controls) to find out sleep disturbance in whole study population. AIS 8 are a self report questionnaire containing 8 items and designed to measure the severity of insomnia. The first 5 items aim to measure sleep induction, awakening during night, final awakening, total sleep duration, and general sleep quality. Remaining 3 items refer to consequences of insomnia during the following day, specifically, wellbeing, functional capacity and sleepiness. All these 8 factors are rated on a 0 – 3 scale and the sleep is finally evaluated from the cumulative score of all factors. A cut off score of > 6 is considered as insomnia [8].

Sample Collection, Separation & analysis of serum

An amount of 5 ml of fasting blood samples was drawn aseptically from the superficial veins of each of the study subjects (Both cases & controls) in plain vials and allowed to clot. Serum was separated at room temperature and later by centrifugation at 800 g for 10 minutes. Separated serum was kept in refrigerator in aliquots to the maximum for 1 week. Serum of all patients and controls included were investigated for Cortisol concentration by Immuno Enzymometric assay [9].

STATISTICAL ANALYSIS

The parameters, cortisol concentration & AIS were expressed in mean \pm SD. The mean values were compared for significance by student's t test. A p value of <0.05 was considered to be significant. The correlation of AIS & Cortisol with severity of depression was assessed by Spearman rank Correlation test. The analysis was done using Free Statistics Software (v1.1.23-r7), Office for Research Development and Education, (URL http://www.wessa.net/rwasp_correlation.wasp/)

RESULT

A total of 212 subjects (106 cases of depression & 106 age & sex matched controls) were enrolled for the study. Among the 106 patients, 4 patients did not turn up for blood collection and 1 patient did not give consent for the study. Thus a total of 101 cases suffering from depression (Male: 24, female: 77) and 106 (Male: 30, female: 76) controls were studied. Among all depression cases, 19 cases were of mild grade, 37 were of moderate and 45 were suffering from severe depression.

It has been found that 91 out of 101 MDD cases had AIS score > 6. Thus the prevalence of

insomnia was calculated to be 90.1%. Mean concentration of cortisol was found to be 10.2 ± 5.5 and 6.8 ± 1.6 $\mu\text{g/dl}$ and AIS score was found to be 11.5 ± 4.3 & 3.8 ± 1.6 in cases & controls respectively. Both the parameters were found to be statistically significant. The cases were further classified according to severity.

Concentration of cortisol and AIS were calculated in each grade, and shown in Table 1. The correlation of cortisol and AIS score with severity of depression was assessed by Spearman rank correlation test, the results of which are shown in Table 2. AIS score was found to be significantly correlated with depression.

Table 1: Serum cortisol and Insomnia score in study population

Study population	Serum Cortisol ($\mu\text{g/dl}$)	Insomnia score (by AIS)
Control (n = 106)	6.8 ± 1.6	3.8 ± 1.6
Cases (n = 101)	$10.2 \pm 5.5^*$	$11.5 \pm 4.3^*$
Mild (n = 19)	8.9 ± 4.9	$7.7 \pm 3.6^*$
Moderate (n = 37)	$10.5 \pm 6.6^*$	$10.5 \pm 3.8^*$
Severe (n = 45)	$10.6 \pm 4.7^*$	$13.8 \pm 3.6^*$

All results are expressed in mean \pm SD; *p < 0.05, when compared by Student t test

AIS: Athens Insomnia Scale

Table 2: Estimation of correlation coefficient of AIS & Cortisol with MDD

Parameter	Spearman rank Correlation coefficient	T value	P value
AIS vs MDD	0.5354	6.31	$p < 0.000001^*$
Cortisol vs MDD	0.1840	1.86	$p = 0.065854$

*Statistically Significant

DISCUSSION

Sleep and depression interact in a complex manner. Insomnia has been found to increase the vulnerability to depression, numerous sleep abnormalities are linked to depressive illness, including decreased REM latency, decreased sleep continuity, decreased slow-wave sleep (SWS), increased REM percentage, decreased sleep efficiency, and increased wakefulness. More than 8 epidemiologic studies have shown that insomnia symptoms at baseline predict an increased risk for depression at follow up, anywhere from 1 to 3 years later [10]. Typical sleep symptoms in depression include, difficulty initiating sleep (initial insomnia), difficulty maintaining sleep (mid insomnia), and early morning waking (terminal insomnia). Hypersomnia is a pattern that is part of the ‘signature’ of bipolar depression [11]. In our study, prevalence of insomnia was found to be 90.1%, which is quite high. Insomnia score was also found to be significantly higher in depression cases.

Moreover, grade wise analysis shows that Insomnia score is significantly elevated in mild, moderate and severe depression. Sleep deprivation has a stimulatory effect on the HPA axis, which causes hyper secretion of CRH. This results in increased cortisol levels in plasma, urine and CSF. Exaggerated cortisol responses to ACTH and enlarged pituitary and adrenal glands also found to occur in individuals suffering from severe mood disorders [12]. One study group showed a significant increase in serum level of cortisol ($14.5 \pm 3.01 \mu\text{g/dl}$) in patients with depression than in control group ($11.56 \pm 1.33 \mu\text{g/dl}$). We also found, significantly high level of cortisol in MDD cases.

However, the elevated level was not found to be significant in cases with mild depression, but it was found to be significantly elevated in cases with moderate and severe depression. The sample size of group containing patients with mild depression is small (n = 19), which may be a reason of such result. It is also possible that in mild cases of depression HPA axis stimulation is not sufficient enough to cause significant rise in cortisol level.

The neuro-endocrine studies are mainly focused on the elevation of cortisol in serum and in the saliva [13–15].

However, correlation analysis (Table 2) shows that Insomnia can be significantly correlated with severity of depression but serum cortisol does not. Maes *et al.*; suggested that about 50% of patients with major depression hyper secrete cortisol, but the rate depends on the population sampled [16]. The findings of Strickland *et al.*; [17] suggested that increased cortisol secretion is not a characteristic feature of depression in the community. Moreover, Julia Ross observed that there are 3 biochemical causes of insomnia in depression. First is deficiency of serotonin or dopamine, second is, deficiency of GABA and third is excess of cortisol. Thus excess cortisol is not the only condition that is associated with Depression [18]. Individuals differ in response to stress and adversity, which is also a determining factor for the level of cortisol in blood.

CONCLUSION

Insomnia is considered as a cause as well as one of the manifestations of Depression. It appears that

HPA axis over activity which is found in stressful events can exert its effect by increased release of cortisol as well as altering sleep pattern. Reducing cortisol levels and stabilizing HPA axis dysfunction can be a very effective approach to control patients with insomnia and depression.

LIMITATION

In addition to Serum cortisol level, Dexamethasone Suppression test could be done for better assessment of the secretion pattern of cortisol in Depression. But due to infra structural constrain, it could not be done.

ACKNOWLEDGEMENT

The authors acknowledge the help from Prof. Saibal Mukherjee, Principal, College of Medicine & Sagore Dutta Hospital & Dipta Kanti Mukhopadhyay, Associate Professor, Department of Community Medicine of the same Institute. The support of The West Bengal University of Health Sciences is also duly acknowledged.

REFERENCES

1. Taylor DJ, Lichstein KL, Durrence HH, Riedel BW, Bush AJ; Epidemiology of insomnia, depression and anxiety. *Sleep* 2005; 28:1457-64.
2. Tsuno N, Besset A, Ritchie K; Sleep and depression. *J Clin Psychiatry* 2005; 66:1254-69.
3. Bunney JN, Potkin SG; Circadian abnormalities, molecular clock genes and chronobiological treatments in depression. *Br Med Bull* 2008; 86: 23-32.
4. Heim C, Owens MJ, Plotsky PM, Nemeroff CB; Persistent changes in corticotropin-releasing factor systems due to early life stress: relationship to the pathophysiology of major depression and post-traumatic stress disorder. *Psychopharmacol Bull* 1997; 33:185-192.
5. DSM-IV; Diagnostic and Statistical Manual of Mental Disorders, Am Psychiatric Assoc., Washington DC, 1994.
6. Bech P, Rasmussen NA, Olsen R, Noerholm V, Abildgaard W; The sensitivity and specificity of the Major Depression Inventory, using the Present State Examination as the index of diagnostic validity. *J Affect Dis* 2001; 66:159-164.
7. Goldberg D; Identifying psychiatric illnesses among general medical patients. *British Medical Journal* 1985; 291: 161-162.
8. Soldatos CR, Dikeos DG, Paparrigopoulos TJ; Athens Insomnia Scale: validation of an instrument based on ICD-10 criteria. *J Psychosom. Res* 2000; 48: 555-560.
9. Burtis CA, Ashwood ER, editors, Tietz Fundamentals of Clinical Chemistry, WB Saunders, 5th ed. Philadelphia: 2001; 862.
10. Riemann D, Voderholzar U; Primary insomnia: a risk factor to develop depression. *J Affect Disord* 2003; 76: 255 - 9.
11. Berk M, Malhi GS, Mitchell PB; Scale matters: the need for a Bipolar Depression Rating Scale (BDRS). *Acta Psychiatr Scand* 2004; 422(Suppl):39-45.
12. Mohamad Sayyed Bakheet; Serum cortisol levels in depression patients. *Int J Med Biomed Sc* 2013; 1(1): 005-008.
13. Heuser I, Deuschle M, Luppa P, Schweiger U, Standler H, Weber B; Increased diurnal plasma concentration of DHEA in depressed patients. *J Clin Endocrinol Metab* 1998; 83: 3130-3133.
14. Weber B, Lewicka S, Deuschle M, Colla M, Vecsei P, Heuser I; Increased diurnal plasma concentrations of cortisone in depressed patients. *J Clin Endocrinol Metab* 2000; 85 (3): 1133-1136.
15. Goodyer IM, Herbert J Altham, PME; Adrenal steroid secretion and major depression in 8 to 16 year-olds: Influence of cortisol/DHEA ratio at presentation on subsequent rates of disappointing life events and persistent major depression. *Psychol Med* 1998; 28: 265-273.
16. Maes M, Calabrese J, Meltzer HY; The relevance of the in-versus outpatient status for studies on HPA-axis in depression: spontaneous hypercortisolism is a feature of major depressed inpatients and not of major depression per se. *Progress in Neuro psycho pharmacology and Biological Psychiatry* 1994; 18: 503 -517.
17. Strickland PL, Deakin J F W, Percival C; The bio-social origins of depression in the community. Interactions between social adversity, cortisol and serotonin neurotransmission. *British Journal of Psychiatry* 2002; 180: 168 -173.
18. Julia Ross; Eliminating the top causes of insomnia: Effective Nutritional Intervention for Neurotransmitter Deficiency and Cortisol Excess. *The Townsend Letter--Insomnia* 2011; 1- 7.