

Review Article

Effects of Cadmium Toxicity in Plants: A Review Article

Pallavi Jali, Chinmay Pradhan*, Anath Bandhu Das

Department of Botany, Utkal University, VaniVihar, Bhubaneswar – 751004, Odisha, India

***Corresponding author**

Chinmay Pradhan

Email: chinmay_pr@yahoo.com

Abstract: Heavy metals, for example, cadmium, copper, lead, chromium and mercury are considered as major environmental pollutants. Among all heavy metals, cadmium (Cd) is a non-essential metal which has pulled in the consideration of specialists in soil science and plant nutrition because of its potential harmfulness and toxicity to living beings, furthermore because of its relative mobility in the soil plant system. This survey underlines the dangerous side effects or toxic symptoms of Cd in plants i.e. growth retardation, alternations of photosynthetic activity, change in stomatal movement, enzymatic activities, protein metabolism, membrane functioning furthermore outlines the mechanisms of cadmium uptake, translocation, deposition and cadmium-induced oxidative stress. This survey/review may likewise help in interdisciplinary study to evaluate/access ecological significance of the metal toxicity and stress.

Keywords: Cadmium, toxicity, photosynthesis, stress.

INTRODUCTION

Heavy metals are characterized as the metals that have a density greater than 5 g cm^3 . Among 90 naturally occurring elements, 53 are designated as heavy metals and some are biologically important [1]. On the basis of their solubility under physiological conditions, 17 heavy metals can be considered for being accessible to the living cells and have significant role in plant and animal communities within various ecosystems [2]. They are considered as significant environmental pollutants that [3,4,5]. As per [6], non-contaminated soil contain Cd concentration ranging from 0.04 to 0.32 mM. Soil that have a Cd concentration ranging from 0.32 to around 1 mM are considered as contaminated to a moderate level. As Cd is naturally present in all soils, so all the food stuffs will contain some amount of Cd and subsequently all living beings including humans are exposed to natural levels of Cd. It has been accounted earlier that leafy vegetables and potato tubers accumulate more elevated amounts of Cd when contrasted with fruits and cereals. In addition, tillage and crop rotation practices have a more noteworthy effect upon the Cd content of food in comparison to that of Cd concentrations in soil [7]. Cadmium accumulation in the environment is presently turning into an essential reason for environmental pollution. Cadmium (Cd) is a standout amongst the most deleterious trace heavy metals both to plants and animals. Industrialization and culture cause Cd the most harmful and widespread pollutants in agricultural soils, and soil-plant-environment system [8]. The maximum

tolerable intake limit of Cd for human beings, suggested by FAO/WHO is $70 \mu\text{g/day}$ [9]. Cd contamination is of expanding investigative interest since Cd^{2+} is taken up readily by the roots of all most all plant species and its harmfulness/toxicity is thought to be 2–20 times higher than other heavy metals [10]. Cd phytotoxicity is a minor, yet an important issue, particularly in exceedingly heavy metal contaminated locales/regions, where a diminishing in agricultural crop productivity has been noticed. The main aim of this review is to focus our present understanding on the factors that utmost the development of plants exposed to Cd treatment.

Mechanism of cadmium uptake, translocation

Plants respond to expanded levels of Cd in soil differentially, regarding the capacity of different plants species uptake and transport Cd. Cd is effectively transported inside plants [11] as metallo-organic complexes. The bio-accessibility of Cd in soil relies on its pH, concentration, redox potential, temperature, and concentration of different elements. Rhizosphere acidification and carboxylase exudation are considered as potential targets for upgrading accumulation of metals [12]. Mechanisms involved in the uptake of Cd by the root of plants generally includes competition for absorption sites between the heavy metals and a few mineral nutrients having comparable/similar chemical properties [13]. The decrease of metals like K, Ca and Mg in tissue because of high level of Cd has been seen

in tomato and cucumber plants [14], maize [15] and lettuce. An antagonistic relation amongst cadmium and zinc and their active absorption has been seen in roots of lettuce [16]. Other mineral nutrients, for example, nitrate, which does not have similar chemical properties with Cd, are also influenced by its presence. Cd first enters into the roots and harms the root system first [17]. The mechanism controlling the Cd uptake by roots of plants and amassing in consumable/edible parts are yet not fully understood. The electrochemical potential difference between the action of Cd²⁺ in the cytosol and in the root apoplasts controls the absorption of Cd across the plasma membrane of root cells. Enough energy is given by the large membrane potential to drive Cd²⁺ uptake even at low levels/doses of Cd²⁺. The energy of Cd²⁺ absorption by roots demonstrates bi-phasic qualities with saturable components at low Cd²⁺ activities in the absorption solution and a linear components at higher Cd activities [18]. Absorption of Cd can likewise happen as inorganic complexes (for example, CdCl⁺, CdCl₂ and CdSO₄) [19] or else as organic complexes, for example, phytometallophore complexes. [20] noticed that Zn(II) phytometallophore complexes were promptly taken up by maize roots however the binding sites in the root plasma membrane are not exceptionally particular for Fe(III) phytometallophores, along these lines likewise permitting the transport of different metals like Cd. In any case, there is no immediate confirmation for Cd binding with phytometallophores during its transport in the root cells. Different metals, particularly Zn²⁺ collaborate with Cd and lessen uptake during Zn-insufficient/deficient conditions and it is clear from the evidence that cereal roots developed under Zn-insufficient /deficient conditions are implicated in diminished Cd uptake following the application of Zn [21]. Cadmium effectively penetrates the root system of xylem through both apoplastic and/or symplastic pathway [22] and therefore reaches tissues of plants specifically aerial [23]. The metal content is greater or more prominent in the root than in the above ground tissues of plants regardless of the distinction of metal ions mobility [24]. Generally Cd particles are held in the roots and just little amounts are transported to the above ground parts [25]. The concentration of Cd in plants diminishes in the order: root > leaves > fruits > seeds/grains [26, 27]. The degree of Cd transport into edible organs varies generally among crops. In soybean

more than 98% of the Cd accumulated was held by roots and just 2% was transported to shoots/above ground parts [25]. Additionally, Cd was effectively transported to the shoots and leaves yet was not detected in fruits of tomato plants [28]. After uptake by the roots Cd is transported to the shoots, through the cells of vascular bundles. Transport of the trace metal is likewise directed by vascular tissues [29]. There are various cell membrane obstructions that Cd must cross to enter consumable/edible plant organs and this is particularly true for seeds and grains.

Cadmium Deposition

The aggravates that bind Cd in mature seeds amid their development are not known. Cadmium may bind to phytate (myo-inositol hexaphosphate) in globoid crystals inside the protein structure of developing seeds. Different metals (Fe, Zn, Mn, Mg and Ca) have been accounted for to be associated with phytate inside globoid crystals of these organelles. [30,31, 32] reported that phytate globular deposits containing Zn was formed in small vacuoles of root cells inside the elongation zone of roots of soybean, maize and wheat. Furthermore, Cd was not bound to phytic acid in these little root-cell vacuoles. Alternatively, Cd could be found to second class metallothioneins in developing seeds and grains since genes for the expression of these sulphhydryl-rich proteins have been accounted for in seeds of some plant species like wheat and maize [33,34]. Further research directed to determine the major type of Cd in edible parts of crops demonstrated that in oat (*Avena sativa* L.) roots, Cd transport from cytosol to the vacuole over the tonoplast is exhibited through Cd²⁺/H⁺ antiport movement and stored in cell compartments [35].

Effect of Cd on stress proteins

Changes in the environment can bring about change in gene expression, in this manner prompting change in the diversity of proteins in the cell. Therefore, changes in level of protein under stress conditions can be molecular markers for the manifestations of the responses to stress in living beings. In plants, the proteomics methodology is developed as a critical strategy for research on stress resistance/tolerance [92]. Heat-shock proteins (HSPs) are known as proteins that have functions to tolerate stress in eukaryotes.

Table 1: Toxic symptoms induced by Cadmium

Parameters	Effects	References
Photosynthesis	Retards photosynthesis	[36,37,38]
	Hindered stomatal opening in <i>Syzygium aromaticum</i> , <i>Medicago sativa</i> , <i>Glucine max</i>	[39,40]
	Chlorotic leaves, changed ratios of chlorophyll a and b, decreasing net photosynthetic rate	[41,42,43,35,44]
	Destructs the photosynthetic apparatus particularly the light harvesting complex II	[45]
	Destructs photosystems I and II	[46,47]
	Inhibition of root Fe(III) reductase	[48]
	Overall destruction of photosynthetic efficiency	[49,50,37]
	Diminished chlorophyll and carotenoids content, and increased non-photochemical quenching in <i>Brassica napus</i>	[51,52,53,54,55, 56,57,58,59]
Fresh weight and dry mass	Diminished the fresh mass in <i>Vigna radiata</i>	[60]
	Decline in root and shoot mass in <i>Vigna ambacensis</i>	[61]
	Decrease in dry mass in <i>Cicer arietinum</i>	[62,63]
Protein	Inhibition of protein synthesis	[64,65,66,67]
Carbonic anhydrase	Retards the activity of carbonic anhydrase	[68]
Proline	Increase in proline in <i>Oryza sativa</i> , <i>Brassica napus</i> , <i>Armeria moritima</i> , <i>Helianthus annus</i> , <i>Brassica juncea</i>	[69,70,71,72,73, 74]
Lipid peroxidation	Membrane leakage, change of lipid composition	[75,76,77]
Cellular concentrations	Changes in cellular concentrations of essential micronutrients like iron, calcium, manganese, zinc	[78,79,80,]
Root ultrastructure	Inhibition of root elongation, increase in volume of cortex cells, damage to epidermis	[81,82,41,83]

Table 2: Cadmium-induced signaling events mediated by reactive oxygen species (ROS) in different plant species

Plant species (References)	Cd concentration	Time of treatment	Signaling events
<i>Oryza sativa</i> [84]	100µM Cd(NO ₃) ₂	13 days	Accumulation of H ₂ O ₂ and modification of the auxin signaling pathway and/or cell-cycle gene expression
<i>Oryza sativa</i> [85]	5mM CdCl ₂	24 h	H ₂ O ₂ accumulation dependent on NADPH-oxidase and phosphatidylinositol 3-phosphate
<i>Oryza sativa</i> [86]	100,200, 400mM CdCl ₂	1 h	Regulation of MAP kinase activity by :non- enzymatic (OH•) and enzymatic ROS production (O ₂ •- or H ₂ O ₂) involving NADPH oxidase, CDPKs, PI3 kinase, and closing of the mitochondrial pore Regulation of NADPH oxidase and CDPKs activity by Ca ²⁺
<i>Pisum sativum</i> [87,88]	50µM CdCl ₂	15days	Accumulation of O ₂ •- and H ₂ O ₂ , Ca ²⁺ -dependent decrease in NO levels, activation of peroxidases and NADPH oxidase
<i>Pisum sativum</i> [89]	100µM CdCl ₂	48 h	Necrotic cell death associated with NO and H ₂ O ₂ generation
<i>Nicotiana tabacum</i> (cell suspension) [90]	5mM CdCl ₂	15 min	Oxidative burst mediated by Ca ²⁺ , calmodulin and protein phosphorylation
<i>Nicotiana tabacum</i> (cell suspension) [91]	3mM CdCl ₂	8 h	Three waves of oxidative stress: (1) transient, NADPH oxidase-dependent accumulation of H ₂ O ₂ (2) increased production of O ₂ •- in mitochondria (3) fatty acid hydroperoxide accumulation concomitant with necrotic type of cell death Regulation of NADPH oxidase activity involving Ca ²⁺ -mediated signaling and protein phosphorylation




Fig-1: A general model for local and systemic stress signaling in plants

Table 1: Effect of Cd on stress proteins

Plant	Treatment	Effects	References
<i>Zea mays</i>	Cd exposure	Production of 70 kDa phosphoprotein (HSP)	[93]
<i>Lycopersicon peruvianum</i>	Pre-treatment with a short heat stress before Cd exposure	Preventing membrane damage. HSP17 (molecular weight 17 kDa) and HSP70 proteins were additionally found in the cytosol of heat-shocked cells.	[94]
<i>Pisum sativum</i>	Cd exposure	Pathogen-related proteins PrP4A and HSP71 were observed.	[88]
<i>Triticum aestivum</i>	50 μ M CdCl ₂ for 48 h	51-kDa soluble protein was found and protein was assigned as a Cd stress-related protein.	[95]
<i>Populus tremula</i>	Cd for a short term (14 days) or a more longer term (56 days) treatment	Stress related proteins, as HSPs, proteinases, and pathogenesis-related proteins, increased in abundance in leaves.	[96]
<i>Oryza sativa</i>	Cd exposure	Affected the synthesis of 36 proteins	[97]
<i>Solanum lycopersicum</i>	Low Cd concentration (10 μ M)	changes in 36 polypeptides	[98]
	Higher Cd level (100 μ M)	changes in 41 polypeptides	[98]
<i>Arabidopsis thaliana</i>	10 μ M Cd	Among 730 determined proteins 21 were up-regulated in response of Cd	[99]

CONCLUSION

Cadmium is a non-essential heavy metal which is phytotoxic and has negative impact on plants. It retards various physiological, morphological, biochemical and molecular activities of plant species. It is easily taken up by the roots of plants and causes severe damage to plant system. At higher concentration it may even cause death of the plant species. At low concentrations it can induce production of stress proteins and other secondary metabolites which indirectly help plants to resist against the oxidative damage caused by cadmium. This review highlights the effects of cadmium on various plant activities.

ACKNOWLEDGEMENT

The authors acknowledge the financial support provided by DST-INSPIRE fellowship, UGC-DRS-SAP scheme and PURSE Grant to P.G Department of Botany, Utkal University Bhubaneswar, India.

REFERENCES

- Hasan SA, Fariduddin Q, Ali B, Hayat S, Ahmad A. Cadmium Toxicity and tolerance in plants. J Environ. Biol. 2009;30(2):165-174.
- Weast RC. CRC Hand book of Chemistry and Physics. 64th edition, CRC Press, Boca Raton.1984.
- Angelone M, C Bini. Trace element concentration in the soil and plants of Western Europe. In: Biogeochemistry of trace metal (ed.: M.I. Arbor). D.C. Adriano, Lewis Publishers. 1992;19-670.

4. Berry WL. Plant factors influencing the use of plant analysis as a tool for biogeochemical properties. In: Mineral exploration : Biological systems and organic matter (ed.: D. Carlisle., I.R. Kaplan and J.R. Watterson).(5)Prentice-Hall, New, Netherland. 1986;13.
5. Kevresan S, Krstic B, Popovi M, Kovacev L, Peyevic S, Kandrak J, Malencici K. Biochemical changes in sugar beet lines in dependence of soil moisture. Biol. Plant. 1998;40:245-250.
6. Wagner G. Accumulation of cadmium in crop plants and its consequences to human health. In: Sparks DL (ed.) Advances in Agronomy. Newark, USA: Academic Press. 1993;173-212.
7. Mench M, Vangronsveld J, Lepp N, Edwards R. Physicochemical aspects and efficiency of trace element in immobilisation by soil amendments. In: Vangronsveld J & Cunningham S (ed.) *In-situ* Inactivation and Phytoremediation of Metal Contaminated Soils, Georgetown TX: Springer-Verlag, Landes Company, 1998;151-182.
8. Dong J, Mao WH, Zhang GP, Cai Y. Root excretion and plant tolerance to cadmium toxicity. Plant Environ. J. 2007;53(5):193-200.
9. Vassilev A, Yordanov I. Reductive analysis of factors limiting growth of cadmium treated plants. Bulg. J. Plant Physiol. 1997;23(3-4):114-133.
10. Shah K, Dubey S. Effect of cadmium on RNA level as well as activity and forms of Ribonuclease in growing Rice seedlings. Plant Physiol. 1995;33:577- 584.
11. Epstein E, Bloom AJ. Mineral Nutrition of Plant Principles and Perspective. 2nd Edn. Sinauer Associates, Inc. Publishers, Massachusetts, 2007.
12. Clemens S, Palmgreen MG, Kramer U. A long way ahead: Understanding and engineering plant metal accumulation. Trends Plant Sci. 2002;7:309-315.
13. Jarvis C, Jones LHP, Hopper MJ. Cadmium uptake from solution by plants and its transport from roots to shoots. Plant and Soil. 1976;44:179-191.
14. Burzynski M. The uptake and accumulation of phosphorus and nitrates and the activity of nitrate reductase in cucumber seedlings treated with Pb and Cd. Acta. Soc. Bot. 1988;57:349-359 .
15. Walker WM, Miller JE, Hassett JJ. Effect of lead and cadmium upon calcium magnesium, potassium and phosphorus, concentration in young corn plants. Soil Sci. 1977;124:145-151.
16. Costa G, Morel JL. Water relations gas exchange and amino acid content in Cd treated lettuce. Plant Physiol. Biochem.1994;32:561-570.
17. Sanita di Toppi L, Lambardi M, Pazzagli L, Cappugi G, Durante M, Gabbriellini R. Response to cadmium in carrot in vitro plants and cell suspension cultures. Plant Sci. 1998;137:119-129.
18. Costa G, Morel IL. Cadmium uptake by *Lypins albus* (L.) cadmium excretion, a possible mechanism of cadmium tolerance. J. Plant Nutr.1993;16:1921-1929.
19. McLaughlin MJ, Tiller KG, Naidu R, Stevens DP. Reviewd: The behaviour and impact of contaminants in fertilizers. Aust. J. Soil Res.1996;34:1-54.
20. Von Wiren N, Marschner H, Romheld V. Roots of iron-efficient maize also absorb phyto siderophore chelated zinc. Plant Physiol. 1996;111:1119-1125.
21. Welch RM. Micronutrient nutrition of plants. Critical Rev. Plant Sci.1995;14:49-82.
22. Salt DE, Rauser WE. MgATP-dependent transport of phytochelatin across the tonoplast of oat roots. Plant Physiol. 1995;107:1293-1301.
23. Yang MG, Lin Xu, Yang XE. Impact of Cd on growth and nutrient accumulation of different plant species. Chin. J. Appl. Ecol. 1998;19:89-94.
24. Ramas I, Esteban E, Lucena JJ, Gorate A. Cadmium uptake and subcellular distribution in plants of *Lactuca* sp. Cd-Mn interaction. Plant Sci. 2002;162:761-767.
25. Cataldo CD, Garland TR, Widung RE. Cadmium uptake, kinetics in intact soybean plants. Plant Physiol. 1983;73:844-848.
26. Blum WH. Cadmium uptake by the higher plants. In: Proceeding of extended abstracts from the fourth International Conference on the Biogeochemistry of Trace Elements, University of California, Berkeley, USA.1997;109-110.
27. Sharma M, Gautam KH, Handique AK. (2006). Toxic heavy metal stress in paddy: metal accumulation profile and development of a novel stress protein in seed. Ind. J. Plant Physiol. 2006;11:227-233.
28. Morel R, Palacios G, Gomez I, Navarro-Pedreno J, Mataix J. Distribution and accumulation of heavy metal (Cd, Ni and Cr) in tomato plants. Fresenius Environ. Bull.1994;3:395-399.
29. Kuppelwieser H, Feller U. Transport of Rb and Sr to the ear in mature excised shoots of wheat effects of temperature and stem length on Rb removal from the xylem. Plant and Soil. 1991;132:281-299.
30. Van Steveninck RFM, Van Steveninck ME, Wells AJ, Fernando DR. Zinc tolerance and the binding of zinc as zinc phytate in *Lemna minor* x-ray microanalytical evidence. J. Plant Physiol. 1990;137:140-146.
31. Van Steveninck, RFM, Van Steveninck ME, Fernando DR. Heavy metal (Zn-Cd) tolerance in selected clones of duck weed (*Lemna minor*). Plant and Soil.1992;146:271-280.
32. Van Steveninck RFM, Babare D, Fernando DR, Van Steveninck ME. The binding of zinc, but not cadmium by phytic acid in roots of crop plant. Plant and Soil. 1994;167:157-164.
33. White CN, Rivin CJ. Characterization and expression of a cDNA encoding a seed specific

- metallothionein in maize. *Plant Physiol.* 1995;108:831-832.
34. Hsieh HM, Liu WK, Chang D, Huang PC. RNA expression patterns of a type 2 metallothioneinlike gene from rice. *Plant Mol. Biol.* 1996;32:525-529.
35. Prasad MN. Cadmium toxicity and tolerance in vascular plants. *Environ. Expt. Bot.* 1995;35:525-545.
36. Greger M, Johansson M, Stihl D, Humza K. Foliar uptake of Cd by pea (*Pisum sativum*) and sugar beet (*Beta vulgaris*). *Physiol. Plant.* 1994;88:563-570.
37. Chugh LK, Sawhney SK. Photosynthetic activities of *Pisum sativum* seedlings grown in presence of cadmium. *Plant Physiol. Biochem.* 1999;37:297-303.
38. Vassilev A, Perez-Sanz A, Semane B, Carteer R, Vangronsveld J. Cadmium accumulation and tolerance of two salix genotypes hydroponically grown in presence of cadmium. *J. Plant Nutr.* 2005;28:2159-2177.
39. Barcelo J, Poschenrieder C, Andreu I, Gunse B. Cadmium induced decrease of water stress resistance in bush bean plants (*Phaseolus vulgaris* L. cv. Vontendev). I. Effect of Cd on water potential, relations water content and cell wall elasticity. *J. Plant Physiol.* 1986;125:17-25.
40. Poschenrieder C, Gunse B, Barcelo J. Influence of cadmium on water relation, stomatal resistance, and abscisic acid content in expanding bean leaves. *Plant Physiol.* 1989;90:1365-1371.
41. Ouzounidou G. Cu-ions mediated changes in growth, chlorophyll and other ion contents in a Cu-tolerant *Koeleria splendens*. *Biol. Plant.* 1995;37:71-78.
42. Viehweger K, Geipel G. Uranium accumulation and tolerance in *Arabidopsis halleri* under native versus hydroponic conditions. *Environ Exp Bot.* 2010;69:39-46.
43. Monnet F, Vaillant N, Vernay P, Coudret A, Sallanon H, Hitmi A. Relationship between PSII activity, CO₂ fixation, and Zn Mn and Mg contents of *Lolium perenne* under zinc stress. *J. Plant Physiol.* 2001;158:1137-1144.
44. Jali P, Pradhan C, Das AB. A comparative analysis of physiological and biochemical responses to low doses of cadmium in two important varieties of *Oryza sativa* L. of Odisha, India. *Int. J. Sci and research.* 2014;12(3):1920-1927.
45. Krupa Z. Cadmium induced changes in the composition and structure of the light-harvesting complex II in radish cotyledons. *Physiol. Plant.* 1988;73:518-524.
46. Siedlecka A, Baszynsky T. Inhibition of electron flow around photosystem I in chloroplasts of cadmium treated maize plants is due to cadmium induced iron deficiency. *Physiol. Plantarum.* 1993;87:199-202.
47. Siedlecka, A, Krupa Z. Interaction between cadmium and iron and its effect on photosynthetic capacity of primary leaves of *Phaseolus vulgaris*. *Plant Physiol. Biochem.* 1996;34:833-841.
48. Alkantara E, Romerza FJ, Canett M, De LaGuardia MD. Effect of heavy metals on both induction and function of root Fe(III) reductase in Fe-deficient cucumber (*Cucumis sativus* L.) plants. *J. Exp. Botany.* 1994;45:1893-1898.
49. Sheoran IS, Signal HR, Singh R. Effect of cadmium and nickel on photosynthesis and the enzymes of photosynthetic carbon reduction cycle in pigeon pea (*Cajanus cajan* L.). *Photosynth. Res.* 1990;23:345-351.
50. Krupa Z, Quist G, Hurner NPA. The effect of cadmium on photosynthesis of *Phaseolus vulgaris* – A fluorescence analysis. *Physiol. Plant.* 1993;88:626-630.
51. Larsen PB, Degenhart J, Stenzler LM, Howell SH, Kochian LV. Aluminium-resistant *Arabidopsis* mutant that exhibit altered pattern of aluminium accumulation and organic acid release from roots. *Plant Physiol.* 1998;117:9-18.
52. Czuba M, Ormond D. Effect of cadmium and zinc on ozone induced phytotoxicity in cress and lettuce. *Can. J. Bot.* 1973;52:645-649.
53. Griffiths PG, Sasse JM, Yokota T, Comeron DW. 6-deoxytyphasterol and 3-dehydro-6-deoxytyphasterol, possible precursors to brassinosteroids in pollen of *Cupressus arizonica*. *Bios. Biotech. Biochem.* 1995;59:956-959.
54. Stiborova M, Doubrovava M, Brezninova A, Frederick A. Effect of heavy metal ion on growth and biochemical characteristics of photosynthesis of barley *Hordeum vulgare* L. *Phytosynthetica.* 1986;20:418-425.
55. Bishnoi NR, Sheoran IS, Singh R. Influence of cadmium and nickel on photosynthesis and water relations in wheat leaves of differential insertion level. *Phytosynthetica.* 1993;28:473-479.
56. Ferretti M, Ghisi R, Merlo L, Dallavecchia F, Passera C. Effect of cadmium on photosynthesis and enzymes of photosynthesis sulfate and nitrate assimilation pathways in maize (*Zea mays* L.). *Phytosynthetica.* 1993;29:49-54.
57. Imai E, Obata H, Shimizu K, Komiya T. Conservation of glutathione into cadystins and their analogs catalyzed by carboxypeptidases. *Biosci. Biotech. Biochem.* 1996;60:1193-1194.
58. Phetsombat S, Kruatrachue M, Pokethitiyook P, Upatham S. Toxicity and bioaccumulation of cadmium and lead in *Salvinia cucullata*. *J. Environ. Biol.* 2006;27:647-652.
59. Pandey S, Gupta K, Mukherjee AK. Impact of cadmium and lead on *Catharanthus roseus* – A phytoremediation study. *J. Environ. Biol.* 2007;28:655-662.

60. Shen YY, Dai JY, Hu DC, Gu WL, He Y, Zheng B. Studies on physiological effects of brassinolide on drought resistant in maize. J. Shenyang Agricultural University. 1990;21:191-195.
61. Al Yemens MN. Effect of cadmium, mercury and lead on seed germination and early seedling growth of *Vigna ambacensis* L. Indian J. Plant Physiol. 2001;6:147-151.
62. Wool House, HW. Encyclopedia of plant physiology. In: Toxicity and tolerance in the responses of plants of metals (Eds.: O.L. Lange, P.S. Nobel, C.B. Osmond and H. Ziegler). New Series, Springer-Verlag, Berlin. 1983;(12):245-300.
63. Rana A, Ahmad M. Heavy metal toxicity in legume microsymbiont system. J. Plant Nutr. 2002; 25: 369-386.
64. Foy CD, Chaney RL, White MG. The physiology of metal toxicity in plants. Ann. Rev. Plant Physiol. 1978;29:511-566.
65. Dubey RS, Dwivedi RS. Effect of heavy metal on seed germination and seedling growth of soybean. Nat. Acad. Sci. Letters. 1987;10:121-124.
66. Tamas L, Huttova J, Zigova Z. Accumulation of stress protein in intracellular spaces of barley leaves induced by biotic and abiotic factors. Biol. Plant. 1997;39:387-394.
67. Salgare SA, Acharekar C. Effect of industrial pollutant on growth and content of certain weeds. J. Nature Conserv. 1992;4:1-6.
68. Siedlecka A, Krupa Z. Interaction between cadmium and iron and its effect on photosynthetic capacity of primary leaves of *Phaseolus vulgaris*. Plant Physiol. Biochem. 1996;34:833-841.
69. Saradhi A, Saradhi PP. Proline accumulation under heavy metal stress. J. Plant Physiol. 1981;138:554-558.
70. Roy D, Bhumnia A, Basu N, Banerjee, SK. Effect of NaCl salinity on metabolism of proline in salt sensitive and salt-resistant cultivars of rice. Biol. Plant. 1992;34:159-162.
71. Chandler SF, Thorpe TA. Characterization of growth water relations, and proline accumulation in sodium sulfate tolerant callus of *Brassica napus* L. cv. Westar (Canola). Plant Physiol. 1987;84:106-111.
72. Farago ME. Metal tolerant plant. Coord. Chem. Rev. 1981;36:155-182.
73. Kastori R, Petrovic M, Petrovic N. Effect of excess lead, cadmium, copper and zinc on water relations in sunflower. J. Plant Nutr. 1992;15:2427-2439.
74. Singh PK, Tiwari RK. Cadmium toxicity induced changes in plant water relations and oxidative metabolism of *Brassica juncea* L. plants. J. Environ. Biol. 2003;24:107-112.
75. Kochian LV. Cellular mechanisms of aluminium toxicity and resistance in plants. Annu Rev Plant Physiol Plant Mol Biol. 1995;46:237-260.
76. Hernandez LE, Cooke DT. Modification of the root plasma membrane lipid composition of cadmium-treated *Pisum sativum*. J Exp Bot. 1997;48:1375-1381.
77. Stefanov K, Seizova K, Popova I, Petkov V, Kimenov G, Popov S. Effect of lead ions on the phospholipid composition in leaves of *Zea mays* and *Phaseolus vulgaris*. J Plant Physiol. 1995;147:243-246.
78. Hernandez LE, Lozano-Rodriguez E, Garate A, Carpena-Ruiz R. Influence of cadmium on the uptake, tissue accumulation and subcellular distribution of manganese in pea seedlings. Plant Sci. 1998;13:139-151.
79. Zhang GP, Fukami M, Sekimoto H. Genotypic differences in effects of cadmium on growth and nutrient compositions in wheat. J Plant Nutr. 2000;23:1337-1350.
80. Viehweger K, Geipel G. Uranium accumulation and tolerance in *Arabidopsis halleri* under native versus hydroponic conditions. Environ Exp Bot. 2010;69:39-46.
81. Kidd PS, Llugany M, Poschenrieder C, Gunse B, Barcelo J. The role of root exudates in aluminium resistance and silicon-induced amelioration of aluminium toxicity in three varieties of maize (*Zea mays* L.). J Exp Bot. 2001;52:1339-1352.
82. Vazquez MD, Poschenrieder C, Barcelo J. Ultrastructural effects and localisation of low cadmium concentrations in bean roots. New Phytol. 1992;120:215-226.
83. Barcelo J, Poschenrieder C, Prasad MNV. Structural and ultrastructural changes in heavy metal exposed plants. In: Heavy metal stress in plants. Springer, Berlin, Heidelberg. 2004;223.
84. Zhao FY, Han MM, Zhang SY, Wang K, Zhang CR, Liu T. Hydrogen peroxide-mediated growth of the root system occurs via auxin signaling modification and variations in the expression of cell-cycle genes in rice seedlings exposed to cadmium stress. J. Integr. Plant Biol. 2012;54:991-1006.
85. Hsu YT, Kao CH. Toxicity in leaves of rice exposed to cadmium is due to hydrogen peroxide accumulation. Plant Soil. 2007;298:231-241.
86. Yeh CM, Chien PS, Huang HJ. Distinct signaling pathways for induction of MAP kinase activities by cadmium and copper in rice roots. J. Exp. Bot. 2007;58:659-671.
87. Rodríguez-Serrano M, Romero-Puertas MC, Zabalza A, Corpas FJ, Gómez M, Del Río LA. Cadmium effect on oxidative metabolism of pea (*Pisum sativum* L.) roots. Imaging of reactive oxygen species and nitric oxide accumulation in vivo. Plant Cell Environ. 2006;29:1532-1544.
88. Rodríguez-Serrano M, Romero-Puertas MC, Pazmino DM, Testillano PS, Risueno MC, Del Río LA. Cellular response of pea plants to cadmium

- toxicity: crosstalk between reactive oxygen species, nitric oxide, and calcium. *Plant Physiol.* 2009;150:229–243.
89. Lehotai N, Peto A, Bajkán S, Erdei L, Tari I, Kolbert Z. *In vivo* and *in situ* visualization of early physiological events induced by heavy metals in pea root meristem. *Acta Physiol. Plant.* 2011;33:2199–2207.
90. Olmos E, Martínez-Solano JR, Piqueras A, Hellín E. Early steps in the oxidative burst induced by cadmium in cultured tobacco cells (BY-2line). *J. Exp. Bot.* 2003;54:291–301.
91. Garnier L, Simon-Plas F, Thuleau P, Agnel JP, Blein JP, Ranjeva R. Cadmium affects tobacco cells by a series of three waves of reactive oxygen species that contribute to cytotoxicity. *Plant Cell Environ.* 2006;29:1956–1969.
92. Nanjo Y, Nouri MZ, Komatsu S. Quantitative proteomic analyses of crop seedlings subjected to stress conditions; a commentary. *Phytochemistry.* 2011;72:1263–1272.
93. Reddy GN, Prasad MNV. Tyrosine is not phosphorylated in cadmium-induced HSP cognate in maize (*Zea mays* L.) seedlings: role in chaperone functions? *Biochemical Archives.* 1993;9:25–32.
94. Neumann D, Lichtenberger O, Gunther D, Tschiersch K, Nover L. Heat-shock proteins induce heavy-metal tolerance in higher plants. *Planta.* 1994;194:360–367.
95. Mitra B, Sharma S, Das A, Henry S, Das T, Ghosh P, Ghosh S, Mohanty P. A novel cadmium induced protein in wheat: characterization and localization in root tissue. *Biologia Plantarum.* 2008;52:343–346.
96. Kieffer P, Schroder P, Dommès J, Hoffmann L, Renaut J, Hausman JF. Proteomic and enzymatic response of poplar to cadmium stress. *J. Proteomics.* 2009;72:379–396.
97. Lee K, Bae DW, Kim SH, Han HJ, Liu X, Park HC, Lim CO, Lee SY, Chung WS. Comparative proteomic analysis of the short-term responses of rice roots and leaves to cadmium. *J. Plant Physiol.* 2010;167:161–168.
98. Rodriguez-Celma J, Rellan-Alvarez R, Abadia A, Abadia J, Lopez-Millan AF. Changes induced by two levels of cadmium toxicity in the 2-DE protein profile of tomato roots. *J. Proteomics.* 2010;73:1694–1706.
99. Semane B, Dupae J, Cuypers A, Noben JP, Tuomainen M, Tervahauta A, Karenlampi S, Van Belleghem F, Smeets K, Vangronsveld J. Leaf proteome responses of *Arabidopsis thaliana* exposed to mild cadmium stress. *J. Plant Physiol.* 2010;167:247–254.