

Original Research Article

Morphological Profile and Hormone Receptor Status in Breast CarcinomasDr. R. Manasa Reddy¹, Dr. Vijaya Gattu², Dr. Soumya Vellanki³, Dr. K.P.A Chandra Sekhar⁴¹Assistant Professor, Department Of Pathology, SVS Medical College, Yenugonda, Mahabubnagar, Telangana, India***Corresponding author**

Dr. R. Manasa Reddy

Email: ravipallymanasa84@gmail.com

Abstract: The data on histological and hormone receptor status in breast cancer in an Indian population is limited as receptor status is not routinely carried out for these patients. In our study, receptor status was analyzed and it was correlated with morphological prognostic parameters. The objective is to analyze the morphological prognostic parameters and its correlation with receptor status in Indian women. The sample consisted of 66 specimens of invasive breast cancer received in Department Of Pathology of SVS Medical College and Hospital, Mahabubnagar between year 2012 and 2014. ER, PR and HER-2 status correlates well with histopathological grading. Higher the tumor grade, the more likely that ductal carcinoma will be Her2 + and ER/PR negative or triple negative. Follow up study of these patients is needed to assess the prognostic significance.

Keywords: Breast cancer, estrogen receptor, HER-2/neu, progesterone receptor

INTRODUCTION

Breast carcinoma is the most common malignant tumor and the leading cause of carcinoma death in women with an estimated life time risk of 13% [1]. Fifty eight percent of patients were 50 years or younger. Morphological classification of breast carcinomas divide these tumours into a number of subtypes . Approximately three fourth of mammary carcinoma are invasive ductal or lobular type. Within these, and also among other morphologic categories , tumors display marked heterogeneity in many of their biologic properties . One is the expression of steroid receptors in concert with the oncogene ErbB2/Human epidermal growth factor receptors 2 (HER2). This has important clinical implications, such as selection of patients for endocrine therapy [2, 3].

A large number studies have correlated the presence or absence of tumour estrogen receptor (ER) or Progesterone Receptor (PR) with ultimate clinical outcomes . In addition to pathological grade and stage , we set out to evaluate tumour ER/PR status and HER2/neu expression by Immunohistochemistry (IHC) as a prognostic and predictive factors in breast carcinoma [1].

AIM AND OBJECTIVES

1. To assess ER , PR status and HER2/neu overexpression by IHC of malignant tumors

2. To correlate the ER/PR and HER2/neu status with modified Bloom - Richardson histopathological grading .
3. To study the relationship between clinicopathological parameters, IHC subtypes and histopathological grading.

METHODOLOGY

This is a prospective study of 66 cases conducted in Department Of Pathology, S.V.S Medical College from 2012 to 2014. This study included lumpectomies, mastectomy, modified radical mastectomy cases. The specimens were thoroughly examined and clinical details were analyzed. The Specimen sent in formalin was sliced at 1 cm interval and fixed immediately in 10% NBF. One dedicated block from the tumor not fixed for more than 24 hours in formalin was used for IHC. Four μ thickness sections were cut and taken on poly-L-Lysine coated slides and stained for evaluating ER, PR receptors and HER2/ new expression. And also sections were routinely stained with H & E and histological grading of tumor was done on H&E stained sections according to Modified Bloom –Richardson grading.

Procedure followed for IHC staining is according to guidelines given in Dako manual. ER or PR was considered positive if finding of more than 1% tumor cell nuclei are immunoreactive. Negative for ER

or PR if finding of less than 1% of tumor cell nuclei are immunoreactive [4] (Figure 1A and 1B)

HER-2/neu status was assessed by a score that includes the intensity and the percentage of positive

tumor cells. HER2 testing results fall into three categories; positive, equivocal and negative [5] [Table 1] (Figure: 2). For the purpose of this study equivocal results were considered negative.




Fig 1A: Showing ER positivity of score 8 in the above tumour.




Fig 1B: Showing PR positivity of score 7 in the above tumour. Inset shows HER2 negative staining

Table 1: HER2 Testing by Validated Immunohistochemistry Assay

Status	Score	Significance
Positive	3+	Uniform intense membrane staining of >30% of invasive tumor cells
Equivocal	2+	Complete membrane staining ,non-uniform or weak in intensity ,in at least 10% of the cells or intense complete membrane staining in 30% or less of tumor cells
Negative	1+	Weak or incomplete membrane staining in any proportion of tumor cells
Negative	0	No staining




Fig 2: Grade III IDC .Showing Strong HER 2 membranous staining of score3 (x Inset shows ER & PR negativity

Statistical analysis:

Statistical analysis was done using spss software version 17. The relationship between clinicopathological parameters, IHC subtypes and histopathological grading was done using chi-square test.

RESULTS:

There were 66 cases of microscopically confirmed invasive breast carcinoma in women. Most women (28 of 66; 42.42%) were between age group of 41 and 50 years. Eighteen (27.27%) were 40 years and younger at diagnosis [Table 2].

Table 2: Age at diagnosis of invasive breast carcinoma

Age in years	Frequency	Percentage
<40	18	27.27
41-50	28	42.42
<50	20	30.30

Most (47 of 66; 71.21%) were invasive ductal carcinoma, not otherwise specified (IDC NOS). [Figure 3, 4] There were other subtypes of invasive carcinoma, including (12 of 66; 18.18%) of medullary carcinoma ,

IDC with Mucinous Type (3 of 66 ; 4.54%),invasive lobular carcinoma(ILC) (2 of 66 ; 3.03%) intracystic papillary carcinoma (1 of 66; 1.51%) ,metaplastic carcinoma(1 of 66 ; 1.51%) .



Fig 3: IDC (NOS): Gross specimen




Fig-5: Grade II IDC showing ER positivity(x400) PR Positivity Of score 8




Fig 4: Grade I IDC (NOS) with tubular pattern




Fig-6: Grade I IDC showing nuclear of score 8 nuclear Inset Shows HER 2 Negativity (x400)

Table 3: Correlation Of Hormone Receptor Status With Histological Subtypes (Figure 5, 6)

Histologic Subtype	Frequency	Percentage (%)	ER		PR		HER-2	
			Pos	Neg	Pos	Neg	Pos	Neg
IDC (NOS)	47	71.21	31	16	29	18	14	33
Medullary carcinoma	12	18.18	00	12	00	12	00	12
Invasive lobular carcinoma	02	3.03	01	01	01	01	01	01
IDC with Intracystic Papillary Carcinoma	01	1.51	01	00	00	01	00	01
IDC with mucinous type	03	4.54	00	03	00	03	00	03
Metaplastic Carcinoma	01	1.51	00	01	00	01	00	01

Table 4: Correlation of ER, PR and HER2 status with clinicopathological data

	ER + n (%)	p value	PR+ n (%)	P value	HER2+ n (%)	P value
Age (years)						
< 50 (38)	20 (52.63)	0.80	18(47.37)	0.90	7(18.42)	0.49
>50 (28)	13 (46.43)		12(42.86)		8(28.57)	
Tumor size (Cms)						
<_5 (41)	20(48.78)	1	18(43.90)	0.94	11(26.82)	0.47
>5 (25)	13(52)		12(48)		4(16)	
Histologic type						
Infiltrating ductal (47)	31(65.96%)	0.72	29(61.70%)	0.64	14(29.79%)	0.87
Infiltrating lobular(2)	1(50%)		1(50%)		1(50%)	
Lymph node						
Positive (27)	17(62.96%)	0.52	15(55.56%)	0.83	5(18.52%)	0.87
Negative(11)	5(45.45%)		5(45.45%)		3(27.27%)	
Tumour grade						
Grade1-2 (41)	24(58.54%)	0.12	23(56.10%)	0.048	12(29.27%)	0.18
Grade 3 (25)	9(36%)		7(28.0%)		3(12%)	

Table 5: ER, PR and HER2/neu status in various types of breast carcinoma

	ER	PR	HER2/neu
Positive	33(50%)	30(45.45%)	15 (22.73%)
Negative	33(50%)	36(54.55%)	49 (74.24%)
Equivocal	-	-	2 (3.03%)
Total	66	66	66

Table 6: Immunohistochemical subtypes

ER/PR and HER2	No.of cases(n=66)	%
ER/PR + HER2 -	15	22.73
ER/PR + HER2 +	13	19.70
ER/PR - HER2 -	31	46.97
ER/PR - HER2+	2	3.03
ER+PR - HER2-	3	4.55
Er/PR +HER2- Equivocal	2	3.03

DISCUSSION

Neoplasms of the breast are one of the common lesions of the breast which though virtually limited to females can occur in males as rare exceptions. Breast cancer is a heterogeneous disease composed of growing number of recognized biological subtypes. Prognostic indicators based on currently available clinical and histopathologic variables such as tumor size, tumor grade, lymph node status and hormone receptor status already exist and are used to predict a patient's clinical outcome in certain situations [6]. It is well known that ER, PR and HER-2 represent

the most acceptable factors for predicting prognosis response or resistance to treatment and the potential use of newer drugs [7]. Assessment of ER/PR and HER2 in breast cancer is mandatory in clinical practice [8].

So we undertook this study to evaluate for estrogen and progesterone receptors and HER2/neu expression in various types of carcinoma breast and correlated with tumor type, histological grade and lymphnode status. Our study comprised of 66 cases of malignant tumours of breast tumors.

Table 7: Comparison of ER, PR and HER2/neu status in breast carcinoma with other studies

ER,PR and HER2 status	Authors		
	Ayadi L <i>et al.</i> ; [7] (n=155)%	Hung HJ <i>et al.</i> ; [9] (n=1362)%	Present Study (n=50)%
ER			
Positive	59.4	81.1	50
Negative	40.6	18.9	50
PR			
Positive	52.3	64.2	45.45
Negative	47.1	35.8	54.55
HER 2			
Positive	18.1	10.9	22.73
Negative	81.9	89.1	74.24
Equivocal	-	-	3.03

In our study, there were equal number of ER+ and ER- cases. PR + cases were less in number than PR- cases. Ayadi *et al.*; and Huang HJ *et al.*; Observed more number of ER and PR Positive cases. In the

present study, number of HER2+ (22.73%) and HER2- (74.24%) cases observed were in consistent with the observation made by Ayadi *et al.*; which showed 18.1% positively and 81.9% negatively. (Table-7).

Table 8: Immunohistochemical Subtypes.

Immunohistochemical Subtypes.	Satti MB <i>et al.</i> ; [8] (%)	Onitilo AA <i>et al.</i> ; [10] (%)	Hung HJ <i>et al.</i> ; [9] (%)	Present Study (%)
ER/PR+, HER 2-	53	68.9	66.4	22.73
ER/PR -, HER 2+	11	10.2	30.9	19.70
ER/PR -, HER 2-	24	13.4	13.8	46.97
ER/PR -, HER 2+	12	7.5	45.6	3.03
ER/PR+./HER 2-	-	-	-	4.55
ER/PR+, HER 2 equivocal	-	-	-	3.03

In our study, triple negative cases were more accounting for 46.97% when compared to other studies. Special variants like medullary carcinoma (18.18%), and Metaplastic carcinoma (1.51%) which showed triple negativity were included in present study.

Apart from these subtypes in present study 3 cases of ER+PR-, HER2- type, one was IDC with intracystic papillary type of histological grade 2 and 2 cases were IDC (NOS) of histological grade 3 and also 2 cases of ER+PR+ and HER2 equivocal, which included IDC (NOS) of grade II and grade III (Table-8).

Frequency of other subtypes observed was consistent with the observation made by Satti MB *et al.*;

Table 9: Correlation of ER status in various types of carcinoma breast with clinicopathological data

	Ayadi L <i>et al.</i> ;[7]		Present Study	
	ER + %	P Value	ER +%	P Value
Age (years)				
< 50	47.5	0.002	52.63	0.80
>50	72		46.43	
Tumour Size(cms)				
< 5	62.7	0.129	48.78	1
>5	48.5		52	
Histologic type				
Infiltrating Ductal	61.1	0.31	65.96	0.72
Infiltrating Lobular	50		50	
Lymph node				
Positive	60	0.88	62.96	0.52
Negative	58.9		45.45	
Tumour grade				
Grade-1-2	72.2	0.000	58.54	0.12
Grade 3	22.5		36	

In our study, ER + tumors were more in the age group of < 50 years. Whereas in the study done by Ayadi *et al.*, ER + tumours were more in the age group > 50 years and Satti MB *et al.*; had similar proportion of ER + cases above and below the age group of 50 years.

A negative correlation between ER expression and histological grade was noted. No association was found between ER expression and tumor size, histologic type and lymphnode involvement which were similar to the observation made by Ayadi L *et al.*; (Table-9).

Table 10: Correlation of PR status in various types of carcinoma breast with clinicopathological data

	Ayadi L <i>et al.</i> ; [7]		Present Study	
	PR +%	P Value	PR+ %	P Value
Age (years)				
< 50	53.8	0.76	47.37	0.90
>50	51.4		42.86	
Tumor Size (cms)				
< 5	53.4	0.72	43.9	0.94
>5	50		48	
Histologic type		0.47		0.64
Infiltrating ductal	53.8		61.7	
Infiltrating Lobular	45.8		50	
Other			0	
Lymph node				
Positive	54.7	0.66	55.56	0.83
Negative	51.1		45.45	
Tumor grade				
Grade 1-2	61.4	0.000	56.10	0.048
Grade 3	27.5		28	

In our study, PR + tumors were more in the age group of < 50 years which was similar to the observation made by Ayadi L *et al.*; A negative correlation between PR expression and histological

grade was noted. No association was found between PR expression and tumor size, histologic type and lymphnode involvement which were similar to the observation made by *et al.*; (Table-10).

Table 11: Correlation of HER2 status in various types of carcinoma breast with clinicopathological data Ayadi L *et al.*; 62 Huang HJ *et al.*;

	Ayadi L <i>et al.</i> ; [7]		Hunjg HJ <i>et al.</i> ; [9]		Present study	
	HER2+ n(%)	P value	HER2+ (%)	P Value	HER2 +	P Value
Age (years)						
< 50	21.3	0.28	12.1	0.344	18.42	0.49
>50	14.7		10.4		28.57	
Tumor Size (cms)						
< 5	15.3	0.104	10.5(<2)	0.617	26.82	0.47
>5	27		11.4(>2)		16	
Histologic type		0.33		-		0.87
Infiltrating ductal	16.8		-		29.79	
Infiltrating Lobular	25		-		50	
Lymph node						
Positive	36.9	0.000	12.2	0.255	18.52	0.87
Negative	4.4		10.2		27.27	
Tumor grade						
Grade 1-2	14.8	0.072	4.6	<0.001	29.27	0.180
Grade 3	27.5		20.8		12	

In our study, HER2/neu overexpression was seen in the age group of < 50 (18.42%) similar observation was made by Ayadi *et al.*; . There was no correlation between HER2 status and tumor size,

axillary lymph node, type and age at diagnosis in the present study and correlated with the study done by Huang HJ *et al.*; Ayadi *et al.*; observed significant

correlation between lymph node status and HER2 overexpression (Table-11).

Table 12: Comparison of Immunohistochemical subtypes with Histologic grading

Immunohistochemical Subtypes (%)	Onitilo AA <i>et al.</i> ; [10]				Present study				
	Histologic Grades				P Value	Grade I	Grade II	Grade III	P Value
	Grade I	Grade II	Grade III						
ER/PR, +HER2-	28.9	44.9	21.5	< 0.001	37.5	27.27	12	0.057	
ER/PR, +HER2+	6.0	41.4	49.1		0	30.3	12		
ER/PR, - HER2-	4.0	12.5	76.3		62.5	30.3	64		
ER/PR, -HER2+	1.2	20.0	77.7		0	6.06	0		

In our study Observed close correlation between tumor grade and immunohistochemical subtypes which was similar to the study done by Onitilo AA *et al.*; HER2+ and basal like subtypes were associated with poorly differentiated tumors (p=0.003) (Table-12).

Correlation between ER and PR status:

In our study positive correlation observed between ER and PR (p = 0.000) which was similar to the study done by Ayadi L *et al.*; [7] and Eisenberg ALA *et al.*; [11]. According to Rosen, most PR positive tumors are also positive for ER [11].

Correlation between HER2 overexpression and hormonal receptors status:

In the study done by Ayadi L *et al.*; [7] the percentage of HER-2 over-expression was sharply weaker in ER positive tumors: 7.6%, compared with 33.3% for ER negative tumors (p:0.000). Over-expression of HER-2 was also inversely related to PR status (p: 0.048).

In our study inverse relationship between HER2/neu and ER/PR expression was observed, similar observation was made by Ambroise *et al.*; [12] and Eisenberg ALA *et al.*; [11]. The inverse association between HER-2/neu and hormone receptors to lower or absent hormone receptors in women with HER-2/neu positive cancers. This is one of the reasons why women who overexpress HER-may be resistant to tamoxifen [13].

CONCLUSION

Our study highlights the importance of histopathological examination in breast lumps but also in predicting the prognosis by typing, staging and grading malignant neoplasms of breast. The interrelationship between ER, PR and HER-2/neu has an important role in the management of breast cancer. Endocrine therapy (tamoxifen) is recommended for tumors expressing ER/PR. Patients with breast carcinoma overexpressing HER-2 do not respond to tamoxifen therapy. Recently anti-HER-2 antibodies (Herceptin) have been shown to be effective against HER-2 overexpressing breast carcinomas. In our study

an attempt was made to understand the correlation of ER, PR & HER-2 status with histopathological grading and clinicopathological parameters.

In conclusion, ER, PR and HER-2 status correlates well with histopathological grading. Higher the tumor grade, the more likely that ductal carcinoma will be Her2 + and ER/PR negative or triple negative. Hence, our study support IHC classification as a clinical tool as ER/PR and HER 2 testing is widely available at a reasonable cost, is a clinically-used, therapeutically informative classification of breast cancer based on immunophenotype / biologic phenotypes, and is prognostic as well as somewhat predictive. Follow up study of these patients is needed to assess the prognostic significance.

REFERENCES

1. Maki DD, Grossman RI. Patterns of disease spread in metastatic breast carcinoma: influence of estrogen and progesterone receptor status. American journal of neuroradiology. 2000 Jun 1; 21(6):1064-6.
2. Gluz O, Liedtke C, Gottschalk N, Pusztai L, Nitz U, Harbeck N. Triple-negative breast cancer—current status and future directions. Annals of Oncology. 2009:mdp492.
3. Helin HJ, Helle MJ, Kallioniemi OP, Isola JJ. Immunohistochemical determination of estrogen and progesterone receptors in human breast carcinoma. Correlation with histopathology and DNA flow cytometry. Cancer. 1989 May 1; 63(9):1761-7.
4. Hammond ME, Hayes DF, Dowsett M, Allred DC, Hagerty KL, Badve S, Fitzgibbons PL, Francis G, Goldstein NS, Hayes M, Hicks DG. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer (unabridged version). Archives of pathology & laboratory medicine. 2010 Jul; 134(7):e48-72.
5. Gutierrez C, Schiff R. HER2 Biology, detection, and clinical implications. Arch Pathol Lab Med 2011; 135:55-62.

6. Munjal K, Ambaye A, Evans MF, Mitchell J, Nandedkar S, Cooper K. Immunohistochemical analysis of ER, PR, Her2 and CK5/6 in infiltrative breast carcinomas in Indian patients. *Asian Pac J Cancer Prev.* 2009 Jan 1; 10(5):773-8.
7. Ayadi L, Khabir A, Amouri H, Karray S, Dammak A, Guermazi M, Boudawara T. Correlation of HER-2 over-expression with clinico-pathological parameters in Tunisian breast carcinoma. *World journal of surgical oncology.* 2008 Oct 22; 6(1):1.
8. Satti MB. Oestrogen receptor/progesterone receptor and human epidermal growth factor receptor 2 status in breast cancer: a 9-year study at Princess Noorah Oncology Center, Saudi Arabia. *Histopathology.* 2011 Sep 1; 59(3):537-42.
9. Kiag DT, Kennedy B.J. Factors effecting estrogen receptors in breast cancer. *Cancer* 1977; 40:1571-6.
10. Onitilo AA, Engel JM, Greenlee RT, Mukesh BN. Breast cancer subtypes based on ER/PR and Her2 expression: comparison of clinicopathologic features and survival. *Clinical medicine & research.* 2009 Jun 1; 7(1-2):4-13.
11. Wynveen CA, Nehhozina T, Akram M, Hassan M, Norton L, Van Zee KJ, Brogi E. Intracystic papillary carcinoma of the breast: an in situ or invasive tumor? Results of immunohistochemical analysis and clinical follow-up. *The American journal of surgical pathology.* 2011 Jan 1; 35(1):1-4.
12. Ambrose M, Ghosh M, Mallikarjuna VS, Kurian A. Immunohistochemical profile of breast cancer patients at a tertiary care hospital in South India. *Asian Pac J Cancer Prev.* 2011 Jan 1; 12(3):625-9.
13. Huang HJ, Neven P, Drijckoningen M, Paridaens R, Wildiers H, Van Limbergen E, Berteloot P, Amant F, Vergote I, Christiaens MR. Association between tumour characteristics and HER-2/neu by immunohistochemistry in 1362 women with primary operable breast cancer. *Journal of clinical pathology.* 2005 Jun 1; 58(6):611-6.