

Original Research Article

## Evaluation of Underlying Factors of Back Pain in Patients with Depression and Psychosis

Halil İbrahim SERİN<sup>1</sup>, Ali İrfan GÜL<sup>2</sup>, İskender Samet DALTABAN<sup>3\*</sup>, Lütfiye TUTKUN<sup>4</sup>, Hakan AK<sup>3</sup>, Ayşe Yeşim Göçmen<sup>4</sup>

<sup>1</sup>Bozok University, School of Medicine, Department of Radiology, Yozgat/TURKEY

<sup>2</sup>Bozok University, School of Medicine, Department of Psychiatry, Yozgat/TURKEY

<sup>3</sup>Bozok University, School of Medicine, Department of Neurosurgery, Yozgat/TURKEY

<sup>4</sup>Bozok University, School of Medicine, Department of Biochemistry, Yozgat/TURKEY

### \*Corresponding author

İskender Samet DALTABAN

Email: [isamet79@hotmail.com](mailto:isamet79@hotmail.com)

**Abstract:** Back pain is a common complaint in depression, anxiety and somatoform disorders. In this prospective study, we aimed to evaluate the underlying factors for back pain in patients with depression and psychosis. 129 patients aged ≥18 years with anxiety disorder and somatoform disorder, including 78 patients with back pain and 51 controls without back pain who presented to our clinic. Visual Analog Score (VAS), tender point count, blood levels of vitamin D, calcium, phosphor, magnesium, and thyroid stimulating hormone (TSH), and thoracolumbar magnetic resonance imaging (MRI) were performed in all subjects to evaluate the possible underlying causes of pain. There were significant differences in weight, VAS score, tender point count, Schmorl's node, Modic's degeneration and blood levels of vitamin D, Ca, P, Mg and TSH between study group and controls. We recommend to perform thoracolumbar MR imaging and to evaluate blood chemistry for vitamin D deficiency and other osteomalacia risk factors in psychiatric patients presenting with back pain and to revalue treatment based on results obtained.

**Keywords:** Low-back pain, depression, psychosis, magnetic resonance imaging, Modic degeneration, endplate degeneration.

### INTRODUCTION

Low-back pain is very common complaint in general population. Its global point prevalence is 9.4% [1]. Although performing many studies about this subject, only a few large representative multinational studies have considered the mental health status of people with back pain [2]. Depression and psychosis is a common and steadily increasing pathology around the world [1, 2].

Osteomalacia is a metabolic bone disease characterized by insufficient or delayed osteoid bone mineralization and lower rate of bone formation, which causes back pain [3]. Vitamin D deficiency is observed to be most common cause of osteomalacia. In psychiatric patients, causes of osteomalacia include decreased vitamin D production, impaired vitamin D absorption from intestine, excessive degradation of vitamin D or 25(OH) vitamin D and phosphate losses [5]. Thyroid hormones are essential for normal skeletal growth and maintenance of bone mass. Hypothyroidism

leads growth retardation and disruption in bone formation whereas hyperthyroidism leads reduction in bone mass, advanced bone age, accelerated growth and increased risk for osteoporotic fracture [6]. Osteomalacia can develop in hypocalcemia, primary hyperparathyroidism and hypercalcemia resulting from malignant disorders [5]. There is a negative feedback mechanism between serum PTH and Mg levels. Moreover, the relationship between bone metabolism and androgens, estrogens or both have been showed in different studies [7, 8].

Endplate abnormalities on lumbar spinal MR imaging were first classified by Modic. Primarily, 3 types of endplate Modic abnormalities can be characterized by T1- and T2-weighted images. Many studies showed that there is a strong association between Modic endplate changes and back pain [9]. Several epidemiological studies showed that incidence of back pain were higher in patients with Schmorl nodules when compared to normal population [10].

Herein, we aimed to investigate the underlying causes of back pain in patients with depression and psychosis.

## MATERIALS AND METHODS

After approval of local ethics committee of Bozok University (date: 08.04.2014; number: 08/33), patients were divided into two groups. Group I was included the patients having depression and/or psychosis and back pain. However, group II included the patients having depression and/or psychosis without back pain. The weight, height, age and other demographic data of the patients were recorded. Patients younger 18 years of age were excluded from the study. Also, patients with diabetes mellitus, hypertension, hypo/ hyperthyroidism, hypo/ hyperparathyroidism, chronic liver and kidney diseases were excluded. Heavy smokers and patients with heavy alcohol consumption were also excluded from the study. The patients with conditions that can be cause of back pain such as congenital anomaly, previous history of trauma, osteoarthritic degeneration, cardiovascular disorders and drug use were excluded.

In all patients, complete blood count, AST, ALT, creatinine, TSH, estradiol, testosterone, vitamin D, Ca, P and Mg levels were measured. VAS scores and body mass index (BMI) and tender point counts were recorded. The pain was assessed by using visual analog scale (VAS). VAS is a linear scale with a continuous line between 2 end points: 0, no pain and 10, worst possible pain. The patients were asked to mark mean severity of pain within prior week. Tender point count was performed in back region at 17 vertebral levels (12 thoracic and 5 lumbar). The tender points can be best assessed by applying 4 kg pressure with thumb which is the pressure that allow fading of nail bed when thumb was compressed on a though layer.

In all patients, thoracolumbar sagittal MRI was obtained to assess Schmorl nodules and endplate degeneration, and other pathologies. T1- and T2-weighted thoracolumbar sagittal MRI sequences were performed by using 1.5 Tesla Ingenia MR Device (Philips Medical Systems, Tilburg, Netherlands). The MR imaging parameters used were as follows: FOV, 180; TR, 2300-2600 and section thickness, 4-mm. All images were assessed by a single radiologist to avoid subjective assessment.

## Biochemical analysis

All the blood samples were centrifuged for 10 min at 3000 RPM, and the serum samples were frozen at  $-80^{\circ}\text{C}$  until the assays were performed by an

investigator who was blind to each patient's status. Commercial enzyme-linked immunosorbent assay (ELISA) kits were used for the measurement of human Vitamin C levels (Elabscience, Wuhan, PRC) using appropriate wavelengths on a micro-plate reader (Biotech Instruments, EL $\times$ 800 TM, Winooski, USA) following the assay instructions. Serum levels of routine biochemistry parameters were measured using an Architect ci2800 integrated system (Abbott Diagnostics, Abbott Park, IL).

## STATISTICAL ANALYSIS

Statistical analysis was performed using SPSS package program (Version 18.0 for Windows, SPSS Inc., Chicago, IL, USA). An analysis of normality of the continuous variables was performed with the Kolmogorov-Smirnov test for multivariate analysis; only variables with a P-value  $< 0.05$  were entered into the model and selected using a stepwise selection procedure.

Continuous variables were expressed as mean  $\pm$  standard deviation (SD) or median while categorical variables were expressed as a percentage. A p value  $< 0.05$  was considered statistically significant. The  $\chi^2$ -test was used to compare proportions. Continuous variables were compared using an independent-groups Student's t-test if normality assumptions were met; otherwise, groups were compared using the Wilcoxon rank-sum test. Pearson correlation analysis was performed.

## RESULTS

The study population included 129 patients (aged  $\geq 18$  years  $\leq 65$ ). Of these, 78 patients consisting study group had back pain while 51 patients consisting control group had no back pain. Mean age was  $45.95 \pm 14.78$  years (range: 18-65 years) in the study group whereas  $41.86 \pm 13.57$  years (range: 20-65 years) in the control group.

There were significant differences in weight ( $p=0.011$ ), VAS score ( $p=0.001$ ), tender point count ( $p=0.001$ ), Schmorl nodule ( $p=0.001$ ), Modic degeneration ( $p=0.000$ ), levels of vitamin D ( $p=0.001$ ), Ca ( $p=0.017$ ), P ( $p=0.028$ ), Mg ( $p=0.001$ ) and TSH ( $p=0.001$ ) in between two groups. However, no significant difference was found in height ( $p=0.592$ ), age ( $p=0.116$ ), levels of estradiol ( $p=0.154$ ), testosterone ( $p=0.387$ ), ALT ( $p=0.378$ ), AST ( $p=0.338$ ), creatinine ( $p=0.257$ ), PTH ( $p=0.348$ ), vitamin C ( $p=0.234$ ), hemoglobin ( $p=0.995$ ), WBC ( $p=0.834$ ), platelet count ( $p=0.854$ ) and erythrocyte count ( $p=0.690$ ) (Table 1).

**Table 1: Showing the data of the patients.**

Parameter	Control (n=51)	Back pain (n=78)	P value
Height	167.80 ± 5.86	167.19 ± 6.95	0.592
Weight	82.49 ± 8.14	85.97 ± 6.18	<b>0.011</b>
Age(Years)	41.86 ± 13.57	45.95 ± 14.78	0.116
VAS score	0.04 ± 0.28	3.97 ± 1.98	<b>0.001</b>
Tender Point Counts	0.04 ± 0.28	5.45 ± 2.17	<b>0.001</b>
Schmorl nodule	0.37 ± 0.63	3.65 ± 1.51	<b>0.001</b>
Modic degeneration	0.51 ± 0.80	4.65 ± 2.14	<b>0.001</b>
Vit. D (ng/ml)	22.62 ± 9.26	8.07 ± 2.31	<b>0.001</b>
Estradiol (pg/ml)	69.32 ± 80.01	52.07 ± 56.40	0.154
Testosterone (ng/ml)	2.00 ± 3.77	1.50 ± 2.01	0.387
Ca (mg/dl)	10.07 ± 2.58	9.09 ± 1.60	<b>0.017</b>
P (mg/dl)	4.19 ± 2.86	3.14 ± 2.10	0.028
Mg (mg/dl)	1.89 ± 0.52	1.58 ± 0.39	<b>0.001</b>
ALT (U/L)	13.22 ± 12.78	11.29 ± 11.58	0.378
AST (U/L)	22.14 ± 13.18	20.10 ± 10.72	0.338
TSH (U/ml)	3.28 ± 2.25	1.23 ± 0.56	<b>0.001</b>
Parathyroid Hormone (pg/ml)	22.78 ± 13.19	20.79 ± 10.75	0.348

## DISCUSSION

In the present study, we aimed to evaluate whether there is a correlation between MR imaging findings and cause of back pain in patients presented with depression and psychosis. We assessed biochemical values and thoracolumbar MR imaging results in patients with back pain in order to investigate the underlying causes of pain in these patients. In our study, biochemical results and thoracolumbar MR imaging findings compatible with osteomalacia were detected in patients with back pain. There were significant differences in weight, VAS score, tender point count, Schmorl nodule, Modic degeneration and vitamin D, Ca, P, Mg and TSH values between patient and control group. The patients with back pain had higher body weight, VAS score, tender point count, number of Schmorl nodule and endplate degeneration while lower vitamin D, Ca, P, Mg and TSH values.

There is limited number of studies on this topic. However, several authors have long been suggested that back pain is commonly presented in psychotic and depressive patients. Back pain can be related to various factors in patients with psychiatric disorders. In many studies, an association was shown between psychiatric disorders and neck or back pain [11]. In an epidemiology study in United Kingdom, it was found that the incidence is 16.9% for association of chronic pain and psychiatric diagnosis [12]. In a study from United States, back pain was detected in 23% of the patients with psychiatric disorder [13].

There is a strong relationship between depression and neck/back pain in developed and developing countries in the studies [14-19]. Jain

reported that back pain could be associated with mood and anxiety symptoms [20]. It was reported that back pain is common in major depression in addition to pain at other body regions [21, 22].

Okamoto *et al.*; reported that psychiatric disorders and decreased bone mineral density (BMD) were encountered in patients with mild hypercalcemia related to primary hyperparathyroidism [23]. Joborn *et al.*; reported that anxiety, depression and cognitive complaints are common in patients with primary hyperparathyroidism and mild hypercalcemia [24]. Driesen *et al.*; found a relationship between secondary hyperparathyroidism and depressive symptoms or cognitive dysfunction [25].

Many studies indicate a relationship between back pain and stress and anxiety [26-28]. It has been reported that psychological distress can be both reason and consequence of neck and back pain [29]. Incidence of depression is 5-8% in general population while it varies from 22% to 78% in patients with chronic pain [30]. However, the rate was found to be 5-50% in studies using strict criteria and techniques of structured interview [31]. Pain prevalence was found to be 65% in patients with depression [32]. In a study on 11 patients (mean age: 55 years) with hypoparathyroidism for more than 9 years Kowdley *et al.*; found cognitive dysfunction in 65%, motor dysfunction in 45% and intracranial calcification in 60% of the patients [33].

More than two decades, efforts to develop a psychological model of multi-disciplinary approach has gained acceptance for better understanding of pain. In recent years, psychological factors have become focus

of interest in pain studies and pain therapy [34]. In back pain, psychological factors are generally more significant than biomedical or biomechanical factors [35]. In a previous study, it was observed that severe depression, anxiety, somatization, avoidance behavior and coping capacity differed between individuals with or without pain. In consistent with previous studies, these results showed significance of psychological variables as risk factor for back and neck pain. It has been reported that chronic back pain is more frequently seen in individuals who are unable to cope with helplessness and hopelessness [36].

#### CONCLUSION:

In conclusion, we found a significant relationship between abnormal values that may cause osteomalacia and endplate degenerations and Schmorl nodule formation in the patients with back pain who were diagnosed as depression and/or psychosis. We recommend to perform thoracolumbar MR imaging and to evaluate blood chemistry for vitamin D deficiency and other osteomalacia risk factors in these patients.

#### ACKNOWLEDGMENT:

The support for this study was provided by the Bozok University Scientific Research Project Funding Unit thanks were extended of it as well.

#### REFERENCES

1. Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, Williams G, Smith E, Vos T, Barendregt J, Murray C. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Annals of the rheumatic diseases*. 2014 Mar 24; annrheumdis-2013.
2. Stubbs B, Koyanagi A, Thompson T, Veronese N, Carvalho AF, Solomi M, Mugisha J, Schofield P, Cosco T, Wilson N, Vancampfort D. The epidemiology of back pain and its relationship with depression, psychosis, anxiety, sleep disturbances, and stress sensitivity: Data from 43 low-and middle-income countries. *General Hospital Psychiatry*. 2016 Dec 31; 43:63-70.
3. Pinheiro MB, Ferreira ML, Refshauge K, Ordoñana JR, Machado GC, Prado LR, Maher CG, Ferreira PH. Symptoms of Depression and Risk of New Episodes of Low Back Pain: A Systematic Review and Meta-Analysis. *Arthritis care & research*. 2015 Nov 1; 67(11):1591-603.
4. Bozkırlı ED, Bozkırlı E, Pourbagher A, Ulaş M, Yücel AE. Osteomalaziye İkincil Gelişen Sakroiliak Eklem Tutulumlu Bir Olgu Sunumu. *Ankara Medical Journal*. 2014.
5. Francis RM, Selby PL. Osteomalacia. *Baillieres Clin Endocrinol Metab*. 1997; 11:145-163.
6. Galliford TM, Murphy E, Williams AJ, Bassett JH, Williams GR. Effects of thyroid status on bone metabolism: a primary role for thyroid stimulating hormone or thyroid hormone?. *Minerva endocrinologica*. 2005 Dec; 30(4):237-46.
7. Kanis JA, Melton LJ, Christiansen C, Johnston CC, Khaltsev N. The diagnosis of osteoporosis. *Journal of bone and mineral research*. 1994 Aug 1; 9(8):1137-41.
8. Leder BZ, LeBlanc KM, Schoenfeld DA, Eastell R, Finkelstein JS. Differential effects of androgens and estrogens on bone turnover in normal men. *The Journal of Clinical Endocrinology & Metabolism*. 2003 Jan 1; 88(1):204-10.
9. Albert HB, Kjaer P, Jensen TS, Sorensen JS, Bendix T, Manniche C. Modic changes, possible causes and relation to low back pain. *Medical hypotheses*. 2008 Dec 31; 70(2):361-8.
10. Takatalo J, Karppinen J, Niinimäki J, Taimela S, Mutanen P, Sequeiros RB, Näyhä S, Järvelin MR, Kyllönen E, Tervonen O. Association of modic changes, Schmorl's nodes, spondylolytic defects, high-intensity zone lesions, disc herniations, and radial tears with low back symptom severity among young Finnish adults. *Spine*. 2012 Jun 15; 37(14):1231-9.
11. Croft PR, Papageorgiou AC, Ferry S, Thomas E, Jayson MI, Silman AJ. Psychologic distress and low back pain: evidence from a prospective study in the general population. *Spine*. 1995 Dec 15; 20(24):2731-7.
12. Benjamin S, Morris S, McBeth JO, Macfarlane GJ, Silman AJ. The association between chronic widespread pain and mental disorder: a population-based study. *Arthritis & Rheumatism*. 2000 Mar 1; 43(3):561.
13. Tschudi-Madsen H, Kjeldsberg M, Natvig B, Ihlebaek C, Dalen I, Straand J, Bruusgaard D. Multiple symptoms and medically unexplained symptoms—closely related concepts in general practitioners' evaluations. A linked doctor–patient study. *Journal of psychosomatic research*. 2013 Mar 31; 74(3):186-90.
14. Chen X, Cheng HG, Huang Y, Liu Z, Luo X. Depression symptoms and chronic pain in the community population in Beijing, China. *Psychiatry Res*. 2012; 200:313-317.
15. Carroll LJ, Cassidy JD, Côté P. Depression as a risk factor for onset of an episode of troublesome neck and low back pain. *Pain*. 2004 Jan 31; 107(1):134-9.
16. Currie SR, Wang J. Chronic back pain and major depression in the general Canadian population. *Pain*. 2004; 107:54-60.
17. Demyttenaere K, Bruffaerts R, Lee S, Posada-Villa J, Kovess V, Angermeyer MC, Levinson D, de Girolamo G, Nakane H, Mneimneh Z, Lara C. Mental disorders among persons with chronic back or neck pain: results from the World Mental Health Surveys. *Pain*. 2007 Jun 30; 129(3):332-42.

18. Trief PM, Elliott DJ, Stein N, Frederickson BE. Functional vs. organic pain: a meaningful distinction? *Journal of clinical psychology*. 1987 Mar 1; 43(2):219-26.
19. Von Korff M, Miglioretti DL. A prognostic approach to defining chronic pain. *Pain*. 2005 Oct 31; 117(3):304-13.
20. Jain R. Pain and the brain: lower back pain. *The Journal of clinical psychiatry*. 2009 Feb 16; 70(2):41-.
21. Trivedi MH. The link between depression and physical symptoms. *Prim Care Companion J Clin Psychiatry*. 2004; 6(Suppl 1):12-6.
22. Baune BT, Caniato RN, Garcia-Alcaraz MA, Berger K. Combined effects of major depression, pain and somatic disorders on general functioning in the general adult population. *Pain*. 2008 Aug 31; 138(2):310-7.
23. Okamoto T, Gerstein HC, Obara T. Psychiatric symptoms, bone density and non-specific symptoms in patients with mild hypercalcemia due to primary hyperparathyroidism: a systematic overview of the literature. *Endocrine journal*. 1997; 44(3):367-74.
24. Jobom C, Hetta J, Lind L, Rastad J. Self-rated psychiatric symptoms in patients operated on because of primary hiperparathyroidism and in patients with long-standing mild hypercalcemia.
25. Driessen M, Wetterling T, Wedel T, Preuss R. Secondary hyperparathyroidism and depression in chronic renal failure. *Nephron*. 1995 Jul 1; 70(3):334-9.
26. Leino P, Magni G. Depressive and distress symptoms as predictors of low back pain, neck-shoulder pain, and other musculoskeletal morbidity: a 10-year follow-up of metal industry employees. *Pain*. 1993 Apr 1; 53(1):89-94.
27. Burton AK, Tillotson KM, Main CJ, Hollis S. Psychosocial predictors of outcome in acute and subchronic low back trouble. *Spine*. 1995 Mar 15; 20(6):722-8.
28. Gatchel RJ, Polatin PB, Kinney RK. Predicting outcome of chronic back pain using clinical predictors of psychopathology: a prospective analysis. *Health psychology*. 1995 Sep; 14(5):415.
29. Jensen JC, Haahr JP, Frost P, Andersen JH. The significance of health anxiety and somatization in care-seeking for back and upper extremity pain. *Family practice*. 2012 Feb 1; 29(1):86-95.
30. Haythornthwaite JA, Sieber WJ, Kerns RD. Depression and the chronic pain experience. *Pain*. 1991 Aug 31; 46(2):177-84.
31. Aslan S, Nazlıel B. Gerilim tipi baş ağrısında anksiyete, depresyon düzeyleri ve tanısal değerlendirme. In *Yeni Symposium 2002* (Vol. 40, No. 1, pp. 10-4).
32. Greist JH, Jefferson JW, Trivedi MH. Depression and pain. *J Clin Psychiatry*. 2008; 69:1970-1978.
33. Kowdley KV, Coull BM, Orwoll ES. Cognitive impairment and intracranial calcification in chronic hypoparathyroidism. *The American journal of the medical sciences*. 1999 May 1; 317(5):273.
34. Waddell G. Psychosocial analysis of low back pain. *Baillieres Clin Rheumatol*. 1992; 6:523-557.
35. Linton SJ. A review of psychological risk factors in back and neck pain. *Spine-Philadelphia-Harper And Row Publishers Then Jb Lippincott Company*. 2000 May 1; 25(9):1148-56.
36. Eriksen HR, Ursin H. Subjective health complaints, sensitization, and sustained cognitive activation (stress). *Journal of psychosomatic research*. 2004 Apr 30; 56(4):445-8.