

Original Research Article

## Marine Fouling Diatoms Identified in the Vellar Estuary of Porto Nova, Cuddalore District, Tamil Nadu, India

E. Suresh and P. Sureshkumar

CAS in Marine Biology, Faculty of Marine Sciences, Annamalai University, Parangipettai, Cuddalore District, India

### \*Corresponding author

P. Sureshkumar

Email: [savegreenenvironment@gmail.com](mailto:savegreenenvironment@gmail.com)

**Abstract:** Marine fouling is an undesirable deposition of abiotic and biotic materials on manmade engineering structures in the marine environment especially contact with sea water. All forms of fouling are detrimental to the marine structures. After molecular fouling habitually the biofouling starts with microfoulers such as bacteria, micro algae, etc., then macro foulers such as algae, plants, mollusk, barnacles, etc will be followed. This study reports the diversity of diatoms involving in the fouling processes in the Vellar estuary of Parangipettai, south east coast of Indian subcontinent using different test panels. Seven species of diatoms such as *Skeletonema costatum*, *Triceratium reticulatum*, *Rhizosolenia setigera*, *Pleurosigma elongatum*, *Nitzschia longissima*, *Thalassione manitzschoides*, and *Thalassiothrix frauenfeldii* were recorded during this study.

**Keywords:** Biofouling, Vellar estuary, Porto Nova, Microfouler, Diatoms.

### INTRODUCTION

Marine fouling is an undesirable deposit of abiotic and biotic materials on natural and artificial substrates in the marine environment especially contact with sea water [1, 2]. The attachment of abiotic materials such as organic and inorganic molecules on immersed surfaces called molecular fouling [3] the attachments of biotic materials are called biofouling [4]. All form of accumulations is most deleterious to marine industries and causes massive structural and economic losses [5-7].

There are more than 250 genera of existing diatoms with possibly 2,00,000 species ranging across three orders of magnitude in the size of 5 µm to 2mm. This rich diversity shows the ability of their adaptation to various environmental conditions. They may be either as free floating or sessile communities [8]. The sessile diatoms are playing vital role in the ecology of fouling communities [9]. However, there were very few work only found on this aspects. Hence, this current study an attempt was made to identify the diatoms which fouled on the test panels of the study area of Vellar estuary.

### MATERIALS AND METHODS

#### Study area

The study area Vellar estuary is located exactly in the Lat. 11° 29' N; 79° 49' E at Parangipettai, south east coast of Indian subcontinent. The origin of Vellar

river on Shervarayan Hills from Salem and travels about 480 kms distance and finally drains into Bay of Bengal at Parangipettai thus forming the Vellar estuary. The average depth of this estuary is 2.5m and the maximum depth noticed during spring tide is 3.75m. The tidal amplitude is about one meter and its influence extends to a distance of 14km up to Bhuvanagiri. The maximum width of the estuary on its mouth is 200m. The Vellar estuary is divided into Marine zone, Gradient zone, Tidal zone, and fresh water zone. The location of present study zone gradiently falls opposite to CAS in Marine Biology institution (Faculty of Marine Sciences) of Annamalai University which is 1.6 km upstream from the mouth of estuary. It is one of the important fish landing centers of this district.

#### Construction, processing and Deployment of panels

Triplicates of the pieces (10 X 15 X 1.5 cm) of *T. paniculata* were procured from Pandian Depot, Tindivanam and aluminum pieces (15 X 20 X 1.5 cm) were procured from Dock Engineers OT, Cuddalore, Tamil Nadu. The pieces were fixed firmly with nylon rope of about 10 m and used them as a test panel. Both the panels were sterilized by 10% HCl, rinsed through with distilled water, dried in hot air oven at 40°C for 24hrs and kept in a dry place. Triplicates of the freshly processed panels were immersed about 1 meter depth in surface waters (~1 m) of the Vellar Estuary at low tide mark.

### Isolation of diatoms

The diatom samples were also collected from the experimental panels on 5th and 7th days of experiment with sterile swabs and then the samples were stored in 5% formalin solution. The isolated diatoms from the panels were identified using the keys given by Sournia [10].

### RESULT

In the present study an attempt was made to find out the diversity of diatoms fouled on all the four experimental panels during seven days of immersion period. The diatoms recorded during the course of experiment were grouped. The overall results indicated a heterogeneous occurrence of diatom on the experimental panel. Seven species of diatoms such as *Skeletonema costatum*, *Triceratium reticulatum*, *Rhizosolenia setigera*, *Pleurosigma elongatum*, *Nitzschia longissima*, *Thalassione manitzschoides*, and *Thalassiothrix frauenfeldii* were identified in both the wooden and aluminum panels.



Fig-1: *Skeletonema costatum*




Fig-2: *Triceratium reticulatum*



Fig-3: *Rhizosolenia setigera*



Fig-4: *Pleurosigma elongatum*



Fig-5: *Nitzschia longissima*



Fig-6: *Thalassione manitzschoides*




Fig-7: *Thalassiothrix frauenfeldii*

### DISCUSSION

Any surfaces immersed in marine environment were rapidly covered by various nonliving and living molecules forming a layer. The initial coats were generally abiotics and the bacteria as well as micro algae. Bacteria and Diatoms secrete diversified structural components called exopolymeric substances (EPS) in which cells are embedded during biofilm development [11-15]. Diatoms were the eukaryotic colonizer during this process [16]. These micro algal communities play a vital role in the development of the subsequent macrofouling community [17-20]. In marine industry, biofilms can have a detrimental impact

on account of the undesirable effect of corrosion promoted by microbial cell accumulation at interfaces [21]. However, there were only few studies only have focused on the fouling diatoms in the oceanic system [22-24].

In this study a preliminary step was taken to record the diversity of initial fouling diatom communities for a short duration of seven days in the Vellar estuary, Parangipettai, Southeast coast of India using wooden panels of *Terminalia paniculata* a chiefly used material for the marine vessel constructions and aluminum panels. The result showed that there were *Skeletonema costatum*, *Triceratium reticulatum*, *Rhizosolenia setigera*, *Pleurosigma elongatum*, *Nitzschia longissima*, *Thalassione nitzschioides*, and *Thalassiothrix frauenfeldii* initially formed on these panels. Exopolymeric secretions and sessile properties were well established in the *S. costatum*, *T. reticulatum*, *R. setigera*, *P. elongatum*, *N. longissima*, *T. nitzschioides*, and *T. frauenfeldii* and confirmed their sessile nature [25-30].

## CONCLUSION

This current study further concludes that the fouling diatoms's identified at Vellar estuary needed to be studied their ecological role and impact on the fouling processes to establish the control of economic losses owing to biofouling crisis successfully and use them beneficially for mankind.

## REFERENCES

1. Clare AS. Marine natural product antifoulants: status and potential. *Biofouling*. 1996 Apr 1;9(3):211-29.
2. Andrewartha J, Perkins K, Sargison J, Osborn J, Walker G, Henderson A, Hallegraef G. Drag force and surface roughness measurements on freshwater biofouled surfaces. *Biofouling*. 2010 May 1;26(4):487-96.
3. Amy G. Fundamental understanding of organic matter fouling of membranes. *Desalination*. 2008 Oct 31;231(1):44-51.
4. Hellio, C. and D. M. Y. Yebra,. Introduction. In *Advances in Marine Antifouling Coatings and Technologies*; Hellio C. and D. M. Y. Yebra, (Eds) Woodshead Publishing: Cambridge, UK. 2009; 1-15pp.
5. Dexter SC, Sullivan JD, Williams JI, Watson SW. Influence of substrate wettability on the attachment of marine bacteria to various surfaces. *Applied microbiology*. 1975 Aug 1;30(2):298-308.
6. Dexter SC. Influence of substratum critical surface tension on bacterial adhesion—in situ studies. *Journal of Colloid and Interface Science*. 1979 Jun 15;70(2):346-54.
7. Bakker DP, Klijnstra JW, Busscher HJ, van der Mei HC. The effect of dissolved organic carbon on bacterial adhesion to conditioning films adsorbed on glass from natural seawater collected during different seasons. *Biofouling*. 2003 Dec 1;19(6):391-7.
8. Mann DG. The species concept in diatoms. *Phycologia*. 1999 Nov;38(6):437-95.
9. Richmond MD, Seed R. A review of marine macrofouling communities with special reference to animal fouling. *Biofouling*. 1991 Apr 1;3(2):151-68.
10. Sournia, A. *Phytoplankton manual*. Monograph on oceanographic methodology 6. UNESCO, Paris, 1978;1-337.
11. Hoagland KD, Rosowski JR, Gretz MR, Roemer SC. Diatom extracellular polymeric substances: function, fine structure, chemistry, and physiology. *Journal of Phycology*. 1993 Oct 1;29(5):537-66.
12. Wustman BA, Gretz MR, Hoagland KD. Extracellular matrix assembly in diatoms (Bacillariophyceae)(I. A model of adhesives based on chemical characterization and localization of polysaccharides from the marine diatom *Achnanthes longipes* and other diatoms). *Plant Physiology*. 1997 Apr 1;113(4):1059-69.
13. Wetherbee R, Lind JL, Burke J, Quatrano RS. Minireview—The first kiss: establishment and control of initial adhesion by raphid diatoms. *Journal of Phycology*. 1998 Feb 1;34(1):9-15.
14. Underwood GJ, Paterson DM. The importance of extracellular carbohydrate production by marine epipelagic diatoms. *Advances in botanical research*. 2003 Dec 31;40:183-240.
15. Marvasi M, Visscher PT, Martinez LC. Exopolymeric substances (EPS) from *Bacillus subtilis*: polymers and genes encoding their synthesis. *FEMS microbiology letters*. 2010 Dec 1;313(1):1-9.
16. Yang C, Wang J, Yu Y, Liu S, Xia C. Seasonal variations in fouling diatom communities on the Yantai coast. *Chinese Journal of Oceanology and Limnology*. 2015 Mar;33:439-46.
17. Mitchell R, Kirchman D. The microbial ecology of marine surfaces. *Marine biodeterioration: an interdisciplinary study*. Naval Institute Press, Annapolis. 1984:49-56.
18. Joint I, Henriksen P, Fonnes GA, Bourne D, Thingstad TF, Riemann B. Competition for inorganic nutrients between phytoplankton and bacterioplankton in nutrient manipulated mesocosms. *Aquatic Microbial Ecology*. 2002 Sep 23;29(2):145-59.
19. Qian PY, Xu Y, Fusetani N. Natural products as antifouling compounds: recent progress and future perspectives. *Biofouling*. 2009 Dec 2;26(2):223-34.
20. Hadfield MG. Biofilms and marine invertebrate larvae: what bacteria produce that larvae use to choose settlement sites. *Annual review of marine science*. 2011 Jan 15;3:453-70.

21. Characklis WT, Cooksey KE. Biofilms and microbial fouling. *Advances in applied microbiology*. 1983 Dec 31;29:93-138.
22. Molino PJ, Wetherbee R. The biology of biofouling diatoms and their role in the development of microbial slimes. *Biofouling*. 2008 Jul 4;24(5):365-79.
23. Sweat LH, Johnson KB. The effects of fine-scale substratum roughness on diatom community structure in estuarine biofilms. *Biofouling*. 2013 Sep 1;29(8):879-90.
24. Yang C, Wang J, Yu Y, Liu S, Xia C. Seasonal variations in fouling diatom communities on the Yantai coast. *Chinese Journal of Oceanology and Limnology*. 2015 Mar;33:439-46.
25. Goto N, Kawamura T, Mitamura O, Terai H. Importance of extracellular organic carbon production in the total primary production by tidal-flat diatoms in comparison to phytoplankton. *Marine Ecology Progress Series*. 1999 Dec 14;190:289-95.
26. Urbani, R., Magaletti, E., Sist, P., & Cicero, A. M. (2005). Extracellular carbohydrates released by the marine diatoms *Cylindrotheca closterium*, *Thalassiosira pseudonana* and *Skeletonema costatum*: effect of P-depletion and growth status. *Science of the Total Environment*. 2005, 353(1), 300-306.
27. Mitbavkar S, Anil AC. Seasonal variations in the fouling diatom community structure from a monsoon influenced tropical estuary. *Biofouling*. 2008 Jul 30;24(6):415-26.
28. Fukao T, Kimoto K, Kotani Y. Production of transparent exopolymer particles by four diatom species. *Fisheries science*. 2010 Sep 1;76(5):755-60.
29. Urbani R, Pletikapić G, Sist P, Radić TM, Žutić V, Svetličić V. Diatom polysaccharides: extracellular production, isolation and molecular characterization. INTECH Open Access Publisher; 2012 Jan 1.
30. Takao Y, Tomaru Y, Nagasaki K, Honda D. Ecological Dynamics of Two Distinct Viruses Infecting Marine Eukaryotic Decomposer Thraustochytrids (Labyrinthulomycetes, Stramenopiles). *PloS one*. 2015 Jul 23;10(7):e0133395.