

Case Report

A Rare Case of Appendicitis

Vijay P Agrawal¹, Nitin Wasnik², Arpit Gupta³

¹Assistant Professor, Department of General surgery, NKPSIMS, Nagpur

²Associate Professor, Department of General surgery, NKPSIMS, Nagpur

³Resident, Department of General surgery, NKPSIMS, Nagpur

*Corresponding author

Dr. Vijay P Agrawal

Email: vijgunnu@gmail.com

Abstract: We present a rare case of 30 year old female presented with chronic lower abdominal pain mainly in suprapubic and right iliac fossa. Diagnostic laparoscopy was done and found 23 cm inflamed appendix involving the right iliac fossa, suprapubic region and part of left iliac fossa.

Keywords: Appendicitis, Diagnostic laparoscopy, lower abdominal pain

INTRODUCTION:

Appendectomy is the most common emergency surgery performed worldwide [1]. Appendicitis can present clinically according various position of appendix like retrocecal, Pelvic, Subcecal, Paracecal, Preileal or Postileal & subhepatic [2-4]. The length of appendix removed from a living person has been 26 cm, 20.5 cm, 18.2 cm and 17.5 cm. Boddeti RK et al removed 28 cm long appendix from a cadaver [5].

CASE REPORT:

A 32 year old female presented with chronic lower abdominal pain of 1-1.5 years. There was no history of urinary and menstrual complaints. There was no significant past history. General examination was normal. On abdominal examination, abdomen was soft and there was tenderness in right iliac fossa and suprapubic region. ultrasonography abdomen and pelvis

was normal and it was done 3-4 times during the course of 1 year as advised by different general practitioners. Her blood investigations and urinalysis was normal. CT abdomen and pelvis was advised but patient was not willing for the same due to financial concern.

Diagnostic laparoscopy was done. Intraoperative finding was 23 cm inflamed appendix involving the right iliac fossa, suprapubic region and part of left iliac fossa. Other abdominal and pelvis findings were normal. There was little difficulty in handling and separating the long appendix laparoscopically but managed successfully. Laproscopic appendectomy was done. The post operative course was uneventful. Histopathology report shows features of chronic appendicitis. The patient was discharged on 3rd post operative day.



Fig 1, 2, 3 & 4: Inflamed long appendix

DISCUSSION:

Appendix can have various sizes and positions. The average length of the appendix is 4.5 cm in neonates and 9.5 cm in adults, but this may vary between 2 cm to 20 cm. Charles McBurney described the classical presentation of appendicitis with point of maximum tenderness. Appendicitis classically present with periumbilical pain, nausea, migration of pain to the right lower quadrant, and later vomiting with fever but it may present differently due to different positions of appendix leading to difficulty in diagnosis. Common causes of appendicitis include fecolith, stricture, carcinoid tumor, pin worm etc [6]. The treatment of choice is appendectomy.

Literature also suggest the appendix length correlated highly significantly with body weight [7]. There are various case reports in literature with various length of appendix has been removed by open appendectomy. The clinical presentation of appendicitis may varies and confuse manytimes even to experienced surgeon. In our case, patient presented with chronic lower abdominal pain managed conservatively many times by general practitioners. Ultrasound, blood investigations and urine analysis were normal. Our case result also suggest the importance of diagnostic laparoscopy in chronic abdominal pain. Various length of appendix was removed previously by various authors by open surgery. There was little difficulty in handling the long appendix laparoscopically but was managed successfully.

REFERENCES:

1. Burkitt DP. The aetiology of appendicitis. *Br J Surg.* 1971 Sep;58(9):695-9.
2. Rothrock SG, Pagane J. Acute appendicitis in children: emergency department diagnosis and

management. *Annals of emergency medicine.* 2000 Jul 31;36(1):39-51.

3. Collins DC. 71000 human appendix specimens. A final report, summarizing forty years' study. *Am j Proctol.* 1963;14:365-81.
4. Solanke TF. The position, length, and content of the vermiform appendix in Nigerians. *British Journal of Surgery.* 1970 Feb 1; 57(2):100-2.
5. Ridipta S Das et al. Case report of a Mega-Appendix: Longest Non-Perforated Appendix: A Case Report *International Journal of Anatomy, Radiology and Surgery.* 2016 Jul; 5(3): SC17-SC18
6. Chaudhary R, Shukla A, Sharma K, Gupta A, Bansal M. Pin worm causing acute appendicitis: case report. *Surgery Curr Res.* 2016; 6(265):2161-1076.
7. Raschka S, Raschka C. On the relationship between body dimensions and appendix length. *Anthropologischer Anzeiger; Bericht uber die biologisch-anthropologische Literature.* 2008 Mar; 66(1):67-72.