

Original Research Article

Evaluation of Dexamethasone with Bupivacaine for infra-umbilical surgeries in children

Bhimireddy Venkata reddy¹, Mohd Inayatulla Khan²¹Associate Professor, Department of Anaesthesia, Prathima Institute of Medical Sciences, Karimnagar, Telangana²Department of Physiology, Rajiv Gandhi Institute of Medical Sciences (RIMS), Adilabad, Telangana***Corresponding author**

Dr. Bhimireddy Venkata reddy

Email: venkatreddy5959@gmail.com

Abstract: Caudal analgesia is one of the most preferred techniques of analgesia in infants and children for infra-umbilical surgeries. However it has disadvantage of short duration. Dexamethasone is a powerful anti-inflammatory agent and has been demonstrated to reduce morbidity after surgery. The aim of this study was to examine the effects of single dose of IV dexamethasone in combination with caudal block with Bupivacaine for infra-umbilical surgeries. This study was done on CAIMS, Karimnagar. 60 children undergoing infra umbilical surgeries were selected and randomly divided into two groups of 30 each. Group I received 0.25% Bupivacaine (1ml/Kg) caudal block. Group II received 0.25% Bupivacaine (1ml/Kg) and 0.5mg/Kg Dexamethasone IV single dose (maximum 10 mg) after surgery rescue analgesic consumption, pain scores and adverse effects were evaluated after 24 hrs. Pain scores measured at the interval of 1, 2, 4, 6, 12 and 24 Hours post-operatively were lower in Group II compared to Group I. Mean duration of analgesia in Group I was 270 ± 57 min and Group II it was 510 ± 78 min and p values were <0.05 . Rescue analgesic requirements were greater in Group I as compared to Group II. No significant incidences of adverse effects were also seen in both Groups. The results shows that IV dexamethasone 0.5mg/kg when used along with caudal block with Bupivacaine increased the duration of post-operative analgesia for pediatric infra-umbilical surgeries. Dexamethasone IV 0.5mg/Kg is good alternative to prolong post-operative analgesia without any adverse effects.

Keywords: Dexamethasone, Bupivacaine, Infra-umbilical Surgeries

INTRODUCTION

Pain defines pain as “An unpleasant sensory and emotional experience associated with actual or potential tissue damage or described in terms of such damage” [1]. If pain is left untreated or not managed properly it can progress to chronic pain There is a large amount of evidence for the use of multi-modal approaches to counteract pain in the pediatric population [2]. Postoperative pain management in children is challenging due to strong emotional component and it is also difficult to assess accurately the postoperative pain in children consequently greater focus is now applied on post-operative pain control in children. Various methods have evolved for providing post-op pain relief in pediatric population, some of which are not used because of side effects in children. In children narcotics could cause respiratory depression, Oral analgesics cannot be given some time after general anesthesia due to fear of vomiting and aspiration, and

fear of needle stick in case of parenteral analgesics. Regional anesthetic techniques significantly decrease the postoperative pain and systemic analgesic requirements. Caudal blocks are most preferred and safe technique for pediatric pain management including infra-umbilical surgeries. It reduces the requirements of inhaled and IV anesthetic agents and facilitates smooth and rapid recovery [3].

Bupivacaine is a long acting Local Anesthetic agent commonly used for caudal analgesia. However despite good initial analgesia from caudal blockade significant number of patients develop moderate to severe pain once duration of action of the Local anesthetic agent is over [4, 5]. Several additives are used in order to improve this drawback and improve its analgesic efficiency [6, 7]. Dexamethasone is a corticosteroid with strong anti-inflammatory effects has shown to provide post-operative analgesia and

improvement in morbidity postoperatively. Dexamethasone due to its potent anti-inflammatory effects, started to be investigated for its analgesic efficacy. In this study we tried to evaluate the analgesic additive effect of dexamethasone when administered IV along with Caudal blocks in children during infra-umbilical surgeries.

MATERIALS AND METHODS

This study was conducted in CAIMS, Karimnagar. Institutional Ethical committee permission was obtained and written informed consent was obtained from parents or guardians. All the patients included in the study were belonging for ASA (American Society of Anaesthesiologists) classification of physical status I and II undergoing herniotomies. All the patients were pre-operatively evaluated to rule out bleeding/coagulation disorders, local infections, sepsis, neurological disorders, allergy to Bupivacaine and drugs to use during and after the surgery. All parents were explained about the anaesthetic technique and perioperative course during a pre-operative visit on the day before surgery. Patients were randomly allocated in two groups. Group I (n=30) they received 0.25% Bupivacaine (1ml/Kg) caudal block. And Group II (n=30) received 0.25% Bupivacaine (1ml/Kg) and 0.5mg/Kg Dexamethasone IV single dose (maximum 10

mg). After inhalational induction of general anaesthesia with sevoflurane 8% and N₂O in 50% O₂ in spontaneous ventilation secured airway with laryngeal mask. Anaesthesia maintained by 2.5% to 3 % sevoflurane. Caudal block was performed by 22G short beveled needle in the lateral decubitus position. The solution was injected slowly with repetitive intermittent aspiration. Surgery was started after 10 min after the block. Standard monitoring was used during anaesthesia and surgery. Pethidine 0.25 mg/kg I.M. was administered as rescue analgesia. All vital signs were continuously monitored during and until the end of surgery. After transferring to the ward, rescue analgesic consumption, pain scores and adverse effects such as post-operative nausea and vomiting, urinary retention were monitored for 24 h. Analgesia when required was provided with paracetamol 15mg/kg postoperatively.

RESULTS

The table 1 shows the details of the patients selected for the study. Group I which received 0.25% Bupivacaine (1ml/Kg) caudal block and Group II received 0.25% Bupivacaine (1ml/Kg) caudal block and 0.5mg/Kg Dexamethasone IV single dose (maximum 10 mg). All the patients selected were belonging ASA classification I or II and the Mean age in years, and other parameters are given.

Table 1: Details of patients selected for the study

	Group I (Mean ± SD)	Group II (Mean ± SD)	P values
No. of patients	30	30	-
ASA I/II	22/8	25/5	0.12
Age in months	36 ± 1.5	34 ± 1.0	0.53
Height in cms	90.8 ± 10.15	89.9 ± 9.51	0.77
Wight in Kgs	12.56 ± 2.20	11.90 ± 3.5	0.94
Duration of surgery	55.6 ± 10.79	59.0 ± 10.11	0.10

The table 2 shows the duration of absolute analgesia in minutes while the value recorded in the Group I was 270 ± 57 minutes in Group II the value was 510 ± 78 the calculated p values were 0.05 and significant. The number of patients receiving rescue Pethidine 0.25mg/kg was 60 % in Group I and it was

13.34% in Group II the p values were significant. Similarly the number of patients receiving rescue paracetamol was 29 (96.67%) in Group I and 19 (63.34%) in Group II and the calculated p values were found to be significant.

Table 2: Duration of variables recorded postoperatively

Variable	Group I (Mean ± SD)	Group II (Mean ± SD)	P value
Duration of absolute analgesia (min)	270 ± 57	510 ± 78	0.05*
Emergence time	306 ± 50.4	360 ± 90	0.02*
No. of patients received rescue Pethidine 0.25 mg/kg	18 (60%)	4 (13.34%)	0.03*
No. of patients received rescue Paracetamol	29 (96.67%)	19 (63.34%)	0.05*

* P values < 0.05 is significant

The post-operative pain score recorded according to Büttner W, Finke W Analysis of behavioral and physiological parameters for the

assessment of postoperative analgesic demand in newborn, infants and children the parameters and scores given in table 3

Table 3: children and infants Postoperative pain score [8]

Item	Structure	Points
Crying	None	0
	Moaning	1
	Screaming	2
Facial Expression	Relaxed/smiling	0
	Dry mouth	1
	Grimace	2
Posture Of Trunk	Neutral	0
	Variable	1
	Rear up	2
Posture Of Legs	Neutral	0
	Kicking about	1
	Tightened legs	2
Motor Restlessness	None	0
	Moderate	1
	Restless	2

Table 4 shows the mean values of the post-operative pain scores recorded in the patients in different groups at the intervals of 1,2,4,6 ,12 and 24

hours. The calculated p values were significant at the interval of 4 hours 6 hours and 24 hours.

Table 4: post-operative pain scores

Time in hours	Group I (Mean ± SD)	Group II (Mean ± SD)	P values
1	2.5 ± 0.5	2.0 ± 0.5	0.12
2	3.0 ± 0.5	2.0 ± 0.5	0.60
4	3.5 ± 0.5	2.5 ± 0.5	0.05
6	4.5 ± 0.5	3.0 ± 0.5	0.03
12	4.5 ± 0.5	4.0 ± 0.5	0.11
24	3.5 ± 0.5	3.0 ± 0.5	0.02

* P values < 0.05 is significant

DISCUSSION

In the present study it was seen clearly that significant prolongation of post-operative analgesia was

achieved by addition of Dexamethasone IV along with caudal Bupivacaine. Steroids have potent anti-inflammatory action and they have shown to reduce

post-operative pain and swelling after various surgeries [9-12]. The biological actions of steroids have been shown to be mediated by altering functional protein synthesized by translocation of hormone receptor complex into the nucleus [13, 14]. However it is not completely understood whether the anti-inflammatory effects of glucocorticoids mediate via binding to a cytoplasmic receptor in target cells and activation of gene expression [15]. They have been shown to suppress formation of levels of bradykinin, neuropeptides from the nerve endings which are responsible for nociception in inflamed tissue [16, 17]. It also inhibits formation of cyclooxygenase isoform-2 in peripheral tissues and in the central nervous system [18]. Hval *et al.*; have demonstrated that the analgesic effect was prolonged for 3 days when dexamethasone is added to a non-steroidal anti-inflammatory drug (rofecoxib) before breast surgery [19].

In the present study there was a decrease in pain score and the number of patients receiving rescue analgesic requirements was significantly less in the Group II which received Dexamethasone IV along with Bupivacaine. The effect of IV dexamethasone is widely popular in the otolaryngological procedures with dose varying from (0.4 -1.0mg/Kg) [20-22]. In one study by H Y Hong *et al.*; using IV dexamethasone 0.5mg/kg with caudal analgesia reported that IV dexamethasone augmented the intensity and duration of postoperative analgesia without adverse effects [23]. In one similar study by Bangash L.R *et al.*; using IV dexamethasone along with caudal block in children orchidopexy has shown that the dexamethasone significantly prolongs the caudal block which is in agreement with the present study [24]. In this study we found that the duration of absolute analgesia in Group II receiving IV dexamethasone was 510 ± 78 mins Hong *et al.*; have shown the duration prolonged up to 646 minutes after IV dexamethasone. The strong anti-inflammatory action of dexamethasone is thought be involved in the process [25, 26]. In one study by Begat *et al.*; addition of 8 mg of dexamethasone to lignocaine (3mg/Kg) for intravenous regional anaesthesia produced significantly lower pain scores and need for analgesia in first 24 hours after surgery [27]. In this study we found that number of patients requiring rescue analgesia were significantly lesser in Group II as compared to Group I similar observations were made by S choudhary *et al.*; where they used caudal dexamethasone with ropivacaine for post-operative pediatric herniotomies [28]. Dexamethasone has added advantages as reported in previous studies that it does not cause problems of prolonged sedation, vomiting or urinary retention with other analgesic agents, it rather prevents postoperative

nausea and vomiting [29, 30]. The anti-emetic action of dexamethasone probably appears due to action via prostaglandin antagonism, serotonin inhibition in the gut and release of endorphins. several studies have also shown that single dose of dexamethasone used perioperatively for pediatric and adult patients undergoing surgical procedures is not associated with a significant increase in the incidence of adverse effect [21, 23]. In the present study we did not recorded any adverse due to use of dexamethasone.

CONCLUSION

The results shows that IV dexamethasone 0.5mg/kg when used along with caudal block with Bupivacaine increased the duration of post-operative analgesia for pediatric infra-umbilical surgeries. Dexamethasone IV 0.5mg/Kg is good alternative to prolong post-operative analgesia without any adverse effects.

Conflict of interest: None

Source of support: Nil

Ethical Permission: Obtained

REFERENCES

1. Strauss M, Lebowitz H, Boyle PJ, Eds. Clinical Anesthesiology, 4thed. New York: McGraw Hill 2006; 361.
2. Rahimzadeh P, Safari S, Imani F. Pediatric Chronic Pain Management: Steps toward a Neglected Area. Journal of Comprehensive Pediatrics. 2012 Oct; 4(1):47-8.
3. Deng XM, Xiao WJ, Tang GZ, Luo MP, Xu KL. The minimum local anesthetic concentration of ropivacaine for caudal analgesia in children. Anesthesia & Analgesia. 2002 Jun 1; 94(6):1465-8.
4. Silvani P, Camporesi A, Agostino MR, Salvo I. Caudal anesthesia in pediatrics: an update. Minerva anesthesiologica. 2006 Jun; 72(6):453-9.
5. Wolf AR. Tears at bedtime: a pitfall of extending paediatric day-case surgery without extending analgesia. British journal of anaesthesia. 1999 Mar; 82(3):319.
6. Khalil S, Campos C, Farag AM, Vije H, Ritchey M, Chuang A. Caudal block in children ropivacaine compared with bupivacaine. The Journal of the American Society of Anesthesiologists. 1999 Nov 1; 91(5):1279-.
7. Heshmati F, Zeinaly MB, Kh M, Mahoori AR. Intra-articular Bupivacaine versus Bupivacaine plus Dexamethasone for Analgesia after Knee Surgery.

- Iranian Red Crescent Medical Journal. 2005 May; 2006(1):39-43.
8. Büttner W, Finke W. Analysis of behavioural and physiological parameters for the assessment of postoperative analgesic demand in newborns, infants and young children: a comprehensive report on seven consecutive studies. *Pediatric Anesthesia*. 2000 May 1; 10(3):303-18.
 9. Schimmer BP, Parker KL. Adrenocorticotropic hormone; adrenocortical steroids and their synthetic analogs; inhibitors of the synthesis and actions of adrenocortical hormones. *Goodman's & Gilman's, The Pharmacological Basis of Therapeutics, Ninth Edition*, McGraw-Hill, New York. 1996:1459-85.
 10. Skjelbred P, Løkken P. Reduction of pain and swelling by a corticosteroid injected 3 hours after surgery. *European journal of clinical pharmacology*. 1982 Mar 1; 23(2):141-6.
 11. Holte K, Kehlet H. Perioperative single-dose glucocorticoid administration: pathophysiologic effects and clinical implications. *Journal of the American College of Surgeons*. 2002 Nov 1; 195(5):694-712.
 12. Bisgaard T, Klarskov B, Kehlet H, Rosenberg J. Preoperative dexamethasone improves surgical outcome after laparoscopic cholecystectomy: a randomized double-blind placebo-controlled trial. *Annals of surgery*. 2003 Nov 1; 238(5):651-60.
 13. Baxter JD, Forshman PH. Tissue effects of glucocorticoids. *Am J Med* 1972; 53(5): 573-89.
 14. Baxter JD, Schambelan M, Matulich DT, SpiNDLER BJ, Taylor AA, Bartter FC. Aldosterone receptors and the evaluation of plasma mineralocorticoid activity in normal and hypertensive states. *Journal of Clinical Investigation*. 1976 Sep; 58(3):579.
 15. Chen TL, Feldman D. Glucocorticoid receptors and actions in subpopulations of cultured rat bone cells: mechanism of dexamethasone potentiation of parathyroid hormone-stimulated cyclic AMP production. *Journal of Clinical Investigation*. 1979 Apr; 63(4):750.
 16. Hargreaves KM, Costello A. Glucocorticoids suppress levels of immunoreactive bradykinin in inflamed tissue as evaluated by microdialysis probes. *Clinical Pharmacology & Therapeutics*. 1990 Aug 1; 48(2):168-78.
 17. Hong D, Byers MR, Oswald RJ. Dexamethasone treatment reduces sensory neuropeptides and nerve sprouting reactions in injured teeth. *Pain*. 1993 Nov 1; 55(2):171-81.
 18. Ferreira SH, Cunha FQ, Lorenzetti BB, Michelin MA, Perretti M, Flower RJ, Poole S. Role of lipocortin-1 in the anti-hyperalgesic actions of dexamethasone. *British journal of pharmacology*. 1997 Jul 1; 121(5):883-8.
 19. Kjetil H, Sem TK, Ellen S, Johan R. The prolonged postoperative analgesic effect when dexamethasone is added to a nonsteroidal antiinflammatory drug (rofecoxib) before breast surgery. *Anesthesia & Analgesia*. 2007 Aug 1;105(2):481-6.
 20. Yousef GT, Ibrahim TH, Khder A, Ibrahim M. Enhancement of ropivacaine caudal analgesia using dexamethasone or magnesium in children undergoing inguinal hernia repair. *Anesthesia: Essays and Researches*. 2014 Jan 1; 8(1):13.
 21. Mohamed SK, Ibraheem AS, Abdelraheem MG. Preoperative intravenous dexamethasone combined with glossopharyngeal nerve block: role in pediatric postoperative analgesia following tonsillectomy. *European Archives of Oto-Rhino-Laryngology*. 2009 Nov 1; 266(11):1815.
 22. Vosdoganis F, Baines D. The effect of single dose intravenous dexamethasone in tonsillectomy in children. *Anaesthesia and intensive care*. 1999 Oct 1; 27(5):489.
 23. Hong JY, Han SW, Kim WO, Kim EJ, Kil HK. Effect of dexamethasone in combination with caudal analgesia on postoperative pain control in day-case paediatric orchiopexy. *British journal of anaesthesia*. 2010 Oct 1; 105(4):506-10.
 24. Bangash L, Afzal F, Hussain S, Ali K. Comparison of caudal block alone with caudal block plus intravenous dexamethasone for postoperative analgesia in children undergoing orchidopexy under general anaesthesia. *Biomedica*. 2014 Oct; 30(4):267.
 25. Jokela RM, Ahonen JV, Tallgren MK, Marjakangas PC, Korttila KT. The effective analgesic dose of dexamethasone after laparoscopic hysterectomy. *Anesthesia & Analgesia*. 2009 Aug 1; 109(2):607-15.
 26. Bajwa SJ, Kaur J, Bajwa SK, Bakshi G, Singh K, Panda A. Caudal ropivacaine-clonidine: A better post-operative analgesic approach. *Indian journal of anaesthesia*. 2010 May 1; 54(3):226.
 27. Bigat Z, Boztug N, Hadimioglu N, Cete N, Coskunfirat N, Ertok E. Does dexamethasone improve the quality of intravenous regional anesthesia and analgesia? A randomized, controlled clinical study. *Anesthesia & Analgesia*. 2006 Feb 1; 102(2):605-9.
 28. Bangash L, Afzal F, Hussain S, Ali K. Comparison of caudal block alone with caudal block plus intravenous dexamethasone for postoperative analgesia in children undergoing orchidopexy

- under general anaesthesia. *Biomedica*. 2014 Oct; 30(4):267.
29. Choudhary S, Dogra N, Dogra J, Jain P, Ola SK, Ratre B. Evaluation of caudal dexamethasone with ropivacaine for post-operative analgesia in paediatric herniotomies: A randomised controlled study. *Indian journal of anaesthesia*. 2016 Jan; 60(1):30.
30. Malhotra D, Gurcoo S, Qazi S, Gupta S. Randomized comparative efficacy of dexamethasone to prevent postextubation upper airway complications in children and adults in ICU. *Indian journal of anaesthesia*. 2009 Jul 1; 53(4):442.