

Case Report

Infantile Yolk Sac Tumor of Testis – A Case ReportDr. N. L. N. Moorthy¹, Dr M. Vithaleshwar Rao², Dr S. Padmaja³, Dr Sangireddy Jayabhaskar Reddy⁴¹Professor Radiology, ²Assistant Professor Radiology, ³Senior Resident Radiology, ⁴Professor and Hod Pathology
Department of radiology and pathology, Apollo institute of medical science and research, Hyderabad-500096***Corresponding author**

Dr M. Vithaleshwar Rao

Email: dr.vithal@yahoo.co.in

Abstract: Yolk sac tumor is the most common tumor of testis in children below 3 years. Most of these tumors present as painless, enlarged testicular masses. They are most frequently associated with elevated alpha fetoprotein (AFP) levels. We report a case of yolk sac tumor in an 11 month old child who presented as a painless testicular mass.

Keywords: yolk sac tumor- testicular masses- alpha fetoprotein.

INTRODUCTION:

Testicular tumors are extremely uncommon in children, forming 1-2 % of all solid childhood tumors. Most of the paediatric testicular tumors are malignant (more than 95 %) and present as painless masses. Testicular tumors are either germ cell tumor or non-germ cell tumors. Germ cell tumor is divided into seminomas and non seminomatous tumors. The non seminomatous tumors which are common in children include teratomas, embryonal carcinomas, chorio carcinoma and yolk sac tumors [1] Of the different testicular tumors, yolk sac tumors are the most common in boys (80 %) with a peak age incidence of 2 years [2]. Malignant testicular tumors typically present as painless solid masses. Serum AFP levels are abnormal in many cases. Ultrasound scrotum is highly sensitive in detecting these tumors.

CASE REPORT:

An 11 month old infant was presented with gradually increasing painless swelling of right hemiscrotum since three months. There was no history of fever, trauma or urinary tract infection. The scrotal mass measured 6x5cm, soft and fluctuant with local rise of temperature. Routine lab investigations were normal.

High resolution ultrasound of scrotum revealed an enlarged, globular right testis with multiple cystic spaces. No calcification was noted within the mass (fig1A &B). Minimal free fluid in right hemiscrotum was present. Color Doppler showed increased vascularity in the testis (fig 2). The right epididymis and spermatic cord were thickened. No significant regional lymphadenopathy detected. Chest radiograph was normal. A provisional diagnosis of solid testicular tumor was made. AFP was normal. Right orchidectomy was done. Histopathology revealed tumor composed of many glomeruloid bodies and few glandular structures and oval to polygonal cells with round hyperchromatic nuclei and many atypical mitoses. Cytoplasm was abundant and pale eosinophilic with focal areas showing intracytoplasmic hyaline globules. Areas of necrosis are seen within the tumor. Tumor was seen infiltrating the capsule in focal area. Tumor emboli are seen within the vessels (fig 3). Adjacent areas showed few uninvolved seminiferous tubules without spermatogenesis. Epididymis appeared normal. No evidence of invasion by the tumor. Section of the spermatic cord was unremarkable and showed no evidence of invasion by the tumor.



Fig 1 & B : Ultrasound images of testis show diffuse enlargement with cystic areas.




Fig 2: Color Doppler image shows increased intratesticular vascular flow




Fig 3: Histopathology shows tumor consisting of glomeruloid bodies and tumor emboli

DISCUSSION:

The pattern of testicular tumors in children and adults are different. The most common cause of testicular tumors in adults is germ cell tumor (95 %) and germ cell tumors account for 60-70 % of testicular tumors in children. Teratoma and yolk sac tumor are the most common type of germ cell tumors in children. The other causes of testicular masses in children include epidermoid cysts, Sertoli cell tumors, Leydig cell tumors, juvenile granulosa cell tumor, gonadoblastoma, cystic dysplasia of testis, lymphoma, leukemia etc. The most common testicular tumor in children is pure yolk sac tumor (upto 80%) occurring mostly below two years of age. Also called as endodermal sinus tumor, these tumors are found within the testis parenchyma. The macroscopic features of the tumor include solid gray myxoid appearance with heterogenous areas of hemorrhage, cysts and necrosis. The microscopic appearance is the presence of spider web like network found in vacuolated cytoplasm [3]. The other microscopic pattern noted in yolk sac tumors is endodermal sinus pattern called Schiller –Duvall body which is pathognomonic of yolk sac tumors. Beside alpha fetoprotein, other markers that are highly sensitive for yolk sac tumor are cytokeratin, vimentin, placental alkaline phosphatase and glypican 3. On ultrasonography the testis is enlarged, the parenchyma is replaced by heterogenous solid tumor. They may be associated with hydrocele. Besides clinical examination, ultrasound scanning is very useful in localizing the tumor especially in detecting small sized lesions. Intratesticular tumor is to be considered as malignant until proved otherwise [4].

By careful analysis of ultrasound findings it is possible to identify the tumor pathology in testicular masses [5-8]. On ultrasound, yolk sac tumors appear as heterogenous solid masses replacing normal testicular parenchyma. Associated cystic areas suggestive of necrosis may be seen. Regional lymphadenopathy is frequent as the tumor spreads by lymphatic system. Besides yolk sac tumor the other most common germ cell tumor in children is teratoma. The ultrasound findings are quite variable depending on the components of the three germinal layers in the teratoma. Teratoma can be cystic with solid components and may also contain calcification and intratumoral fat. A solid intratesticular mass with increased vascularity is mostly a testicular tumor as the sensitivity and specificity on ultrasound in detecting malignancy is 92-98% and 95-98% respectively [9, 10]. Seminomas are generally hypoechoic and homogenous while non seminomatous germ cell tumors are more heterogenous with cystic spaces and show calcifications frequently [11].

Leukemias and lymphomas are rare causes of testicular masses in children. On ultrasound they appear as single or multifocal hypoechoic masses. Benign conditions can sometimes appear similar to tumors in testis. Epidermal cysts typically seen as round hypoechoic lesion with a hyperechoic rim. Other benign conditions that mimic testicular tumors in children include adrenal rests, testicular infarcts, tunica albuginea and intratesticular Cysts [1].

D.Y.Huang [12] reviewed the different imaging modalities in the diagnosis of testicular malignancies and found that contrast enhanced ultrasound and elastography will help in further improving the characterization of malignant and benign testicular masses. The most commonly elevated tumor marker in testicular tumors is alpha fetoprotein (AFP) seen in yolk sac tumors and in some cases of teratomas, whereas pure choriocarcinoma and pure seminoma do not produce AFP. The other tumor markers produced in testicular masses are human chorionic gonadotropin, lactate dehydrogenase. The diagnosis of testicular tumor is made only on histopathology alone and not based on tumor marker as they lack sensitivity and specificity [3]. They are helpful in monitoring the treatment response.

Clinico pathological and treatment protocols and management of yolk sac tumors of testis were reviewed by many authors in the literature [14-17]. Liu and others [15] in the review of 11 cases of testicular yolk sac tumors found Glypican- 3 as a very sensitive diagnostic antibody for yolk sac tumors. Radical inguinal orchiectomy followed by cisplatin based chemotherapy is adequate for stage 1 yolk sac tumor of testis in children [18-22].

CONCLUSION:

Painless testicular masses in children should always be considered as very important symptom and histopathological diagnosis along with AFP levels should be available always for further management.

REFERENCES:

1. Sung EK, Setty BN, Castro-Aragon I. Sonography of the pediatric scrotum: emphasis on the Ts—torsion, trauma, and tumors. *American Journal of Roentgenology*. 2012 May; 198(5):996-1003.
2. Aso C, Enriquez G, Fité M, Torán N, Piró C, Piqueras J, Lucaya J. Gray-scale and color Doppler sonography of scrotal disorders in children: an update. *Radiographics*. 2005 Sep; 25(5):1197-214.
3. Ximing J Yang, chief editor Liang Cheng MD Yolk sac tumor pathology Medscape jan 19, 2015

4. Skoog SJ. Benign and malignant pediatric scrotal masses. *Pediatric Clinics of North America*. 1997 Oct 1; 44(5):1229-50.
5. McDonald MW, Reed AB, Tran PT, Evans LA. Testicular tumor ultrasound characteristics and association with histopathology. *Urologia internationalis*. 2012; 89(2):196-202.
6. Hebert SC, Chong WK, Deurdulian C. Essentials of scrotal ultrasound: A review of frequently encountered abnormalities. *Applied Radiology*. 2012 Sep 7; 41(9):7.
7. Shtricker A, Silver D, Sorin E, Schreiber L, Katlowitz N, Tsivian A, Katlowitz K, Benjamin S, Sidi AA. The value of testicular ultrasound in the prediction of the type and size of testicular tumors. *International braz j urol*. 2015 Aug; 41(4):655-60.
8. Orlando Catalano. Antonio Nunziata, Alfredo Siani. Chapter 6.9 in *Fundamentals in Oncologic Ultrasound: Sonographic Imaging and Intervention*: Springer Italy, 2009: 320-330.
9. Yagil Y, Naroditsky I, Milhem J, Leiba R, Leiderman M, Badaan S, Gaitini D. Role of Doppler ultrasonography in the triage of acute scrotum in the emergency department. *Journal of Ultrasound in Medicine*. 2010 Jan 1; 29(1):11-21.
10. Guthrie JA, Fowler RC. Ultrasound diagnosis of testicular tumours presenting as epididymal disease. *Clinical radiology*. 1992 Dec 1; 46(6):397-400.
11. Coursey Moreno C, Small WC, Camacho JC, Master V, Kokabi N, Lewis M, Hartman M, Mittal PK. Testicular tumors: what radiologists need to know—differential diagnosis, staging, and management. *Radiographics*. 2015 Mar 12; 35(2):400-15.
12. Huang DY, Sidhu PS. Focal testicular lesions: colour Doppler ultrasound, contrast-enhanced ultrasound and tissue elastography as adjuvants to the diagnosis. *The British journal of radiology*. 2012 Nov; 85(special_issue_1):S41-53.
13. Milose JC, Filson CP, Weizer AZ, Hafez KS, Montgomery JS. Role of biochemical markers in testicular cancer: diagnosis, staging, and surveillance. *Open access journal of urology*. 2012; 4:1.
14. Khan IU, Jose J, Fawazy T, Hadi WA, Sharma PK. Testicular yolk sac tumour in an eight month old child: A case report. *Gulf Medical Journal*. 2012; 1 (1): 37-40.
15. F.Liu, S. Cen, X. Gong, X.Xie Testicular yolk sac tumours in infants: A clinicopathological study of 11 cases. *Researchgate.net/ publication*, July 2012.
16. Wei Y, Wu S, Lin T, He D, Li X, Liu J, Liu X, Hua Y, Lu P, Wei G. Testicular yolk sac tumors in children: a review of 61 patients over 19 years. *World journal of surgical oncology*. 2014 Dec 29; 12(1):400.
17. Cornejo KM, Frazier L, Lee RS, Kozakewich HP, Young RH. Yolk sac Tumor of the Testis in Infants and Children: A Clinicopathological Analysis of cases. *Am J Surg Pathol*. 2015 Aug; 39(8); 1121-31.
18. Liu HC, Liang DC, Chen SH, Liu FL, Chang PY, Sheu JC, Wang NL. The stage I yolk sac tumor of testis in children younger than 2 years, chemotherapy or not? *Pediatr Hematol Oncol*. 1998 May-Jun; 15(3): 223-8.
19. Kuo JY, Hsieh YL, Chin TW, Wei CF, Chen KK, Chang LS. Testicular yolk sac tumors in children. *Zhonghua Yi Xue Za Zhi (Taipei)*. 1999 Feb; 62(2): 92-7.
20. Karnak I, Ciftci AO, Şenocak ME, Tanyel FC, Büyükpamukçu N. Colonic atresia: surgical management and outcome. *Pediatric surgery international*. 2001 Nov 8; 17(8):631-5.
21. Jonathan H. Ross MD Testicular tumors in *Pediatric Urology* 1-4. <http://pediatricurology/book.com/tumors.html>.
22. Zahran MH, Helmy TE, Hafez AT, Dawaba M. Prepubertal testicular tumours: Should testicular-sparing surgery be considered? A single-institution experience and review of the literature. *Arab journal of urology*. 2014 Jun 30; 12(2):130-6.