

## A Study of Etiology and Outcome of Acute Respiratory Distress Syndrome in a Rural Tertiary Care Center

Dr. Poola Sriharika<sup>1</sup>, Dr. R. Lakshmi Visruja<sup>2\*</sup>, Dr. Srinivasa Rao<sup>3</sup>, Dr. Y. J. Visweswara Reddy<sup>4</sup>

<sup>1</sup>Postgraduate Resident, Department of General Medicine, PESIMSR/NTRUHS, Kuppam, India

<sup>2</sup>Assistant Professor, Department of General Medicine, PESIMSR/NTRUHS, Kuppam, India

<sup>3,4</sup>Professor, Department of General Medicine, PESIMSR/NTRUHS, Kuppam, India

### Original Research Article

#### \*Corresponding author

Dr. R. Lakshmi Visruja

#### Article History

Received: 26.02.2018

Accepted: 09.03.2018

Published: 30.03.2018

#### DOI:

10.21276/sjams.2018.6.3.42



**Abstract:** Acute respiratory distress syndrome (ARDS) is one of the major causes for acute respiratory failure characterized by bilateral chest infiltrates and decreased lung compliance. Several etiological factors cause ARDS, both direct and indirect injury to lungs. It has a high mortality rate. This study was a prospective observational study and included 50 patients with a diagnosis of ARDS. This study was undertaken to elucidate the causes and outcome of patients admitted with ARDS in the intensive care unit (ICU) of a tertiary care center. New Berlin criteria were used to classify them as Mild, Moderate and Severe ARDS. Severe ARDS was found in 30%, moderate ARDS in 58% and mild ARDS was seen in 12% of subjects. Pneumonia followed by tropical infections was the major etiological causes. Tropical infections contributed to 26% of ARDS, of which Dengue fever contributed to 12% of ARDS in the subjects and malaria was seen to be causing ARDS in 8% of subjects. Most of the patients (96%) required ventilator support, 52% of them requiring invasive ventilation and 44% requiring non – invasive ventilation. Mortality was 62% in this study. Higher mortality rates were seen in age groups above 40 years and males had slightly higher mortality rates than females. ABG (Arterial blood gas analysis) showed higher PaCO<sub>2</sub> (partial pressure of carbon dioxide) and lower PaO<sub>2</sub>/FiO<sub>2</sub> (PaO<sub>2</sub>-partial pressure of oxygen/FiO<sub>2</sub>- fraction of inspired oxygen) were seen in non-survivor group. Tropical infections, especially in rural areas, contribute significantly to ARDS which needs to be stressed up on. These findings may help intensivists in instituting early effective empirical regimens depending on the regional tropical diseases.

**Keywords:** Acute respiratory distress syndrome, Berlin criteria, Arterial blood gas, PaO<sub>2</sub>/FiO<sub>2</sub>.

### INTRODUCTION

In 1967, Ashbaugh first identified acute respiratory distress syndrome (ARDS), described 12 patients who had bilateral chest infiltrates and reduced lung compliance [1] with acute respiratory failure refractory to oxygen therapy. ARDS can be caused by direct and indirect injury. The direct causes most commonly are lung related infection and aspiration pneumonia. Indirect causes are extra pulmonary among which systemic sepsis is the most common one [2].

Chronic liver disease, non-pulmonary organ dysfunction, sepsis and advanced age have been identified to predict the risk of death in ARDS [3, 4]. Significant causes of ARDS other than pulmonary, in rural areas are tropical infections like malaria, dengue and leptospirosis [5].

Mortality rates for ARDS vary widely; according to a few studies it was about 58 % [3, 4]. Several studies have shown that initial oxygenation

level does not have any prognostic significance in determining mortality, but if there is no improvement in oxygenation it further leads to poor outcome.

This study was undertaken to elucidate the causes and outcome of patients admitted with ARDS in the intensive care unit (ICU) of a tertiary care and medical college hospital catering to a rural population of south India.

### MATERIALS AND METHODS

The study done was a prospective observational study conducted in ICU of PESIMSR (Peoples education society institute of Medical Sciences and Research) during the year 2016-2017. The hospital is located in a rural area (Kuppam) of Andhra Pradesh, south India.

A total of 50 patients aged above 18years with a medical diagnosis leading to ARDS were included in

the study. Patients with surgical causes of ARDS like trauma and burns were excluded.

Diagnosis of ARDS was done based on history, physical examination, chest radiograph and arterial blood gas analysis. Based on the New Berlin criteria [6], ARDS was categorized into mild, moderate and severe ARDS. Echocardiography was done for all the patients to rule out cardiac causes. Relevant laboratory investigations were done. Bacterial pneumonia was diagnosed by chest radiograph and examination of sputum. Viral pneumonia was diagnosed when total leukocyte counts were normal, chest radiograph suggestive of pneumonia and all other tests including gram stain, culture and sensitivity tests were negative.

The ethical committee of the medical college hospital approved this study. The data were entered into MS Excel sheet and SPSS software was used for statistical analysis. Descriptive inferential statistical analysis has been carried out. Chi-square/ Fischer exact test has been used to find the significance of study parameters on categorical scale between groups .P value < 0.05 was taken as significant value.

**RESULTS**


During the study period, 50 patients were admitted with ARDS as diagnosis to medical ICU. Most of the subjects were relatively young with a mean age of 41.78 ± 16.87years. Nearly a quarter of them were in the age group 31-40 years (24%) and 21-30 years (22%) each. Distribution of Male (52%) and female (48%) were as shown in table 1.

**Table-1: Correlation of age & gender with incidence of mortality of patients**

variables	Mortality		Total (n=50)	P value
	No (n=19)	Yes (n=31)		
Age in years				
• <20	2(10.5%)	2(6.5%)	4(8%)	0.013*
• 20-30	8(42.1%)	3(9.7%)	11(22%)	
• 31-40	6(31.6%)	6(19.4%)	12(24%)	
• 41-50	1(5.3%)	7(22.6%)	8(16%)	
• 51-60	0(0%)	7(22.6%)	7(14%)	
• >60	2(10.5%)	6(19.4%)	8(16%)	
Gender				
• Female	10(52.6%)	14(45.2%)	24(48%)	0.608
• Male	9(47.4%)	17(54.8%)	26(52%)	

Most of them presented with respiratory complaints (80% of subjects) with 54% of subjects presenting with breathlessness. Majority (90%) of them did not have pre-existing lung diseases. Patients were classified according to New Berlin Criteria based on arterial blood gas (ABG) as mild, moderate and severe ARDS. Severe ARDS was found in 30%, moderate ARDS in 58% and mild ARDS was seen in 12% of subjects.

Significantly large number of patients presented with infections. (As shown in the Figure1). The commonest etiology was pneumonia (44%). Tropical infections contributed to 26% of ARDS, of which Dengue fever contributed to 12% of ARDS in the subjects and malaria was seen to be causing ARDS in 8% of subjects.



**Fig-1: Clinical Diagnosis with ARDS**

Most of the patients (96%) required ventilatory support, 52% of them requiring invasive ventilation and 44% requiring non – invasive ventilation. Mortality was 62% in this study. Higher mortality rates were seen in age groups above 40 years and males had slightly higher mortality rates than females did (52% vs 48%). Mortality rates for patients on mechanical invasive ventilation were 48.4% compared to 44% in patients on non-invasive ventilation.

ABG showed significant PCO<sub>2</sub> (Partial pressure of carbon dioxide) and PaO<sub>2</sub> / FiO<sub>2</sub> (PaO<sub>2</sub>-partial pressure of oxygen, FiO<sub>2</sub>- fraction of inspired oxygen) ratio in the present study. PaCO<sub>2</sub> was higher in non-survivor group as compared to survivor group (37.28±12.77 VS 28.23±4.94) which was statistically significant (as shown as table 2).

**Table-2: Comparison of ABG according to Mortality of patients**

Parameters	Mortality		Total	P value
	No	Yes		
pH	7.23±0.21	7.10±0.28	7.15±0.26	0.070+
PaO <sub>2</sub>	99.36±24.14	90.39±48.08	93.80±40.61	0.454
PaCO <sub>2</sub>	28.23±4.94	37.28±12.77	33.84±11.33	0.005**
HCO <sub>3</sub>	19.07±4.34	17.46±3.65	18.07±3.96	0.165
PaO <sub>2</sub> /FIO <sub>2</sub>	212.14±61.00	166.92±55.23	184.10±61.04	0.010**

PaO<sub>2</sub>-partial pressure of oxygen, PaCO<sub>2</sub>-partial pressure of carbondioxide, HCO<sub>3</sub>- Bicarbonate, FiO<sub>2</sub>- fraction of inspired oxygen

PaO<sub>2</sub>/FiO<sub>2</sub> ratio was lower in non-survivor group (166.92 ± 55.23) and had high mortality as compared to survivor group (212.14±61) was seen in severe ARDS (as shown in table 3).

**Table-3: Comparison of PaO<sub>2</sub>/FIO<sub>2</sub> with incidence of mortality of patients**

PaO <sub>2</sub> /FIO <sub>2</sub>	Mortality		Total
	No	Yes	
<100 SEVERE ARDS	3(15.8%)	12(38.7%)	15(30%)
100-200 MODERATE ARDS	11(57.9%)	18(58.1%)	29(58%)
>200-300 MILD ARDS	5(26.3%)	1(3.2%)	6(12%)
Total	19(100%)	31(100%)	50(100%)

PaO<sub>2</sub>-partial pressure of oxygen, FiO<sub>2</sub>- fraction of inspired oxygen

**DISCUSSION**

This study focused on the etiology and outcome of ARDS. Mean age of the subjects was 41.78years (SD ± 16.87). This was similar to studies from other parts of India [2,7]. A few studies showed high mean age above 50 years [8-10]. Our study showed near equal distribution in males and females which is unlike some national and international studies [2, 8, 9, 11].

The commonest presentation was with respiratory complaints, breathlessness being present in 54% of subjects [12], followed by cough in 24% of subjects. Most of the patients did not have co-morbidities. Most of our patients were in moderate ARDS (58%) as described by the New Berlin criteria [6].

Infective pneumonia was the commonest cause of direct lung injury (44%) [2]. Dengue, malaria and systemic sepsis contributed majorly to indirect lung injury [2]. High incidence of tropical infections was noted (26%). This may be because of favorable climate for infective agent and the vector. One more study from South India showed similar results, 26% of total ARDS cases were contributed by tropical infections [8].

Streptococcus pneumonia was the commonest organism isolated in culture- positive cases in our subjects followed by pseudomonas.

Non – infectious causes contributed to 22% of the etiology. Acute pancreatitis, aspiration pneumonia, organophosphorus poisoning contributed 2% each to the etiological causes of ARDS. Dengue fever contributed to 12% of the etiology. Recently many dengue related ARDS case reports have been reported [13]. Wang *et al.* found an incidence of 1.8% ARDS in a total of 606 dengue patients [14]. A few other studies have shown that dengue hemorrhagic fever can result in ARDS [15-17].

Malaria contributed to 8% of the ARDS cases. ARDS has been described as a major cause of death in adults with severe malaria [18, 19]. Mortality is as high as 80% in malaria associated ARDS, without ventilatory support [19]. Ventilatory support was required in 52% of the ARDS patients. Mortality rate estimated in many studies varied between 37% and 57 % [2, 7, 20]. In our study, mortality rate was 62% in 2007. Lancet review mentioned 25-30% as the mortality rates [21] while in 2008, Phua *et al* described it to be around 30-60% [22]. Our study had a higher mortality

rate probably due to differing baseline characteristics, etiology and differing severity of ARDS compared to other studies. Higher mortality rates were seen in age groups above 40 years.

Higher mortality rates were seen in patients with severe ARDS (12 out of 15 patients expired) followed by moderate ARDS (18 out of 29 patients expired). Only one patient with mild ARDS succumbed out of 6 patients [7]. Severity can be considered as individual predictor of mortality, findings similar to patients in a study conducted by Bhadade and Rajbanshi [7, 12]. ABG revealed significant association with PCO<sub>2</sub>/FiO<sub>2</sub> ratio in the present study.

#### LIMITATION

This study had some limitations. The study was an observational study and sample size was small. APACHE, Lung injury scoring was not done. Due to the lack of institutional facility we could not do viral PCR.

#### CONCLUSION

The commonest cause of ARDS is pneumonia. Tropical infections, especially in rural areas, contribute significantly to ARDS which needs to be stressed up on. Lower PaO<sub>2</sub>/FiO<sub>2</sub> ratio contributed to higher mortality. These findings may help intensivists in instituting early effective empirical regimens depending on the regional tropical diseases.

#### REFERENCES

1. Ware LB, Matthay MA. The acute respiratory distress syndrome. *New England Journal of Medicine*. 2000 May 4; 342(18):1334-49.
2. Agarwal R, Aggarwal AN, Gupta D, Behera D, Jindal SK. Etiology and outcomes of pulmonary and extrapulmonary acute lung injury/ARDS in a respiratory ICU in North India. *Chest*. 2006 Sep 1; 130(3):724-9.
3. Doyle RL, Szaflarski N, Modin GW, Wiener-Kronish JP, Matthay MA. Identification of patients with acute lung injury. Predictors of mortality. *American journal of respiratory and critical care medicine*. 1995 Dec; 152(6):1818-24.
4. Monchi M, Bellenfant F, Cariou A, Joly LM, Thebert D, Laurent I, Dhainaut JF, Brunet F. Early predictive factors of survival in the acute respiratory distress syndrome: a multivariate analysis. *American Journal of Respiratory and Critical Care Medicine*. 1998 Oct 1; 158(4):1076-81.
5. George T, Viswanathan S, Karnam AHF, Abraham G. Etiology and outcomes of ARDS in a rural-urban fringe hospital of South India. *Crit Care Res Pract*. 2014; 1-8.
6. Force AD. Acute respiratory distress syndrome. *Jama*. 2012 Jun 20; 307(23):2526-33.
7. Bhadade RR, De Souza RA, Harde MJ, Khot A. Clinical characteristics and outcomes of patients

- with acute lung injury and ARDS. *Journal of postgraduate medicine*. 2011 Oct 1; 57(4):286.
8. Estenssoro E, Dubin A, Laffaire E, Canales H, Saenz G, Moseinco M, Pozo M, Gomez A, Baredes N, Jannello G, Osatnik J. Incidence, clinical course, and outcome in 217 patients with acute respiratory distress syndrome. *Crit Care Med*. 2002;30(11):2450-2456.
9. Zilberberg MD, Epstein SK. Acute lung injury in the medical ICU: comorbid conditions, age, etiology, and hospital outcome. *American journal of respiratory and critical care medicine*. 1998 Apr 1; 157(4):1159-64.
10. Irish Critical Care Trials Group. Acute lung injury and the acute respiratory distress syndrome in Ireland: a prospective audit of epidemiology and management. *Critical care*. 2008 Feb 1; 12(1):R30.
11. Brun-Buisson C, Minelli C, Bertolini G, Brazzi L, Pimentel J, Lewandowski K, Bion J, Romand JA, Villar J, Thorsteinsson A, Damas P. Epidemiology and outcome of acute lung injury in European intensive care units. *Intensive care medicine*. 2004 Jan 1; 30(1):51-61.
12. Lalit Kumar Rajbanshi, Shambhu Bahadur Karki, Batsalya Arjyal and Rakesh Mandal Rajbanshi. Etiology, clinical profile and outcome of acute respiratory distress syndrome patients in eastern nepal. *EJPMR*. 2017; 4(8): 541-545.
13. Belagavi C, Sunil HS, Sudhir U & Punith K. Adult respiratory distress syndrome in Dengue -a case report. *Al Ameen Journal of Medical Science*. 2011; 4: 405-407.
14. Wang CC, Liu SF, Liao SC, Lee K, Liu JW, Lin AS, Wu CC, Chung YH, Lin MC. Acute respiratory failure in adult patients with dengue virus infection. *The American journal of tropical medicine and hygiene*. 2007 Jul 1; 77(1):151-8.
15. Lum LC, Thong MK, Cheah YK, Lam SK. Dengue-associated adult respiratory distress syndrome. *Annals of tropical paediatrics*. 1995 Dec 1; 15(4):335-9.
16. Sharma SK, Gupta BS, Devpura G, Agarwal A, Anand S. Pulmonary haemorrhage syndrome associated with dengue haemorrhagic fever. *JOURNAL-ASSOCIATION OF PHYSICIANS OF INDIA*. 2007 Oct 5; 55(T):729.
17. Setlik RF, Ouellette D, Morgan J, McAllister KC, Dorsey D, Agan BK, Horvath L, Zimmerman MK, Purcell B. Pulmonary hemorrhage syndrome associated with an autochthonous case of dengue hemorrhagic fever. *Southern Medical Journal*. 2004 Jul 1; 97(7):688-92.
18. World Health Organization. Severe falciparum malaria. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2000 Apr 1; 94:1-90.
19. Taylor WR, White NJ. Malaria and the lung. *Clinics in chest medicine*. 2002 Jun 1; 23(2):457-68.

20. Singh G, Gladdy G, Chandy TT, Sen N. Incidence and outcome of acute lung injury and acute respiratory distress syndrome in the surgical intensive care unit. *Indian journal of critical care medicine: peer-reviewed, official publication of Indian Society of Critical Care Medicine*. 2014 Oct; 18(10):659.
21. Wheeler AP, Bernard GR. Acute lung injury and the acute respiratory distress syndrome: a clinical review. *The Lancet*. 2007 May 5; 369(9572):1553-64.
22. Phua J, Stewart TE, Ferguson ND. Acute respiratory distress syndrome 40 years later: time to revisit its definition. *Critical care medicine*. 2008 Oct 1; 36(10):2912-21.