

To Find Out Various Clinical Characteristics of Patients with Rheumatic Mitral Stenosis Registered in Cardiology OPD from 1992 To 2013

Dr. Nitin Rawat¹, Dr. Vinayak vishwanath Jatale^{2*}, Dr. Nitin Rawat Clinic, Barwani¹

¹General Physician, Dept. of Medicine, Index Medical College, Indore, India

²Asst. Prof, Dept. of Medicine, Index Medical College, Indore, India

Original Research Article

*Corresponding author

Dr. Vinayak Vishwanath
Jatale

Article History

Received: 12.04.2018

Accepted: 23.04.2018

Published: 30.04.2018

DOI:

10.21276/sjams.2018.6.4.25



Abstract: Mitral Stenosis is a sequel to Rheumatic fever which follows GAS Pharyngitis. Recurrent hospitalization for heart failure, development of atrial fibrillation, systemic embolization and cerebral embolization are associated with high morbidity and mortality seen during natural history of Mitral Stenosis. The Rheumatic Mitral Stenosis affect young people particularly females. Unoperatic patients with significant Mitral Stenosis carry high risk of Cardio embolism and recurrent hospitalization with associated high morbidity and mortality.

Keywords: Clinical, Rheumatic, Mitral Stenosis & Cardiology.

INTRODUCTION

The initial mitral valvulitis, as part of endocarditis occurring during rheumatic fever, results in cicatrization and fibrosis during healing process results in commissural fusion and disease of sub-valvular apparatus causing obstruction to left ventricular inflow.

Mitral valve disease, especially MS is associated with high risk of Cardioembolic events. Numerous clinical and echo cardiographic factor are associate with a high risk thromboembolism in these patients. These include atrial fibrillation, left atrial size, left appendage dysfunction, duaration of mitral valve disease, and presence of spontaneous contrast (SEC) in the left atrium [1-4]. Nearly one fifth of all embolic event in patient mitral stenosis occur in patient in sinus rhythm. However, despite this higher risk, the limited data on predictors of clinical stroke in these patients.

Moreover there is no unanimity amongst various guidelines in the use of anti-coagulants in patients with MS. Currently there is paucity of data on whether to initiate anti-coagulation in patients with mitral stenosis who are in NSR or not as the risk marker for thromboembolism are poorly defined.

MATERIALS & METHODS

This Study was carried out in the Department of Medicine, Mahatma Gandhi Memorial Medical College and Maharaja Yashwant Rao Hospital Indore, from January 2013 to October 2014. We analyzed case files of all the patients who were diagnosed to have Rheumatic Mitral Stenosis as dominant lesion and were registered in cardiology OPD of Maharaja Yashwant Rao Hospital Indore during period of 1992 to 2013, case files with complete information were accepted for analysis. Case files with incomplete information were not analyzed. If required, we tried to obtain information on hospitalizations, Cardioembolic events and mortality by contacting patient's families. The consent for permission to use patient's related

data, so obtained, was taken from surviving patients or their relatives.

Inclusion criteria

Case files of patients with diagnosis of mitral stenosis as dominant lesion registered in cardiology OPD from January 1992 to December 2013 were analyzed.

Exclusion criteria

- Case files with incomplete data in form of inadequate clinical and laboratory Information including patient contact details and physical examination were excluded.
- Patients with more than mild mitral regurgitation or more than mild aortic valve stenosis / regurgitation were not included.
- After analyzing case records of these patients, the following information was recorded: age, gender, Cardiac examination findings, laboratory findings, Echocardiographic findings, hospitalization record, OPD follow up record, Therapy received,

Cardioembolic events (cerebral or peripheral), change in NYHA class, change in rhythm, various hospitalization episodes with reasons and mortality.

- Statistical Methods: The data were analyzed using statistical software SPSS IBM Versions 20, The data was subjected to Kaplan Meire analysis and various survival curves were drawn. The complications including mortality were compared among various subgroups.

Criteria for Diagnosis

Definition

Mitral Stenosis - Mitral stenosis is a narrowing of heart's mitral valve orifice which is 4-6 cm². When the orifices is reduced to approximately 2cm², which is considered to represent mild MS. When the mitral valve opening is reduced to 1.5 to 1 cm², which is considered to moderate MS.

When the mitral valve opening is reduced to 1 cm², which is considered to severe MS. CLINICAL DIAGNOSIS of mitral stenosis was based upon the presence of rough, rolling rasping diastolic or presystolic murmur.

Strokes [5]

“Rapidly developing clinical sign of focal (or global) disturbance of cerebral function, with symptoms lasting 24 hours or longer or leading to death, with no apparent cause than of vascular origin

- Ischemic stroke [6] are caused by sudden occlusion of arteries supplying the brain either due to a thrombus at the site of occlusion or formed in another part of the circulation.
- Transient ischemic attack: TIAs are episodes of stroke symptoms that last only briefly; the standard definition of duration is <24 hours, but most TIAs last <1 hour.

- Silent Stroke: The stroke events diagnosed with MRI and CT examination in patients with no overt symptoms or sign of neurological deficit.
- Cardioembolism is responsible for around 20% of all ischemic strokes. Stroke caused by heart disease is primarily due to embolism of thrombotic material forming on the atrial or ventricular wall or the left heart valves. These thrombi then detach and embolism into the arterial circulation. The thrombus may fragment or lyse quickly, producing only a TIA. Alternatively, the arterial occlusion may last longer, producing stroke. Embolic strokes tend to be sudden in onset, with maximum neurologic deficit at once. With reperfusion following more prolonged ischemia, petechial hemorrhage can occur within the ischemic territory. This is usually of no clinical significance and should be distinguished from frank intracranial hemorrhage into a region of ischemic stroke where the mass effect from the hemorrhage can cause a decline in neurologic function.
- Emboli from the heart most often lodge in the MCA, the posterior cerebral artery (PCA), or one of their branches; infrequently, the anterior cerebral artery (ACA) territory is involved. Emboli large enough to occlude the stem of the MCA (3–4 mm) lead to large infarcts that involve both deep gray and white matter and some portions of the cortical surface and its underlying white matter. A smaller embolus may occlude a small cortical or penetrating arterial branch. The location and size of an infarct within a vascular territory depend on the extent of the collateral circulation.

OBSERVATIONS & RESULTS

Severity of Mitral Stenosis

Among these patients, 26 (17.2%) had mild MS, 47 (31.1%) had moderate MS while 78 (51.7%) had severe MS.

Table-1: Severity of Mitral Stenosis

Severity of Mitral Stenosis	Female	Male	Total
Mild (MVA 1.5-2.0 cm ²)	18 15.9%	8 21.1%	26 17.2%
Moderate (MVA 1.0-1.5 cm ²)	38 33.6%	9 23.7%	47 31.1%
Severe (MVA <1.0 cm ²)	57 50.4%	21 55.3%	78 51.7%
Total	113 100.0%	38 100.0%	151 100.0%




Fig-1: Severity of mitral stenosis and event free survival

Patients of severe mitral stenosis remained event free for mean period of 4.15 years (SD \pm 1.47), while patients of moderate MS remained event free for mean period of 4.43 years (SD \pm 1.7). Patients with mild MS were event free for mean period of 5.6 years (SD \pm 1.58).

DISCUSSION

Mitral stenosis is a sequel to Rheumatic fever which follows GAS Pharyngitis. The initial mitral valvulitis, as part of endocarditis occurring during rheumatic fever, results in cicatrization and fibrosis during healing process results in commissural fusion and disease of sub-valvular apparatus causing obstruction to left ventricular inflow [7].

The recurrent GAS pharyngitis leads to play important role in its causation. Mitral Stenosis is a progressive disease, if rheumatic fever recurrences are not prevented. Its natural history consists of progressive obstruction to left atrial emptying resulting in progressive pulmonary hypertension and its resultant complications.

Recurrent hospitalization for heart failure, development of atrial fibrillation, systemic embolization and cerebral embolization are associated with high morbidity and mortality seen during natural history of Mitral Stenosis.

The study was done to assess the natural history of mitral stenosis in non-operated cases. Severe Mitral stenosis is associated with high mortality and morbidity, if not operated/ intervened in suitable cases. We retrospectively analyzed case records data of patients with Mitral Stenosis registered in cardiology OPD form 1992 to 2013.

Although there were 771 such cases, we could only analyze 151 of the records as other records were not complete. We however, supplemented information

on complication and details by contacting patients or their families. Our study has a selection bias introduced by completeness of record.

CONCLUSION

The Rheumatic Mitral Stenosis affect young people particularly females. Unoperated patients with significant Mitral Stenosis carry high risk of Cardio embolism and recurrent hospitalization with associated high morbidity and mortality.

REFERENCES

1. Bonow R W, Blasé CA, Chatterjee K, de Leon AC, Faxon DP, Freed MD, ACC/AHA 2006 guidelines for the management of patient with valvular heart disease : A report of the American college of cardiology / American Heart Association Task force on practice guidelines (writing committee to revise the 1998 Guideline for the management of patient with valvular heart disease) developed in collaboration with the society of cardiovascular Anesthesiologist Endorsed by the Society for cardiovascular Angiography and Interventions and the society of thoracic surgeons. *Circulation* 2006; 114: e84-e231.
2. Nishimura RA, Carabello BA, Faxon DP, Freed MD, Lytle BW, O’Gara PT, O’Rourke RA, Shah PM. ACC/AHA 2008 Guideline update on vascular heart disease: focused update on infective endocarditis: a report of the American college of Cardiology/American Heart Association Task for practice guidelines endorsed by the society of cardiovascular Anesthesiologists, Society for cardiovascular angiography and Interventions, and Society of thoracic surgeons. *J Am Coll Cardiol* 2008; 52:676-85.
3. Vahanian A, Baumgartner H, Bax J, Butchart E, Dion R, Filippatos G, Flachskampf F, Hall R, Jung B, Kasprzak J, Nataf P. Guidelines on the management of valvular heart disease: The Task

Force on the Management of Valvular Heart Disease of the European Society of Cardiology. European heart journal. 2007 Jan 1;28(2):230-68.

4. Salem DN, O'Gara PT, Madias C, Pauker SG. Valvular and structural Heart Disease: American College of Chest Physicians Evidence Based Clinical Practice Guidelines (8th Edition) Chest 2008;133:593S-629S.
5. Tripathy A, Jeemon P, Ajay V, Prabhakar D, Reddy K. CVD profile of India, IC Health and WHO 2007: APW SE/06/226543
6. Feigin V, Lawes C, Benner D, Barker C, Parag V. Worldwide stroke incidence and early case fatality: a systematic review. Lancet Neurology 2009; 8(4): 355-369
7. Krasuski RA, Assar MD, Wang A, Kisslo KB, Pierce C, Harrison JK, Bashore TM. Usefulness of percutaneous balloon mitral commissurotomy in preventing the development of atrial fibrillation in patients with mitral stenosis. American Journal of Cardiology. 2004 Apr 1;93(7):936-9.