

## Study of Socio-Economic & Clinical Profile of Fibroids Undergoing Surgical Treatment in a Tertiary Care Hospital

Dr. S.A. Shinde<sup>1</sup>, Dr. U.S. Shinde<sup>1\*</sup>, Dr. G.S. Aher<sup>2</sup>

<sup>1</sup>Associate Professor, Department of Obstetrics & Gynecology, DVVPPF's Medical College Hospital, Ahmednagar, India

<sup>2</sup>Professor & HOD, Department of Obstetrics & Gynecology, DVVPPF's Medical College Hospital, Ahmednagar, India

### Original Research Article

#### \*Corresponding author

Dr. U.S. Shinde

#### Article History

Received: 10.06.2018

Accepted: 19.06.2018

Published: 30.06.2018

#### DOI:

10.21276/sjams.2018.6.6.31



**Abstract:** Uterine leiomyomas (fibroids) are the commonest benign tumors in women. Leiomyomas are the most common tumors in women of reproductive age. Objective of this study is to assess socio-economic & clinical profile of fibroid cases undergoing surgical treatment. This cross sectional prospective study was conducted at Department of Obstetrics and Gynecology in DVVPPF's medical college hospital, Ahmednagar, Maharashtra, over a period 4 years from June 2013 to may 2017. In this study 400 cases of uterine fibroid who were admitted & fulfilling inclusion criteria were included. The incidence of fibroid uterus was 12.34 % of all gynaecological admission. Most cases were seen in 41-45 years age group (47%), followed by 28% in 36-40 years age group. Parity  $\geq 3$  accounted for 83.25 % of all cases. Majority of cases were from lower socioeconomic class (53%), followed by lower middle class (29.75%). Most common presenting symptom was menorrhagia 45.25 %, followed by dysmenorrhoea (18 %). Most of the patients were having interstitial type of fibroid (66.5%). Uterine size was between 10-14 weeks (40.75%), followed by 6-10 weeks size (33.5%). Total abdominal hysterectomy with bilateral salpingo oophorectomy (TAH with BSO) was the most common surgical procedure done (37.75%). Post operative pyrexia was the most common complication seen post operatively (7%), followed by urinary tract infections (6.25%). Wound dehiscence was seen in 5 % of all cases. Uterine fibroid are commonly seen in women of reproductive age group and commonly present with menstrual disorders and may be associated with pelvic pain, infertility, pressure symptoms. The treatment is hysterectomy and myomectomy.

**Keywords:** Uterine Leiomyomata, Menstrual disorder, Hysterectomy, myomectomy.

### INTRODUCTION

Uterine leiomyomas (fibroids) are the commonest benign tumors in women [1]. Leiomyomas are the most common tumors in women of reproductive age [2]. They are most commonly multiple & asymptomatic. Several etiological factors have been suggested, with subtle variations in clinical presentation. The incidence of uterine fibroid tumours increases as woman grows older and they may occur in more than 30 percent of woman between 40 to 60 years of age [3]. It is benign, monoclonal tumour of the smooth muscle cells of the myometrium. The incidence is about 30% to 70% according to different diagnostic methods [4,5].

Uterine fibroids are common in nulliparous or relatively infertile women, African population & it is often associated with excessive oestrogen stimulation.

Fibroids may occur in any of the three layers of the uterus, could be intramural, submucosal and serosal. They can occur at unusual locations such as

the uterine cervix or within the layers of the broad ligament [6].

Woman with fibroids can be asymptomatic or may present with menorrhagia, pelvic pain with or without dysmenorrhoea or pressure symptoms, subfertility and recurrent pregnancy loss [7]. Diagnosis of fibroid is mainly clinical & can be confirmed by ultrasonography [8, 9]. Management of fibroid includes expectant management, medical management, myomectomy, and hysterectomy, embolotherapy depending on age, parity, location & presentation.

Objective of this study is to assess socio-economic & clinical profile of fibroid.

### MATERIALS & METHODS

This was a cross sectional prospective study, carried out in Obstetrics and Gynecology Department of DVVPPF's medical college & hospital, Ahmednagar, Maharashtra, over a period 4 years from June 2013 to may 2017. In this study 400 cases of uterine fibroid

who were admitted in this period were included after fulfillment of inclusion criteria.

**Inclusion criteria**

- Age 20-50 years
- Symtomatic patients
- Patients required operative procedure

**Exclusion criteria**

- Postmenopausal patients
- Asymptomatic patients

- Pregnancy

Each case was scrutinized for socio-economic, clinical profile, surgical procedure underwent and other necessary information. Data were collected in pre-designed form. Data were expressed as number & percentage in tabular form. Appropriate statistical analysis was done to interpret the results. Observations from this study were compared with other similar studies.

**RESULTS**

**Table-1: Incidence of fibroid uterus requiring operative procedure**

| Total gynaecological admissions | Admissions due to fibroid requiring operative procedure | Percentage |
|---------------------------------|---|------------|
| 3239                            | 400   | 12.34      |

During the study period total number of admission was 3239. Out of which 400 fibroid cases were managed surgically. The incidence of fibroid uterus requiring operative procedure was 12.34 % of all gynaecological admission.

Most cases were seen in 41-45 years age group (47%), followed by 28% in 36-40 years age group (Table-2).

**Table-2: Age wise distribution of cases**

| Age group(years) | Number of patients | Percentage |
|------------------|--------------------|------------|
| 20-25            | 11                 | 2.75       |
| 26-30            | 29                 | 7.25       |
| 31-35            | 10                 | 2.5        |
| 36-40            | 112                | 28         |
| 41-45            | 188                | 47         |
| 46-50            | 50                 | 12.5       |
| Total            | 400                | 100        |

**Table-3: Parity wise distribution of cases**

| Parity    | Number of patients | Percentage |
|-----------|--------------------|------------|
| Nullipara | 18                 | 4.5        |
| Para 1    | 20                 | 5          |
| Para 2    | 29                 | 7.25       |
| Para 3    | 159                | 39.75      |
| ≥ para 4  | 174                | 43.5       |
| Total     | 400                | 100        |

Majority of cases were having parity ≥3 i.e. 333 cases (83.25 %). Only 18 cases (4.5%) were nulliparous (Table-3).

Majority of cases were from lower socioeconomic class (53%), followed by lower middle class (29.75%) (Table-4).

**Table-4: Socioeconomic class wise distribution of cases**

| Socioeconomic class | Number of patients | Percentage |
|---------------------|--------------------|------------|
| Upper               | 11                 | 2.75       |
| Upper middle        | 13                 | 3.25       |
| Middle              | 45                 | 11.25      |
| Lower middle        | 119                | 29.75      |
| Lower               | 212                | 53         |
| Total               | 400                | 100        |

**Table-5: Distribution of cases according to clinical presentation**

| Clinical presentation       | Number of cases | Percentage |
|-----------------------------|-----------------|------------|
| Menorrhagia                 | 181             | 45.25      |
| Metrorrhagia                | 34              | 8.5        |
| Dysmenorrhoea               | 72              | 18         |
| Lump in abdomen             | 30              | 7.5        |
| Infertility                 | 20              | 5          |
| Pain /discomfort in abdomen | 96              | 24         |
| Pressure symptoms           | 26              | 6.5        |

Most common presenting symptom was menorrhagia (45.25 %), followed by dysmenorrhoea

(18 %).Lump in abdomen was seen in 7.5% of all cases. Infertility was seen in 5 % cases (Table-5).

**Table-6: Location of fibroid**

| Location of fibroid | Number of patients | Percentage |
|---------------------|--------------------|------------|
| Interstitial        | 266                | 66.5       |
| Submucous           | 61                 | 15.25      |
| Subserous           | 30                 | 7.5        |
| Cervical            | 5                  | 1.25       |
| Broad ligament      | 6                  | 1.5        |
| Fibroid polyp       | 11                 | 2.75       |
| Mixed               | 96                 | 24         |

Most of the patients were having interstitial type of fibroid (66.5%).Submucous & subserous fibroids were seen in 15.25% & 7.5% cases

respectively.11 cases were due to cervical & broad ligament fibroid.

**Table-7: Distribution of cases according to uterine size**

| Size of fibroid (weeks) | Number of patients | Percentage |
|-------------------------|--------------------|------------|
| 6-10                    | 134                | 33.5       |
| >10-14                  | 163                | 40.75      |
| >14-18                  | 62                 | 15.5       |
| >18-24                  | 28                 | 7          |
| >24                     | 13                 | 3.25       |
| Total                   | 400                | 100        |

Most of the patients were having uterine size between 10-14 weeks (40.75%), followed by 6-10

weeks size (33.5%).10.25% cases were having fibroid size more than 18 weeks.

**Table-8: Modality of surgical treatment given**

| Operation  | Number of cases | Percentage |
|--|-----------------|------------|
| Total Laparoscopic Hysterectomy(TLH)                     | 52              | 13         |
| Laparoscopic Assisted Vaginal Hysterectomy (LAVH)        | 18              | 4.5        |
| Total Abdominal Hysterectomy (TAH)                       | 112             | 28         |
| Total Abdominal Hysterectomy with Salpingo- oophorectomy | 151             | 37.75      |
| Subtotal hysterectomy                                    | 2               | 0.5        |
| Vaginal hysterectomy                                     | 34              | 8.5        |
| Myomectomy   | 20              | 5          |
| Polypectomy  | 11              | 2.75       |
| Total  | 400             | 100        |

Total abdominal hysterectomy with salpingo-oophorectomy was the most common surgical procedure done (37.75%), followed by Total

abdominal hysterectomy (TAH) done in 28 % cases.17.5 % cases underwent laparoscopic procedure.20 cases (5%) required myomectomy.

**Table-9: Postoperative complications**

| Complication  | Number of cases | Percentage |
|---|-----------------|------------|
| Pyrexia   | 28              | 7          |
| UTI (Urinary Tract Infections)                      | 25              | 6.25       |
| Paralytic ileus                                     | 12              | 3          |
| Wound dehiscence                                    | 20              | 5          |
| Blood transfusion due to intra-operative blood loss | 14              | 3.5        |
| Blood transfusion reaction                          | 4               | 1          |
| Post dural puncture headache                        | 4               | 1          |
| Others  | 12              | 3          |

Post-operative pyrexia was the most common complication seen post operatively (7%), followed by urinary tract infections (6.25%). Wound dehiscence was seen in 5 % of all cases. 14 cases (3.5%) required blood transfusion post-operatively due to intra operative blood loss. Blood transfusion reactions were seen in 4 cases.

## DISCUSSION

During the study period total number of admissions was 3239. Out of which 400 cases were of fibroid managed surgically. The incidence of fibroid uterus was 12.34 % of all gynaecological admission. This incidence is more than that of Verma *et al.* (6.8%) [10], Okogbo *et al.* (9.3%) [11], Rinku Das *et al.* (9.3%) [12].

In present study, most cases were seen in 41-50 years age group (59.5%), followed by 31-40 years age group (30.5%). Higher incidence in these age groups was also noted in other studies. Verma *et al.* [10] reported incidence of 42.12 % in 31-40 years & 37.7 % in 41-50 years age group. Rinku Das *et al.* [12] reported incidence of 18 % & 40 % in 30-39 years & 40-49 years age groups respectively. So from all these studies it is evident that incidence of fibroid is higher in third & fourth decade of life.

In our study majority of cases were having parity  $\geq 2$  (90.5 %). These findings were consistent with that of seen in study of Verma *et al.* [10], where 88.9 % cases were having parity  $\geq 2$ .

In this study, most common presenting symptom was menorrhagia (45.25 %), followed by dysmenorrhoea (18 %). Lump in abdomen was seen in 7.5% of all cases. Infertility was seen in 5 % cases. Similar findings were documented by Okogbo *et al.* [11], where menorrhagia, dysmenorrhoea & metrorrhagia accounted for 30.6 %, 18.9% & 17.1% cases respectively. Verma *et al.* [10] also reports menorrhagia as a most common presenting symptom seen in 60.4 % of all cases.

Most of the patients in present study were having interstitial type of fibroid (66.5%). Submucous & subserous fibroid were seen in 15.25% & 7.5% cases respectively. 11 cases were due to cervical &

broad ligament fibroids. Similar pattern is seen in study of Rinku Das *et al.* [12] who have reported incidence of 62.4%, 18.4 %, 7.6% in interstitial, submucous & subserous fibroids respectively.

Most of the patients in this study were having uterine size between 10-14 weeks (40.75%), followed by 6-10 weeks size (33.5%). 10.25% cases were having fibroid size more than 18 weeks. Verma *et al.* [10] also report similar findings regarding uterine size. In that study 36.9% & 35.2 % cases were having uterine size between 6-10 weeks & 10-14 weeks respectively. Okogbo *et al.* [11] reported maximum uterine size between 13-20 weeks in 74.4 % of cases.

Total abdominal hysterectomy with salpingo-oophorectomy was the most common surgical procedure done (37.75%), followed by Total abdominal hysterectomy (TAH) done in 28 % cases. 17.5 % cases underwent laparoscopic procedure. 20 cases (5%) required myomectomy. These findings were different from study of Verma *et al.* [10], where total abdominal hysterectomy with bilateral salpingo-oophorectomy was the most common surgery done in 94.8 % of all cases. Verma *et al.* [10] reported myomectomy done in 4.3 % of all cases, which is consistent with our study. Findings in our study were very much dissimilar with study of Okogbo *et al.* [11], where myomectomy was the common operation performed (65.41%). Rinku Das *et al.* [12] reported operation rates of 64 %, 24% & 12% for TAH, myomectomy & LAVH respectively.

Post-operative pyrexia was the most common complication seen post operatively (7%) in our study, followed by urinary tract infections (6.25%). Wound dehiscence was seen in 5 % of all cases. 14 cases (3.5%) required blood transfusion postoperatively due to intra operative blood loss. Blood transfusion reactions were seen in 4 cases. Rinku Das *et al.* [12] documented higher incidence of post operative complications compared with present study, where post operative pyrexia was seen in 12 % , followed by wound infection (10%), intraoperative blood loss requiring blood transfusion (8%), prolonged hospital stay (6%). Okogbo *et al.* [11] reports complications like postoperative pyrexia (13.5%), blood loss warranting transfusion (12.8%), postoperative anaemia (10.4%),

wound infection (8.7%), vault infection (5.1%) and prolonged hospital stay (4.2%). Other complications recorded were wound dehiscence (2.5%), vesicovaginal fistula (0.3%), ureteric injuries (0.3%), bladder injuries (0.6%), and death (0.2%).

College Journal, May 2014, Volume 13, Issue 2, Pages 36-41.

## CONCLUSION

Uterine fibroid is commonest benign tumour in female body, mostly asymptomatic. Uterine fibroid are commonly seen in women of reproductive age group. Menstrual complaints are the leading presenting symptoms. The treatment is usually hysterectomy. Myomectomy may be considered choice when family is not complete.

## REFERENCES

1. Wise LA, Palmer JR, Stewart EA, Rosenberg L. Age-specific incidence rates for self-reported uterine leiomyomata in the black women's health study. *Obstet Gynecol.* 2005; 105:563-8.
2. Stewart EA. Uterine fibroids. *Lancet.* 2001; 357:293-8.
3. Manor C, Brunsell MS. Uterine fibroid tumors: Diagnosis and treatment. *Am FAM Physician.* 2007; 75(10):1503-1508.
4. Goodwin SC, Spices JB, Worthington Kirsch R, Peterson E, Pron G. Uterine artery embolization for treatment of leiomyomata: Long term outcomes from the Fibroid Registry. *Obstet. Gynecol.* 2008.111(1): 22-33.
5. Cramer SF, Patel A. The frequency of uterine leiomyomas. *Am J Clin Pathol.* 1990; 94:435-8.
6. Meniru GI, Wasdahl D, Onuora CO. Vaginal leiomyoma co-existing with broad ligament and multiple uterine leiomyomas. *Arch Gynecol Obstet.* 2001; 265: 105-107
7. Lumsden M A, Wallace E M. Clinical presentation of uterine fibroids. *Baillieres Clin Obstet Gyne.* 1998;12(2): 177-195
8. Anate M. Uterine fibroids in Federal medical centre, Lokoja: a five year review 2002-2006. *The Nigerian Clinical Review Journal.* 2007: Jan/Feb: 5-12.
9. Ogunniyi SO, Fasubaa OB. Uterine Fibromyoma in Ilesha, Nigeria. *Nigerian Medical Practitioner.* 1990; 19(6): 93-95.
10. Verma Madhurima, Gupta Ritu, Porwal Sanjay, Swarnkar Madhusudan & Porwal Varsha. Uterine leiomyomata a clinical study. *J Pharm Biomed Sci.* 2013. March; 28 (28):672-676.
11. Okogbo FO, Ezechi O, Loto O, Ezeobi P. Uterine Leiomyomata in South Western Nigeria: a clinical study of presentations and management outcome. *African Health Sciences.* 2011; 11(2):271-278.
12. Rinku Das, Rummana Jafrin, Banita Biswas. Clinical Presentation & Operative Treatment of Leiomyoma of Uterus and its Outcome. *Chattagram Maa-O-Shishu Hospital Medical*