

Spectrum of Primary Paediatric Brain Tumors –A Single Institutional Experience

Dr. Sanidhaya Tak¹, Dr. Shweta Choudhary^{2*}¹Resident, Department of Pulmonary Medicine, S.N Medical College, Jodhpur, Rajasthan, India²Assistant Professor, Department of Pathology, S.N Medical College, Jodhpur, Rajasthan, India

Original Research Article

***Corresponding author**

Dr. Shweta Choudhary

Article History

Received: 02.07.2018

Accepted: 11.07.2018

Published: 30.07.2018

DOI:

10.21276/sjams.2018.6.7.19



Abstract: Tumors of the central nervous system are the second most common childhood tumor after haematological malignancies [1]. These are the leading cause of cancer-related deaths in children and constitute approximately 35% of all childhood malignancies [2]. The present study analyses the age, sex, location, and histological spectrum of primary paediatric brain tumors in a tertiary care hospital of western India. This is a retrospective study which included 120 primary paediatric brain tumors (<15 yrs of age) in two year duration at tertiary care centre of western India. All the cases of CNS tumors were studied by histopathological examination using H and E staining. Immunohistochemistry was done as and when required. Most of the patients were in the age group of 11-15 yrs. male to female ratio was 1.3:1. Most of the paediatric brain tumors were infratentorial in location. The most common primary paediatric brain tumors were astrocytic tumors (41.6%), followed by medulloblastoma and craniopharyngiomas. The most common astrocytic tumor was pilocytic astrocytoma. There is increase in the incidence of paediatric CNS tumors but a few studies on primary paediatric CNS tumors have been reported in india . The current study is from single largest institution in western India. As compared to western data, incidence of brain tumors in children was found to be less in the present study.

Keywords: paediatric, central nervous system ,tumors.

INTRODUCTION

Tumors of the central nervous system (CNS) are the second most common childhood tumor after haematological malignancies[1]. Tumors of the nervous system are the leading cause of cancer-related deaths in children and constitutes approximately 35% of all childhood malignancies [2]. Paediatric CNS tumors differ significantly from adult brain tumors with respect to sites of origin, clinical presentation, histological features, tendency for early dissemination and prognosis. The predominant CNS tumors in adults are glial neoplasms, meningiomas and metastases, whereas in children, besides gliomas, other major tumors include embryonal neoplasms .The present study therefore analyses the histological spectrum of primary paediatric brain tumors in a single largest tertiary care hospital of western india.

MATERIALS AND METHODS

A hospital based retrospective study on biopsy specimens received at tertiary care centre of western

India in two year. A total of 120 biopsy specimens of paediatric brain tumors (<15 yrs of age) were reviewed. Infectious, metastatic lesions and vascular malformations were excluded .The specimens were processed routinely and IHC performed wherever necessary. Results were expressed as percentage and proportions.

RESULTS & DISCUSSION

Paediatric CNS tumors accounted on an average 10.56% of total intracranial tumors. Male to female ratio was 1.3:1. Most of the paediatric brain tumors were infratentorial(60%) in location. Astrocytomas(41.6%) were the commonest tumors followed by medulloblastomas (26.6 %). Craniopharyngiomas (10%)were the third most common tumors followed by and oligodendroglioma(6.6%), ependymal tumors (5%), nerve sheath tumors (3.3%) germ cell tumor(3.3%) and meningeal tumors (3.3%).{Table 1}. The most common astrocytic tumor was pilocytic astrocytoma.

Table-1: Spectrum of primary paediatric CNS Tumors

Pediatric Cns Tumor	No of cases	Percentage (%)
Astocytoma	50	41.6
Ependymoma	6	5
Oligodendroglioma	8	6.66
Medulloblastoma	32	26.66
Cranioparngioma	12	10
Meningioma	4	3.33
Nerve sheath tumor	4	3.33
Germ cell tumor	4	3.33
total	120	

Pediatric CNS tumors form the most concerning group of tumors due to high incidence and mortality. There is rise in incidence of CNS tumors in developing countries with increased availability of diagnostic and healthcare facilities. The exact tumor burden of CNS tumors is underestimated due to lack of complete registration of newly diagnosed cases with cancer registries. Hospital-based prevalence studies therefore are required for estimating the disease burden, management, prognosis and for assessing geographical differences in their molecular and genetic profiles. In contrast to the Western literature, there are few data on pediatric CNS tumors in India. Pediatric CNS tumors accounted for 10.6 % of total intracranial tumors in the present study. It was higher in the study by Jain A *et al.* (14.8%) [4]. As compared to western data, incidence of brain tumors in children was found to be less in the present study.

Majority were in infratentorial location in our study (60%). This was in concordance with studies by Nasir S *et al.* (64.1%) and Hanif G *et al.* and Shah SH *et al.* (43.5%) [9,5,11]. Whereas Asirvatham JR *et al.* concluded that pediatric CNS tumors were present in supratentorial compartment more often than the infratentorial compartment, in Rickert *et al.* study, supratentorial-to-infratentorial ratio was 1.1 [7,3].

Male to female ratio was 1.3:1 in the present study. It was similar to studies done by Jain A *et al.* (1.6:1), Shah SH *et al.* (1.1:1), Hanif G *et al.* 3:1), Ahmed N *et al.* (2.5:1), Asirvatham JR *et al.* (1.7:1), Das U *et al.* (1.58:1). whereas in Rickert CH *et al.* study male to female ratio was 1:1. Most of the patients were in the age group of 11-15 yrs [4, 11,5,7,8].

The most common primary pediatric brain tumors were astrocytic tumors (41.6%), followed by medulloblastoma (26.6%) and craniopharyngiomas (10%) in the present study.

This was in concordance with the study by Chen *et al.* who found astrocytomas to be leading tumors (29.2%). The most common primary pediatric brain tumors were astrocytic tumors in Jain A *et al.* (34.7%), Shah SH *et al.* (39%), Hanif G *et al.* (75.8%), Asirvatham JR *et al.* (47.3%), Rickert CH *et al.*

(33.6%) studies [12,4,11,5,7,4]. While medulloblastoma was most common in Nasir S *et al.* (33.6%), Ahmed N *et al.* (49.4%), Das U *et al.* studies and ependymoma in Azad TD *et al.* study (17.5%) [9,6,8,10].

CONCLUSION

The present study revealed the histopathological spectrum of primary pediatric CNS tumors in children at largest hospital in western India. There is a rising incidence of pediatric CNS tumors and hence more comprehensive population-based studies are required to determine the cancer burden, management in developing countries.

REFERENCES

- Rosemberg S, Fujiwara D. Epidemiology of pediatric tumors of the nervous system according to the WHO 2000 classification: a report of 1,195 cases from a single institution. *Child's Nervous System*. 2005 Nov 1;21(11):940-4.
- Jemal A, Siegel R, Ward E, Murray T, Xu J, Smigal C, Thun MJ. Cancer statistics, 2006. *CA: a cancer journal for clinicians*. 2006 Mar 1;56(2):106-30.
- Rickert CH, Probst-Cousin S, Gullotta F. Primary intracranial neoplasms of infancy and early childhood. *Child's nervous system*. 1997 Oct 1;13(10):507-13.
- Jain A, Sharma MC, Suri V, Kale SS, Mahapatra AK, Tatke M, Chacko G, Pathak A, Santosh V, Nair P, Husain N. Spectrum of pediatric brain tumors in India: A multi-institutional study. *Neurology India*. 2011 Mar 1;59(2):208.
- Hanif G, Shafqat S. Morphological pattern and frequency of intracranial tumours in children. *Journal of the College of Physicians and Surgeons-Pakistan: JCPSP*. 2004 Mar;14(3):150-2.
- Ahmed N, Bhurgri Y, Sadiq S, Shakoork KA. Pediatric brain tumours at a tertiary care hospital in Karachi. *Asian Pacific Journal of Cancer Prevention*. 2007 Sep;8(3):399.
- Asirvatham JR, Deepti AN, Chyne R, Prasad MS, Chacko AG, Rajshekhar V, Chacko G. Pediatric tumors of the central nervous system: a retrospective study of 1,043 cases from a tertiary care center in South India. *Child's Nervous System*. 2011 Aug 1;27(8):1257-63.

8. Das U, Appaji L, Kumari BA, Sirsath NT, Padma M, Kavitha S, Avinash T, Lakshmaiah KC. Spectrum of pediatric brain tumors: a report of 341 cases from a tertiary cancer center in India. *The Indian Journal of Pediatrics*. 2014 Oct 1;81(10):1089-91.
9. Nasir S, Jamila B, Khaleeq S. A retrospective study of primary brain tumors in children under 14 years of age at PIMS, Islamabad. *Asian Pac J Cancer Prev*. 2010 Jan 1;11(5):1225-7.
10. Azad TD, Shrestha RK, Vaca S, Niyaf A, Pradhananga A, Sedain G, Sharma MR, Shilpakar SK, Grant GA. Pediatric central nervous system tumors in Nepal: Retrospective analysis and literature review of low-and middle-income countries. *World neurosurgery*. 2015 Dec 1;84(6):1832-7.
11. Shah SH, Soomro IN, Hussainy AS, Hassan SH. Clinico-morphological pattern of intracranial tumors in children. *JPMA. The Journal of the Pakistan Medical Association*. 1999 Mar;49(3):63-5.
12. Chen L, Zou X, Wang Y, Mao Y, Zhou L. Central nervous system tumors: a single center pathology review of 34,140 cases over 60 years. *BMC clinical pathology*. 2013 Dec;13(1):14.